

Phosphorus speciation in Swedish agricultural clay soils

Influence of fertilisation and mineralogy

Ann Kristin Eriksson

*Faculty of Natural Resources and Agricultural Sciences
Department of Soil and Environment
Uppsala*

Doctoral Thesis
Swedish University of Agricultural Sciences
Uppsala 2016

Acta Universitatis agriculturae Sueciae

2016:25

Cover: Soil after plowing. The word cloud contains the words most frequently occurring in the abstract of this thesis. Also shown are the bulk soil XRD patterns of the Fors soil.

(photo: A.K. Eriksson)

ISSN 1652-6880

ISBN (print version) 978-91-576-8552-0

ISBN (electronic version) 978-91-576-8553-7

© 2016 Ann Kristin Eriksson, Uppsala

Print: SLU Service/Repro, Uppsala 2016

Phosphorus speciation in Swedish agricultural clay soils. Influence of fertilisation and mineralogy

Abstract

Phosphorus (P) is an important element for crop production, but build-up of excess soil P can promote P leaching and eutrophication of surface waters. To better understand the dynamics of P release from soil to waters, more knowledge is needed about sorption patterns and P speciation in agricultural soils.

Two new indices were developed to assess the importance of P sorption to hydroxy-interlayered clay minerals, and to evaluate the amount of hydroxy-interlayering and hydroxy-interlayer stability. A strong relationship was found between oxalate-extractable aluminium (Al) and the amount of hydroxy-interlayering in soil, suggesting a common source. This makes it difficult to analytically distinguish between phosphate (PO_4) adsorbed to hydroxy-interlayers from PO_4 adsorbed to Al hydroxide-type precipitates.

Application of X-ray absorption near-edge structure (XANES) spectroscopy to evaluate P speciation in soil profiles and the effects of P depletion and fertilisation revealed a distinct change in P speciation with increasing soil depth in an agricultural clay soil profile. The results indicated that the subsoil P predominantly occurred as apatite, whereas PO_4 adsorbed with Al-hydroxides or hydroxy-interlayers dominated in the topsoil. Organic P and PO_4 adsorbed to iron (Fe)-(hydr)oxides were observed only in the topsoil. This can be explained by long-term weathering of apatite and silicates, particularly ferromagnesian forms such as amphibole, in the upper soil horizons, causing an association of the released PO_4 with secondary metal (hydr)oxides.

Collectively, the XANES results showed that the most important phases governing P retention and release in agricultural clay soils are adsorption of PO_4 to Al and to Fe in (hydr)oxide minerals or interlayers. After fertilisation, P was adsorbed to Al hydroxide phases in six different soil types studied, but in two of the soils there was also an increase in calcium phosphate. Moreover, P solubility was shown to be lowest at pH values ranging from 4.5 to 7.5 and increased with decreasing pH, probably as a result of the dissolution of apatite and PO_4 -bearing hydroxy-Al precipitates at low pH.

Keywords: Soils, XANES spectroscopy, phosphorus, phosphate, clays, apatite, secondary iron and aluminium (hydr)oxides, clay mineralogy, hydroxy-interlayered minerals, X-ray diffraction

Author's address: Ann Kristin Eriksson, Swedish University of Agricultural Sciences (SLU), Department of Soil and Environment, P.O. Box 7014, 750 07 Uppsala, Sweden
E-mail: ann.kristin.eriksson@slu.se

Dedication

To my grandmother, Hildur

“Environments are not just containers, but are processes that change the content totally”

Marshall McLuhan

so...

“What we plant in the soil of contemplation, we shall reap in the harvest of action”

Meister Eckhart

Contents

List of Publications	7
Abbreviations	9
1 Introduction	11
2 Aims and hypotheses	13
3 Background	15
3.1 Phosphorus in soils	15
3.1.1 Phosphorus in minerals	15
3.1.2 Adsorbed phosphate	16
3.1.3 Organic P forms	17
3.2 Speciation of P in soils	17
3.3 The pH dependence of PO ₄ sorption/desorption in soils	18
3.4 Clay mineralogy and effect on P sorption	19
3.5 Research questions	20
4 Materials and Methods	21
4.1 Site description	21
4.1.1 Long-term fertility experiments	21
4.1.2 Lanna	21
4.1.3 Bornsjön	22
4.1.4 Lilla Böslid	22
4.1.5 20E	22
4.2 Soil sampling and preparation	23
4.3 General soil characterisation	24
4.4 Mineralogy	24
4.4.1 Bulk soil mineralogy	24
4.4.2 Clay mineralogy	25
4.4.3 Characterisation of hydroxy-interlayered minerals	25
4.5 Speciation of phosphorus	29
4.5.1 Clay fractions	29
4.5.2 Bulk soils	30
4.6 Solubility experiments	31
4.7 Sorption experiments	33

5	Results	35
5.1	Mineralogy (Paper I)	35
	5.1.1 Bulk soil mineralogy	35
	5.1.2 Clay mineralogy	36
	5.1.3 A new index to characterise hydroxy-interlayered minerals	36
5.2	The effect of weathering and soil development, an example from Lanna (Paper III)	40
	5.2.1 Mineralogy affected by weathering	40
	5.2.2 Phosphorus speciation affected by soil development	41
	5.2.3 Phosphorus solubility patterns affected by soil development	42
5.3	Phosphorus speciation affected by long-term fertilisation (Paper II and IV)	43
	5.3.1 Phosphorus speciation in soils and clay fractions with no added phosphorus	43
	5.3.2 Phosphorus speciation in soils and clay fractions after long-term fertilisation	44
5.4	Dissolved phosphate as affected by long-term fertilisation (Paper IV)	45
5.5	Phosphate sorption affected by long-term fertilisation (Paper IV)	49
6	Discussion	51
6.1	Characterising hydroxy-interlayering in clay fractions and soils (Paper I)	51
6.2	How does soil development affect P speciation? (Papers I and III)	52
	6.2.1 Formation of hydroxy-interlayers and (hydr)oxides	52
	6.2.2 Weathering of apatite	53
	6.2.3 Organic P and PO ₄ adsorbed to Fe (hydr)oxides in the topsoil	54
6.3	Phosphorus speciation, which P is available? (Papers II, III and IV)	54
6.4	Is it possible to develop a standard procedure for XANES data treatment?	55
6.5	Evaluation of hypotheses	57
7	Conclusions	59
8	Implications and future perspectives	61
	References	63
	Acknowledgements	71

List of Publications

This thesis is based on the work contained in the following papers, referred to by Roman numerals in the text:

- I Eriksson, A.K., Hillier, S., Simonsson, M. and Gustafsson J.P. X-ray diffraction using *in situ* heating to characterise hydroxy-interlayering in clay minerals: A new method evaluated on Swedish agricultural soils (manuscript).
- II Eriksson, A.K., Hesterberg, D. and Gustafsson, J.P. (2015). Phosphorus speciation of clay fractions from long-term fertility experiments in Sweden. *Geoderma* 241-242, 68-74. [doi:10.1016/j.geoderma.2014.10.023](https://doi.org/10.1016/j.geoderma.2014.10.023)
- III Eriksson, A.K., Hillier, S., Hesterberg, D., Klysubun, W., Ulén, B. and Gustafsson J.P. Evolution of phosphorus speciation with depth in an agricultural soil profile (submitted to *Geoderma*).
- IV Eriksson, A.K., Hesterberg, D., Klysubun, W. and Gustafsson, J.P. Phosphorus dynamics in Swedish agricultural soils as influenced by fertilisation and mineralogical properties: insights gained from batch experiments and XANES spectroscopy (manuscript).

Paper II is reproduced with the permission of the publishers.

My contribution to paper I-IV included in this thesis was as follows:

Planned the experimental work together with the co-authors. Performed the practical laboratory work with assistance from co-authors. Performed data analysis and writing, with assistance from the co-authors.

Abbreviations

Al	Aluminium
BNL	Brookhaven National Laboratory
BL	Beamline
Ca	Calcium
COG	Centre of gravity
DOM	Dissolved organic matter
e.s.d.	Equivalent spherical diameter
Fe	Iron
Ha	Hectare
HIV	Hydroxy-interlayered vermiculite
K	Potassium
LCF	Linear Combination Fitting
Mg	Magnesium
N	Nitrogen
NH ₄	Ammonium
NMR	Nuclear Magnetic Resonance spectroscopy
NSLS	National Synchrotron Light Source
P	Phosphorus
P-HCl	Phosphorus digested in HCl
P-AL	Phosphorus extracted in ammonium lactate and acetic acid
PCA	Principal Component Analysis
PO ₄	Orthophosphate
PO ₄ -P	Orthophosphate phosphorus
PsTotP	Pseudo-total phosphorus
RIR	Reference Intensity Ratio
SLRI	Synchrotron Light Research Institute
XANES	X-ray Adsorption Near-edge Structure
XRD	X-ray Diffraction
Yr	Year

1 Introduction

Phosphorus (P) is a macronutrient and an important element for crop production. Too little P may limit crop yield, whereas too much may promote P leaching and contribute to eutrophication of surface waters. Chemical fertilisers are a primary source of P in agricultural soils, but application of animal wastes to soil also contributes to build-up of P in soils. Apatite is commonly mined for use in fertiliser, but it is a non-renewable resource. Current reserves may be depleted in 50-100 years (Cordell *et al.*, 2009). In general, soil P management in crop production must be responsive both to crop needs and water quality protection.

The largest terrestrial source of P entering surface waters from Sweden is agricultural fields (Boesch *et al.*, 2006). Phosphorus in soil water occurs as inorganic orthophosphate (PO_4) ions (*e.g.* H_2PO_4^- , HPO_4^{2-}), as dissolved organic P or as inorganic or organic P bound to suspended colloids. Leachate from clay soils commonly contains a larger fraction of particulate P than leachate from coarse-textured soils (*e.g.* Djodjic *et al.*, 2004).

The plant availability of P is generally related to soil P solubility. The processes affecting soil P availability to the crop are not all fully understood. According to the established paradigm of soil P availability, P is most available to plants at neutral pH (Brady & Weil, 1999). However, many recent studies have shown that this paradigm does not hold for clay-rich soils and that in fact the P solubility in laboratory experiments is *lowest* at neutral pH (*e.g.* Gustafsson *et al.*, 2012; Devau *et al.*, 2011; Weng *et al.*, 2011). It has been suggested that this solubility behaviour may be due to the instability of hydroxy-aluminium (Al) polymers of clay mineral interlayers at low pH (Gustafsson *et al.*, 2012), but this has not been confirmed. Furthermore, the roles of particle size distribution and fertilisation history in P availability are not clear (Gustafsson *et al.*, 2012).

Different sequential chemical extraction methods have commonly been used for P speciation (*e.g.* Hedley *et al.*, 1982; Williams *et al.*, 1967; Chang & Jackson, 1957). The current limited knowledge of P speciation in soils (despite many years of research on the topic) can be attributed to the lack of methods that permit direct identification of the various P-containing phases. However, in recent years Nuclear Magnetic Resonance (NMR) spectroscopy and X-ray Adsorption Near-edge Structure (XANES) spectroscopy have emerged as powerful techniques that allow more specific information regarding P speciation to be obtained (McLaughlin *et al.*, 2011). Correct characterisation of P speciation in soils is a challenge, not least due to the presence of many different P phases, such as organic P, calcium (Ca) and Al phosphates, and PO₄ adsorbed to iron (Fe) and Al (hydr)oxides. Nevertheless, for process-oriented modelling of PO₄ sorption/desorption in soils, it is essential to know which processes dominate. Another prerequisite for successful modelling is that the P dynamics of each individual P-containing phase must be sufficiently well known (*e.g.* Devau *et al.*, 2011; Weng *et al.*, 2011), and any interactions between phases in these multi-phase systems must be understood.

Clay mineralogy is important for studies of soil P chemistry for at least two reasons: (i) There may be a direct role of hydroxy-interlayered minerals in P sorption (as suggested by Gustafsson *et al.*, 2012); and (ii) variations in clay mineralogy in a pedon or among soils may be linked to a number of soil-forming processes that affect P chemistry (*e.g.* pH, weathering, formation of secondary solid phases *etc.*). However, clay mineral identification and quantification are subject to a number of analytical difficulties. One property that is particularly difficult to determine accurately and consistently from one soil to another is the amount of hydroxy-Al-interlayering.

2 Aims and hypotheses

The overall aim of this thesis was to investigate the sorption patterns and speciation of P in agricultural clay soils, in order to better understand the dynamics of P release from soils to waters (see review in Figure 1).

Specific objectives of the work were to:

1. Develop a new method to characterise hydroxy-interlayered clay minerals semi-quantitatively with regard to the amount and stability of interlayers.
2. Investigate P speciation in a clay-rich soil profile as influenced by mineralogical composition and soil-forming processes.
3. Evaluate changes in P speciation caused by long-term P fertilisation for a range of typical Swedish agricultural soils.
4. Study the pH dependence of sorption/desorption patterns (for the same soils).

The hypotheses tested were:

1. Soils with a large proportion of hydroxy-Al-interlayered clay minerals have higher P sorption capacity than those with a small proportion.
2.
 - a) In clay soils with low P status, P is mainly sorbed to Al hydroxide phases;
 - b) In clay soils with high P status, P is bound to both Al and Fe (hydr)oxide phases;
 - c) Ca-bound P as apatite is only reactive in calcareous soils.
3. Apatite increases with depth in the soil profile as a result of its dissolution in the upper horizons during pedogenic weathering.

4. The solubility of phosphate (PO_4) in Swedish agricultural clay soils is lowest at near neutral pH and increases with both increasing and decreasing pH.




Figure 1. Illustration of how the different studies described in Papers I-IV in this thesis relate to each other (photo and drawing by A.K. Eriksson).

3 Background

3.1 Phosphorus in soils

Phosphorus in soils is present both in the solid phase of the soil matrix and in the soil solution. In the solid phase, P can be present as: (1) a constituent of primary and secondary minerals; (2) surface complexes on oxides [mainly Al- and Fe (hydr)oxides] and on clay minerals; and (3) organic forms. Adsorption/desorption is considered the single most important process controlling soil P solubilisation during the season. However, precipitation/dissolution of minerals and immobilisation/mineralisation of organic forms are also important (Pierzynski *et al.*, 2005). According to the conventional paradigm (*e.g.* Brady & Weil, 1999), at low soil pH PO_4 is adsorbed to Al- and Fe (hydr)oxides or precipitated as Al and/or Fe phosphates, whereas at higher pH PO_4 is mostly fixed as Ca phosphates. This paradigm states that PO_4 has its greatest solubility at neutral soil pH (pH 6-7), where solubility is defined as the concentration of soluble P after a given time. An overview of the different P forms and reactions in soils is provided in Figure 2.

3.1.1 Phosphorus in minerals

Phosphorus is present in both primary and secondary minerals. The weathering of primary minerals, and subsequent release of P, is usually a very slow process. Some examples of P-containing primary minerals are variscite, strengite and apatite. Secondary minerals occur predominantly as poorly crystalline precipitates, for example less crystalline Al, Fe and Ca phosphate minerals. The secondary mineral phases usually have a larger surface area than the primary minerals, and hence they are more reactive and more easily dissolved (Pierzynski *et al.*, 2005).




Figure 2. Pools and reactions of P in the soil system (modified from Pierzynski *et al.*, 2005; photo A.K. Eriksson).

3.1.2 Adsorbed phosphate

Phosphate is commonly adsorbed to Al and Fe (hydr)oxides, *e.g.* goethite, ferrihydrite, gibbsite and boehmite (Pierzynski *et al.*, 2005). Previous research indicates that amorphous Al-hydroxide and aluminosilicate phases are among the most important P-bearing particles (Pierzynski *et al.*, 1990) and may also control the solubility of inorganic P in soils (*e.g.* Cui & Weng, 2013).

A number of factors affect PO_4 adsorption, for example; (1) the variable charge properties of adsorption sites, which cause decreased PO_4 adsorption when the pH is increased (*e.g.* Antelo *et al.*, 2010; Hansen *et al.*, 1999); (2) the reaction time, as a prolonged reaction time will increase the quantity of adsorbed PO_4 (Hansen *et al.*, 1999); (3) competition for adsorption sites by *e.g.* organic acids (Oburger *et al.*, 2011), carbonate (Sø *et al.*, 2011) and, arsenate (Gustafsson *et al.*, 2012); and (4) the ionic strength, with increased ionic strength decreasing PO_4 adsorption at low pH, but increasing adsorption at high pH (Barrow, 2008; Barrow *et al.*, 1980).

The adsorption process is usually divided into two different types of reactions, one fast and one slow. Fast reactions dominate in the early stage (during the first minute) and are reversible. Slow reactions dominate in the later stages and are partly reversible (depending on the time scale for desorption). The slow reactions are probably due to slow diffusion of PO_4 into

the interior of Fe and Al (hydr)oxide precipitates (Strauss *et al.*, 1997; Willett *et al.*, 1988), but the mechanisms are not yet completely understood.

3.1.3 Organic P forms

Between 15 and 80 % of the P in the surface horizon of soils is usually present in organic forms, mainly originating from plant residues. Inositol phosphates (phytates), sugar phosphates, phospholipids and nucleic acids are the most frequently found organic P species. However, less than 50 % of the organic P in soil can be characterised into known compounds. Organic P is mainly found in the topsoil and decreases with soil depth (Stevenson, 1994).

3.2 Speciation of P in soils

Traditional techniques for determining P speciation in soils involve various extraction procedures, *e.g.* sequential extractions (*e.g.* Hedley *et al.*, 1982; Williams *et al.*, 1967; Chang & Jackson, 1957). However, extraction results are only indicative (at best), as they provide only indirect information about soil P speciation. It is probably impossible to design extraction procedures that target specific P forms.

A number of challenges are involved in measuring P in soils and especially P speciation (McLaughlin *et al.*, 2011). The heterogeneity of the soil material, with many different surfaces and minerals, gives a high matrix complexity, resulting in many different P species. For example, it has been shown that a particular form of extraction can release different kinds of P species in different soils (Kar *et al.*, 2011).

More recently, NMR (Leinweber *et al.*, 1997) and XANES (*e.g.* Beauchemin *et al.*, 2003; Hesterberg *et al.*, 1999) spectroscopy have been used. These techniques potentially represent a vast improvement over extractions, as they permit more direct identification of the species involved. With the XANES technique, it is possible to estimate the dominant P species in the soil by means of linear combination fitting (LCF), in which the spectra of known standards are compared against the spectrum of a specific soil sample (Kelly *et al.*, 2008).

The principle employed in XANES spectroscopy is that a sample is exposed to an X-ray beam and scanned across an energy range corresponding to the binding energy of core electrons (*e.g.* K-shell electrons) in atoms of the element of interest. A synchrotron X-ray beam of high intensity is used in XANES spectroscopy of soils to increase the analytical sensitivity. The high X-ray intensity causes excitation of the core electrons into the continuum and a core hole is formed in the orbital that the electron occupied. However, this is

an unstable state for the atom, and therefore one electron from a higher energy level drops down to the core hole, releasing fluorescence X-rays (or secondary electrons). The emitted fluorescence can be measured by a sensitive fluorescence detector, and the intensity of this radiation is proportional to the amount of absorbed X-rays, *i.e.* the energy-dependent absorption coefficient. The energy level at which K-shell electrons are excited corresponds to the so-called K edge of the element. For phosphorus, the K edge ranges between 2145 eV for elemental P and 2153 eV for P(V).

In a typical K-edge XANES experiment on soil P, the soil sample is exposed to the X-ray beam, the energy of which is varied from a level below the edge (usually about 2100 eV) to an energy level above the edge (in this case usually 2320 eV). The amount of absorbed energy is measured continuously as the fluorescence signal. The exact binding energy of the electron and variations in signals due to back-scattering of the excited electrons by surrounding atoms above the edge are affected by the oxidation state and molecular coordination environments of P atoms. Average absorption coefficients vary depending on the local molecular structure of the P species present, and hence different combinations of species will give rise to different spectra (Kelly *et al.*, 2008).

Franke & Hormes (1995) identified the properties giving rise to the different features of the K-edge XANES spectra for P. They found that phosphate associated with transition metal cations (*e.g.* Mn or Fe) shows a pre-edge resonance, while alkaline, alkaline earth and condensed phosphates (P bound to Ca, K or Na) show a wider peak base. They also found a high intensity of the so-called white line (the main absorption edge) in samples that contained ionic and covalently bound phosphates and a less intense white line for metal-bound phosphates.

McLaughlin *et al.* (2011) rated XANES spectroscopy as one of the best methods for direct speciation of inorganic P. However, the method offers very limited information on organic P species due to lack of sensitivity. Negassa & Leinweber (2009) suggested a combination of NMR and XANES spectroscopy for full information about P speciation. Kruse *et al.* (2009) also suggested the use of L_{2,3}-edge XANES spectroscopy, which seems to give more information about organic P species.

3.3 The pH dependence of PO₄ sorption/desorption in soils

As mentioned above, PO₄ is usually considered to be most readily dissolved at around neutral pH. However, a number of recent studies (*e.g.* Gustafsson *et al.*, 2012; Devau *et al.*, 2011; Weng *et al.*, 2011) have shown the opposite trend for

clay soils, with the lowest P solubility near neutral pH. There are different explanations for these observations, *e.g.* (1) instability of hydroxy-Al polymers of clay mineral interlayers at low pH (Gustafsson *et al.*, 2012); (2) interaction between Ca^{2+} and metal oxides leading to increased P sorption, especially at high pH (Weng *et al.*, 2011); (3) adsorption of Al^{3+} to organic material at lower pH, which removes P-adsorbing Al hydroxides or hydroxy-Al-interlayers (Gustafsson *et al.*, 2012; Weng *et al.*, 2011); and (4) competition between phosphate and organic matter for sorption surfaces.

However, it seems that other factors also have an effect. After long-term fertilisation, Gustafsson *et al.* (2012) found that the soluble $\text{PO}_4\text{-P}$ increased. They also found that the solubilisation behaviour differed depending on whether the P was freshly added in the laboratory or in long-term experiments in the field, as a result of occlusion processes. Moreover, the P solubility was found to increase with decreasing pH for soils with > 20 % clay and to decrease with increasing pH for soils with < 10 % clay (Gustafsson *et al.*, 2012).

3.4 Clay mineralogy and effect on P sorption

Previous research has indicated that clay mineralogy is rather uniform on regional scale in Sweden. Illite is commonly the dominant mineral phase in the clay fraction, followed by vermiculite and chlorite (Wiklander, 1950). However, the county of Skåne in southern Sweden is an exception, as smectite-rich and kaolinite-rich soils are also frequently found in this area (Ahlberg *et al.*, 2003). Most clay fractions present in Swedish agricultural soils also contain large amounts of primary minerals, *e.g.* quartz (Kirchmann *et al.*, 2005; Stevens & Bayard, 1994; Wiklander, 1950).

The following hypotheses have been forwarded to explain the observed similarities on regional scale (Stevens & Bayard, 1994): (1) most clay soils originate from glacial and postglacial deposits; (2) the landscape formed after the glacial period was not eroded down to the bedrock, and therefore there is a limiting effect of the underlying rock type as a parent material; (3) long-range transport of minerals (including clay minerals) took place with ice-melt water; (4) the ice milled down pre-glacial exposed bedrock further north from which fine-grained primary mineral particles were produced and transported.

Penn *et al.* (2005) found that soils containing hydroxy-interlayered vermiculite (HIV) also commonly showed stronger PO_4 sorption. Hydroxy-interlayered vermiculite can be considered an intermediate between pure vermiculite and aluminium chlorite. It is commonly formed by deposition of hydroxy-Al polymers in the interlayer area of vermiculite or as a weathering

product of Al chlorite (Barnhisel & Bertsch, 1989). However, the presence of carbonate inhibits the weathering of clay minerals and the formation of these hydroxy-interlayers (Stevens & Bayard, 1994).

3.5 Research questions

It has been suggested that sorption of PO_4 onto hydroxy-interlayered clay minerals can explain the observed lowest P solubility at near neutral pH (Gustafsson *et al.*, 2012). Moreover, Penn *et al.* (2005) observed that soils containing hydroxy-interlayered clay minerals commonly had higher P sorption capacity than other soils. In order to obtain more knowledge about the effect of mineralogy, more information is needed. In this thesis XRD was used for this purpose, in collaboration with the James Hutton Institute, which has a well-established method for quantifying mineralogy and clay mineralogy (Omotoso *et al.*, 2006; Hillier, 2003; Hillier, 1999).

To the best of my knowledge, there is no simple way to estimate the proportion and stability of hydroxy-interlayered minerals. Heating is one way to quantify the amount of interlayering (Harris *et al.*, 1992). In this approach, XRD is used with an *in situ* heating stage that also prevents rehydration, which has otherwise been observed to affect quantification (Sayegh *et al.*, 1965). To estimate the influence of these clay minerals on the PO_4 sorption properties of the whole soil, an index to characterise the hydroxy-interlayering of the whole soil is needed. For comparison, minerals with different amounts of interlayering need to be synthesised in the laboratory. The results can then be compared with those for the clay fraction of soils.

To understand the differences in the pH-dependent sorption/desorption patterns of P in agricultural clay soils (Gustafsson *et al.*, 2012), more knowledge is needed about the P speciation in these soils. The XANES method seems to be the most prominent approach used to date for such analyses of inorganic phosphates. Using information about P speciation and mineralogy, pH-dependent P solubility and PO_4 adsorption can be studied. Gustafsson *et al.* (2012) showed for one soil that fertilisation may affect the pH-dependent solubility behaviour, which is further investigated in the present thesis.

Higher P content has been reported in the subsoil of agricultural soil profiles, but the origin of this P and how it may affect P leaching is not explained (*e.g.* Andersson *et al.*, 2015; Andersson *et al.*, 2013). To investigate this issue, in this thesis the P speciation in a soil profile was determined by XANES spectroscopy and the influence of mineralogical composition on P speciation was analysed by XRD.

4 Materials and Methods

4.1 Site description

4.1.1 Long-term fertility experiments

The Swedish long-term soil fertility experiments were initiated in the period 1957-1969 and comprise different fertilisation treatments and crop rotations (Carlgren & Mattsson, 2001). The samples used in this thesis were from six sites in southern and central Sweden, namely Fors, Kungsängen, Vreta Kloster, Bjertorp, Ekebo and Fjärdingslöv (Table 1). Soil was sampled from three different P and potassium (K) fertiliser application treatments: 'A', where no P and K were added; 'C', where P and K were applied to replace the amounts removed with the previous year's harvest plus a surplus of 20 kg P and 50 kg K ha⁻¹ yr⁻¹ (except for the Ekebo and Fjärdingslöv sites, where the surplus was 15 kg P and 40 kg K ha⁻¹ yr⁻¹); and 'D', where P and K were applied to replace the amounts removed with the previous year's harvest plus a surplus of 30 kg P and 80 kg K ha⁻¹. All samples were taken from a non-livestock crop rotation with the highest N fertiliser application rate ('3'), 125 kg N ha⁻¹ yr⁻¹ (150 N ha⁻¹ yr⁻¹ at Ekebo and Fjärdingslöv). Selected treatments in the soil fertility experiments have been investigated previously in the context of P studies, e.g. regarding speciation of organic P (Ahlgren *et al.*, 2013) and P leaching (e.g. Djodjic *et al.*, 2004; Svanbäck *et al.*, 2013). Temporal changes in P status in these experiments have also been investigated (Bergström *et al.*, 2015).

4.1.2 Lanna

Lanna is an experimental farm on the largest agricultural plain (Vara plain) in south-western Sweden (Table 1). The experimental farm was started in 1929 and the location was chosen to represent the conditions usually found on the Vara plain (Johansson, 1944). The soil profile shows an increase in P content with depth (Andersson *et al.*, 2015; Andersson *et al.*, 2013;

Mattson *et al.*, 1950) and is silty clay in the topsoil and clay in the subsoil. Both horizons have a strong coarse subangular blocky structure (Bergström *et al.*, 1994). The samples used in this thesis were collected in conjunction with the studies by Andersson *et al.* (2013) and Andersson *et al.* (2015). Previous investigations on P at this site have addressed the effects of liming on soil P (Mattson *et al.*, 1950), the influence on P losses of different soil management practices and cropping systems (Aronsson *et al.*, 2011; Neumann *et al.*, 2011), and the effects on P leaching of the chemical composition of the subsoil (Andersson *et al.*, 2015; Andersson *et al.*, 2013).

4.1.3 Borsjön

Borsjön is an experimental field situated in central Sweden (Table 1), on a soil of marine origin (Ulén *et al.*, 2014). The field is situated in a catchment area where research concerning P losses and P chemistry is carried out. The field experiment was established in 2006 to study the effect of different mitigation options for nutrient (primarily P) and pesticide leaching (Svanbäck *et al.*, 2014). Soil samples used in this thesis were taken from the topsoil in four plots with conventional tillage and conventional P fertilisation. The P added by fertilisation corresponded to the expected crop requirement in the current year. Previous studies at the site have examined the importance of subsoil conditions for P losses (Andersson *et al.*, 2015; Andersson *et al.*, 2013) and P retention in constructed wetlands (Kynkäänniemi *et al.*, 2013).

4.1.4 Lilla Böslid

Lilla Böslid is an experimental farm situated in south-western Sweden (Table 1). It is divided into two areas with experimental leaching plots. The experimental activities were initiated in 2002 (Aronsson *et al.*, 2011). The area contains both sandy and clayey soils. The samples used in this thesis originated from a clay loam soil. The leaching experiments were started in 2010, when mineral fertiliser was applied (supplying 21 kg P ha⁻¹ yr⁻¹).

4.1.5 20E

The 20E field is included in the Swedish environmental monitoring programme of single observation fields (Table 1). It belongs to a farm with pig production situated in south-eastern Sweden. The soil is managed according to conventional practices by the commercial farmer (Ulén *et al.*, 2012).

Table 1. *Coordinates and soil classification of the field experiments at different sites in Sweden from which samples were taken for this thesis. n.d.= not classified*

Site	Coordinates	Soil taxonomy	Soil type
Fors	60°20'N, 17°29'E	Calcaric Phaeozem ¹	Silt loam
Kungsängen	59°50'N, 17°40'E	Gleyic Cambisol ¹	Clay
Vreta Kloster	58°30'N, 15°30'E	Haplic Phaeozem ²	Silty clay
Bjertorp	58°14'N, 13°08'E	<i>n.d.</i>	Silty clay loam
Ekebo	55°59'N, 12°52'E	Haplic Phaeozem ³	Loam
Fjärdingslöv	55°24'N, 13°14'E	Haplic Phaeozem ³	Sandy loam
Lanna	58°21'N, 13°07'E	Udertic Haploboroll ⁴	Silty clay / clay
Bornsjön	59°14'N, 17°41'E	Eutric Cambisol ⁵	Silty clay
Lilla Böslid	56°35'N, 12°56'E	<i>n.d.</i>	Clay loam
20E	<i>n.d.</i>	<i>n.d.</i>	Clay

¹Kirchmann (1991)

²Kirchmann *et al.* (2005)

³Kirchmann *et al.* (1999)

⁴Bergström *et al.* (1994)

⁵Ulén & Persson, (1999)

4.2 Soil sampling and preparation

Samples of topsoil (0-20 cm) were collected using a core sampler (approx. 2 cm diameter) within a circle of 1 m in diameter and combined into 1-2 kg bulk samples. At the long-term fertility experiment sites, all samples were taken at least 0.5 m from the edge of the plots. The samples from the Lanna soil profile were collected from five different depths (0-10, 10-30, 30-50, 50-70 and 70-100 cm) at five different points along a linear transect of approximately 10 m. The five samples from each depth were mixed to form one bulk sample with approximately the same amount of soil from each sampling point. The samples were air-dried (approx. 30 °C), sieved (2 mm) and stored under dry conditions until analysis.

Clay fractions (< 2 µm in equivalent spherical diameter (e.s.d.)) were separated from the soil samples by sedimentation according to Stokes' law. In brief, subsamples of 20 g soil were mixed with 200 mL water and treated twice with ultrasonic dispersion for 5 minutes. The suspension was stirred thoroughly with a spatula between the ultrasound treatments. The suspension was then transferred to a 1-L cylinder, mixed thoroughly and left to sediment for 16 h before the top 20 cm was siphoned off. The water level was then refilled and the sedimentation was repeated. The clay fraction suspension was freeze-dried and stored under dry conditions. The clay fraction samples used for oxalate

extraction and for K and magnesium (Mg) saturation were separated in the same way, but dried at approx. 30-40 °C.

4.3 General soil characterisation

The particle size distribution of soil samples was analysed by sieving and sedimentation analysis according to ISO 11277 (2009). A summary of the soil chemical characterisation methods used is provided in Table 2. Soil pH was measured using a Radiometer PHM93 pH meter with a GK2401C combination pH electrode in a suspension containing 6 g air-dried soil and 18 mL de-ionised water (1:3 soil:water). Organic carbon (OrgC) content was determined after combustion using a LECO CNS-2000 analyser (LECO, St Joseph, MI). Aluminium, Fe and P were extracted in ammonium oxalate according to van Reeuwijk (1995). The extract was filtered through a 0.2 µm single-use filter and diluted 1:5 in ultra-pure H₂O. The concentrations of Al and Fe in the extracts were analysed using ICP-OES on an ICP Optima 7300 DV instrument. The extractant is assumed to dissolve Fe phases with low crystallinity and surface-reactive Al, such as Al in organic complexes, allophane, imogolite, Al hydroxide and other forms of non- and paracrystalline inorganic Al. In this thesis, no effort was made to differentiate non-crystalline Al(OH)₃ from non- and paracrystalline Al silicates, such as allophane and imogolite; all are referred to as 'Al hydroxide'. The Al and Fe were also extracted in pyrophosphate where the extract was filtered through a 0.2 µm single use filter and diluted 1:5 in H₂O. The pyrophosphate extract was analysed using ICP-OES on a Perkin Elmer 5300 DV instrument. The extraction is assumed to dissolve organically bound Fe and Al, although it may also to some extent dissolve non- and paracrystalline inorganic phases (Kizewski *et al.*, 2011; Kaiser & Zech, 1996). A summary of these methods is provided in Table 2.

4.4 Mineralogy

4.4.1 Bulk soil mineralogy

Air-dried soil samples (< 2 mm) were micronised in ethanol (10 mL ethanol to 3 g air-dried soil) in a McCrone mill. The suspension was formed into a random powder by spray-drying (Hillier, 1999). The powder was packed into a metal holder and diffraction patterns were recorded with a Philips Xpert Pro diffractometer using Ni-filtered Cu K α radiation. The XRD patterns were recorded from $2\theta = 3^\circ$ to 70° by scanning in steps of 0.0167° and counting for 300 s per step. The recorded patterns were quantitatively analysed by full pattern fitting, as described in Omotoso *et al.* (2006). The group of clay

Table 2. *Methods used for general soil characterisation in this thesis*

Method	Element	Soil (mg)	Solution (mL)	Solution specification	Reaction time	Measure -ment	Reference
P-AL	P	5	100	0.1 M NH ₄ lactate, 0.4 M acetic acid (pH 3.75)	1.5 h	ICP	Egnér <i>et al.</i> (1960)
P-HCl	P	2	50	2 M HCl	2 h	ICP	KLS (1965)
Oxalate	P, Al, Fe	1	100	0.2 M NH ₄ oxalate (pH 3.0)	4 h ¹⁾	ICP ²⁾	van Reeuwijk (1995)
Pyro- phosphate	Al, Fe	1	100	0.1 M Na ₂ P ₂ O ₇ ·10H ₂ O	16 h	ICP	
Pseudo- total P	Al, Fe	3	30	<i>Aqua regia</i> 3.95 M HNO ₃ , 9 M HCl	2 h	ICP	ISO 11466 (1995)

¹⁾ In darkness

²⁾ PO₄ analysed colorimetrically according to Wolf & Baker (1990)

minerals used in this method included vermiculite, hydrobiotite, chlorite, illite, kaolinite and various types of mica/smectite mixed-layer clay minerals.

4.4.2 Clay mineralogy

Mounts of orientated clay fractions (<2 µm e.s.d.; see section 4.2) were prepared on glass slides using the filter transfer method (Moore & Reynolds, 1997). The XRD patterns were recorded on a Siemens D5000 diffractometer using Fe-filtered Co K α radiation after air-drying, glycolation (ethylene glycol vapour solvation at 60 °C) and heating (300 °C for 1 h on a hotplate). Diffraction patterns were recorded from $2\theta = 3^\circ$ to 45° by scanning in steps of 0.02° and counting for 1 s per step. The XRD patterns were analysed semi-quantitatively using a peak area-based reference intensity ratio (RIR) method (see Table 3) (Hillier, 2003). All measurements were made using the EVA DIFFRAC^{plus} software (Bruker, 2005) and calculations based on RIRs were calculated in NEWMOD[®] (Reynolds, 1985).

4.4.3 Characterisation of hydroxy-interlayered minerals

All methods used to characterise the amount of hydroxy-interlayering rely on measurements of the decrease of the *d*-spacing (Å) with different treatments at the laboratory. Two existing methods for determining the amount of hydroxy-interlayering were used as references; both involve K- and Mg-saturation (Esser, 1990; Matsue & Wada, 1988). In this thesis work, two new methods were developed. The first is a simple method ranking the COG values of the clay fractions after K-saturation and heating.

Table 3. Reference intensity ratio (RIR) values used for semi-quantitative analysis of clay minerals

Clay mineral	Peak position (\AA)	Peak order	RIR	Used trace
Kaolinite	7.10	001	2.45	Air-dried
	3.58	002	1.70	
	2.38	003	0.12	
Chlorite	14.20	001	0.59	Air-dried
	7.10	002	3.32	
	4.72	003	0.81	
	3.52	004	1.77	
Illite	10.00	001	1.00	Glycolated
Expandable minerals	10.00	001	1.00	Heated

Clay fractions from soils and synthesised hydroxy-interlayered minerals were investigated by all described methods. The synthesised hydroxy-interlayered minerals were used to verify the amount of hydroxy-interlayering. In the second method, an *in situ* heating stage was used to assess the amount and stability of the hydroxy-interlayers. To evaluate these methods both vermiculite with synthesised hydroxy-interlayers and clay fractions from soils were investigated.

Preparation of synthetic hydroxy-interlayered vermiculite

Hydroxy-interlayered vermiculite can be synthesised from pure vermiculite using one of three different approaches: (1) The Al hydroxide polymer is prepared in a separate solution, which is mixed with the clay mineral to incorporate the aluminium polymer into the interlayer space (*e.g.* Carstea, 1968; Slaughter & Milne, 1960). (2) The clay mineral is saturated with an easily exchangeable cation, followed by ion exchange of aluminium ions and titration with sodium hydroxide to form polymers in the interlayer space (*e.g.* Janssen *et al.*, 1997; Rich, 1960; Sawhney, 1960). (3) The clay mineral is artificially weathered (using *e.g.* acid in a reaction chamber), resulting in the formation of hydroxy-interlayers from weathered Al (Mareschal *et al.*, 2009).

The second method was used because it was deemed more likely that the Al^{3+} ion is able to enter the interlayer space to form aluminium polymers than for the formed aluminium polymers to diffuse into the interlayer space, both because of charge and size. The third method may lead to the formation of hydroxy-interlayers that not only contain aluminium polymers.

Synthetic hydroxy-interlayered minerals were prepared from pure trioctahedral vermiculite (<20 μm e.s.d; sample KL2 in the paper by Marwa *et al.*, 2009). First, vermiculite was sodium-saturated, and afterwards it was

loaded with two different levels of hydroxy-interlayers (0.032 and 0.67 mol Al kg⁻¹ vermiculite, respectively) according to the titration method described by Janssen *et al.* (1997).

K and Mg saturation

To detect hydroxy-interlayering in a sample, different pre-treatment steps were performed before XRD analysis. The samples were K- and Mg-saturated by adding 20 mL 1 M KCl or MgCl₂ to approximately 100 mg clay and equilibrating overnight. In the morning, the suspension was filtered (0.45 µm) and rinsed twice with the same amount of the saturating solution. The sample was then washed three times with 20 mL deionised H₂O and the clay fraction was mounted and analysed as described in section 4.4.2.

In situ heating treatment

The XRD patterns were recorded upon *in situ* heating using an Anton Paar XRK900 reaction chamber from 25 to 300 °C in 25 °C steps and from 300 to 550 °C in 50 °C steps.

Method 1: Change in COG upon K saturation (using the Mg-saturated sample as a reference)

This method was developed by Esser (1990) and quantifies the shift in COG (x-scale, *d*-distance (Å)) after K-saturation compared with an Mg-saturated reference sample (measured at room temperature). Instead of RIR values, a value of 0 is set at 9.5 Å, increasing to 1 at 15.5 Å. The calculation is as follows:

$$\Delta\bar{x}_{Mg-K} = \bar{x}_{Mg} - \bar{x}_K \quad (1)$$

where \bar{x}_{Mg} is the COG of the Mg-saturated sample (Å), \bar{x}_K is the COG of the K-saturated sample (Å), and $\Delta\bar{x}_{Mg-K}$ is the difference between these (Figure 3a).

Method 2: Intensity ratios of Mg- and K-saturated samples

This index was developed by Matsue and Wada (1988) and is defined as:

$$I_K/I_{Mg} = \frac{(I_{14}/(I_{14}+I_{10}))_K}{(I_{14}/(I_{14}+I_{10}))_{Mg}} \quad (2)$$

where I_{14} represents the height of the 14 Å peak, I_{10} is the height of the 10 Å peak, and the subscripts K and Mg denote the K-saturated and Mg-saturated

samples measured at room temperature, respectively. The ratio I_K/I_{Mg} describes the resistance of the 14 Å peak of an Mg-saturated sample to collapse upon saturation with K (Figure 3b).

Method 3: COG after K saturation and heating (300 °C)

This is a newly developed index, which is defined as the COG between 9.5 and 15.5 Å measured after K-saturation and heating to 300 °C on a hot plate (Figure 3c). This method was developed as a simple and quick method to characterise the amount of hydroxy-interlayering without any special equipment.

Method 4: Changes in COG upon in situ heating (250 to 450 °C)

This new method was developed to get more precise characterisation of the hydroxy-interlayers (both amount and stability).

This index was measured on the *in situ* heated samples. It is divided into two different parts, describing (4a) the amount of hydroxy-interlayering; and (4b) the thermal stability of the hydroxy-interlayers. The first part is defined as the COG between 8 Å (approx. 11° 2θ) and 25 Å (approx. 3.5° 2θ):

$$x_{amount} = \bar{x}_{450^{\circ}\text{C}} - 10.2\text{Å} \quad (3)$$

where x_{amount} represents the amount of hydroxy-interlayering, $\bar{x}_{450^{\circ}\text{C}}$ is the COG at 450 °C and 10.2 Å is the reference (COG) value for a completely collapsed 2:1 mineral.

The second part of the index describes the thermal stability of the hydroxy-interlayers and is defined as the change in COG, $\Delta\bar{x}_{temp.}$, when the sample is heated from 250 °C to 450 °C (Figure 3d):

$$\Delta\bar{x}_{temp.} = \bar{x}_{250^{\circ}\text{C}} - \bar{x}_{450^{\circ}\text{C}} \quad (4)$$




Figure 3. Measurements to determine the amount of hydroxy-interlayered minerals in clay fractions according to: a) Esser (1990), comparing COG after K saturation with Mg saturation as a reference; b) Matsue & Wada (1988), using an intensity ratio comparing the intensities of the 14 Å peak (I_{14}) and 10 Å peak (I_{10}); c) a new index measuring centre of gravity after K saturation and heating and d) another new index measuring centre of gravity after *in situ* heating at 250 and 450 °C. The horizontal (x) axis represents d -distance (Å), and the vertical axis the diffracted intensity.

4.5 Speciation of phosphorus

4.5.1 Clay fractions

Phosphorus K-edge XANES spectra of the clay fractions from the long-term fertility experiments were collected on Beamline X-15B at the National Synchrotron Light Source (NSLS), Brookhaven National Laboratory (BNL) in NY state, U.S. The data were collected using an Si(111) monochromator. Fluorescence signals were collected in a helium (He) atmosphere using a solid-state Ge-detector. To minimise the signal from silicon (Si) in the samples, the detector window was covered with an approx. 40 µm thick polypropylene film. A minimum of 13 and 9 scans was made for unfertilised and fertilised treatments, respectively. Baseline correction, normalisation and LCF analysis were carried out using the Athena Software (Athena v. 0.8.056 and Demeter v. 0.9.18; Ravel & Newville, 2005). Three sets of standards were used in the LCF analysis: (1) mineral samples from Ingall *et al.* (2011); (2) mineral standards, standards with adsorbed PO_4 and organic P standards from Hesterberg *et al.* (1999); and (3) mineral standards and standards with adsorbed phosphate from Eveborn *et al.* (2009). For the unfertilised treatments, a maximum of three

standards (selected as described in Paper II) was allowed in the fit. For the fertilised treatments, there was also a maximum of three standards, but the best-fit model spectrum generated for the unfertilised treatment was used as a mandatory standard which assumes that fertiliser may add new P species to the unfertilised soil P species.

4.5.2 Bulk soils

Phosphorus K-edge XANES spectra were collected at Beamline BL-8 at the Synchrotron Light Research Institute (SLRI) in Nakhon Ratchasima, Thailand, for the soil samples from the long-term soil fertility experiments and the bulk soil and clay fraction samples from the Lanna site. The data were collected using an InSb(111) crystal monochromator, and fluorescence signals were measured using a solid-state 13-element Ge detector. The sample was held in a He gas atmosphere. Baseline correction and normalisation were performed using the Athena software, version 0.9.020 (Ravel & Newville, 2005). LCF analysis was conducted using weighted combinations of 31 known standards (Figure 5), including amorphous calcium phosphate (ACP), octacalcium phosphate (OCP), three different natural apatites, synthetic hydroxyapatite, brushite, monetite, amorphous aluminium phosphate, phosphate adsorbed to aluminium hydroxide, phosphate adsorbed to gibbsite, variscite, amorphous iron phosphate, phosphate adsorbed to ferrihydrite, phosphate adsorbed to goethite, strengite, struvite, lecithin and phytate. In the topsoil sample of the unfertilized treatments, four standards were allowed in the fit, including lecithin and apatite (Taiba) as mandatory standards. For the fertilised treatments, a maximum of three standards was allowed, but the best-fit model spectrum generated for the unfertilised treatment was used as a mandatory standard to investigate the speciation of the P bound from added fertiliser.

The absolute fluorescence intensity, defined as the edge step in the XANES spectrum (Leri *et al.*, 2006), was compared to the amount of P as determined by different extractions (Figure 4). The most significant relationship ($r = 0.81^{***}$) was found between P digestible in HCl (KLS, 1965) and the absolute fluorescence intensity. In principle, the X-ray fluorescence is directly related to the total concentration of sample P; however, a lower correlation was found for the fluorescence signal with pseudo-total P (measured as digestion with *aqua regia*; $r = 0.69^{***}$) than with HCl digestion (Figure 4c).




Figure 4. Scatter plots for correlations between absolute fluorescence intensity and a) P extracted with ammonium lactate, b) P extracted with oxalate, c) P solubilised by HCl digestion, and d) P solubilised by *aqua regia* digestion from samples included in the long-term fertility experiments (treatments A3 and D3) and the Lanna soil profile.

4.1 Solubility experiments

Batch experiments were carried out to determine pH-dependent P solubility from the soil samples by adding acid or base to soil samples at a rate ranging from 0.075 mol acid kg⁻¹ soil to 0.03 mol base kg⁻¹. The suspensions were equilibrated for 7 d in darkness on an end-over-end shaker. After centrifugation, the pH was measured in the supernatant, which was then filtered (0.2 μ m) and analysed colorimetrically for molybdate reactive PO₄-P using a Tecator Aquatec 5400 spectrophotometer with flow injection analysis, based on the method by Murphy & Riley (1962). In addition, Ca, Al and Fe were determined using ICP-OES on an ICP Optima 7300 DV instrument.

To calculate the activity of free Ca²⁺ and PO₄³⁻, Visual MINTEQ ver. 3.1 software was used (Gustafsson, 2014). The solubility constants of various P-containing mineral phases used were those detailed by Gustafsson *et al.* (2012).




Figure 5. XANES spectra for the standards used for LCF analysis.

4.2 Sorption experiments

Batch sorption isotherm experiments were also performed on the soil samples, to which PO_4 was added in a range from 0 to $0.0045 \text{ mol kg}^{-1}$. Equilibration and measurements were performed as described in section 4.6 at natural soil pH. Afterwards, pH was measured in the supernatant and $\text{PO}_4\text{-P}$ was analysed colorimetrically (Seal Analytical AAS Autoanalyzer).

A Freundlich isotherm was fitted to the data:

$$n_{\text{sorb}} = K_F \cdot c^m - n_{\text{init}} \quad (5)$$

where n_{sorb} represents the amount of PO_4 sorbed during the experiment, n_{init} is the amount of PO_4 sorbed before the start of the experiment (mol kg^{-1}), K_F and m are coefficients that have to be optimised during the fitting process, and c is the equilibrium concentration of total dissolved P (mol L^{-1}). Equation 5 can be logarithmically transformed to:

$$\log n = \log K_F + m \cdot \log c \quad (6)$$

where $n = n_{\text{init}} + n_{\text{sorb}}$. This equation is linear, where m is the slope and $\log K_F$ the intercept. In this thesis, the value of m was set to 0.33 as suggested by Tolner & Füleky (1995). This was shown to give values of n_{init} comparable with isotopically exchanged P. Values of $\log K_F$ and $\log n$ were determined after adjusting n_{init} to the best fit as judged by the R^2 value.

5 Results

5.1 Mineralogy (Paper I)

5.1.1 Bulk soil mineralogy

The results of the quantitative bulk mineralogical analysis are shown in Table 4. A finding in common for all soil samples was that more than 40 % of the mineral soil consisted of quartz, plagioclase and K-feldspar. Quartz was the single most common mineral in all soils. Only one of the soils, Fors, had a substantial amount of calcite.

Table 4. *Bulk fine earth (<2 mm) topsoil mineralogy in soil samples (% of bulk soil) of Swedish field experiments, analysed by full XRD pattern fitting after micronising and spray drying*

	Quartz	Plagioclase	K-feldspar	Calcite	Amphibole	Iron oxides	Clay minerals ¹
	<i>(% of bulk soil)</i>						
Fors	30	18	14	6.4	3.4	2.0	15
Kungsängen	17	14	11	<i>n.d.</i> ²	2.9	3.8	34
Vreta Kloster	32	14	12	<i>n.d.</i> ²	2.2	2.6	23
Bjertorp	33	18	16	<i>n.d.</i> ²	2.6	2.7	15
Ekebo	48	14	14	<i>n.d.</i> ²	1.7	1.1	11
Bornsjön	18	15	12	<i>n.d.</i> ²	2.9	3.7	32
Lilla Böslid	24	21	17	<i>n.d.</i> ²	3.2	2.6	17
20E	21	13	11	<i>n.d.</i> ²	2.4	3.6	31

¹Including chlorite, muscovite, biotite, hydrobiotite, vermiculite, illite, mixed layer minerals with illite/smectite and kaolinite.

²Not detected.

5.1.2 Clay mineralogy

All soils had a high content of expandable minerals (> 50 %), especially the soil from Fjärdingslöv, for which a high content of smectite was observed. The clays also commonly had a high content of mica/illite (> 20 %), except for the sample from Fjärdingslöv. The soils had a low content of kaolinite (< 5 %) with the exception of Ekebo, which had a content of 20 % (Table 5).

Most of the samples showed an increase in d-spacing of the 14 Å peak after ethylene glycol solvation, which is an indication of low layer charge (Brindley & Brown, 1980). However, the d-spacing did not increase in Ekebo, Bjertorp and Lilla Böslid, soils that contained obvious hydroxy-interlayers.

5.1.3 A new index to characterise hydroxy-interlayered minerals

Clay mineral response to increased heating

As a starting point for evaluation of the heating effects, the response of synthetic hydroxy-interlayered vermiculite was investigated. Below 100 °C, the pure vermiculite had a d-spacing of 14.4 Å, consistent with two layers of water molecules in the interlayer. Between 100 and 200 °C this decreased to a d-spacing of 11.8 Å, corresponding to one layer of water molecules. Above 200 °C the peak collapsed to 10 Å, consistent with a completely dehydrated interlayer.

Table 5. *Semi-quantitative XRD analysis of parallel orientated clay mineral specimens prepared by the filter transfer method (% of clay fraction, <2 µm) from topsoil samples*

	Kaolinite	Chlorite	Mica	Expandable minerals ¹
	<i>(% of clay fraction)</i>			
Fors	2	3	44	51
Kungsängen	1	3	40	56
Vreta Kloster	4	1	23	73
Bjertorp	4	<i>n.d.</i> ²	27	69
Ekebo	20	<i>n.d.</i> ²	23	57
Fjärdingslöv	6	<i>n.d.</i> ²	8	86
Bornsjön	1	2	25	71
Lilla Böslid	4	3	36	60
20E	3	2	41	54

¹Including minerals that contract from 14 to 10 Å upon heating, *e.g.* smectite, vermiculite, hydroxy-interlayered minerals and mixed layer illite-smectite minerals.

²Not detected.

In contrast to the stepwise contraction of synthetic HIV, in natural clay fractions with hydroxy-interlayers there was a progressive movement of the peak towards 10 Å with increasing temperature. The collapse to 10.3-10.4 Å was not achieved until heating to 500 °C. In the remainder of this thesis, the resistance to collapse is described by the term 'stability'. As discussed by Tolpeshta *et al.* (2010), there is no single definition of the term stability, as it has been used to describe the response to a multitude of treatments, such as displacement by cations, added dissolving reagents, heating or ageing. Here, the term relates to thermal stability, and a certain degree of stability (*i.e.* resistance to collapse upon heating) was observed in all soils, indicating some hydroxy-interlayering.

There are different ways to describe the amount of hydroxy-interlayering in the soil: (1) Determining the amount of hydroxy-interlayers in a specific clay mineral; (2) determining the amount of hydroxy-interlayered mineral in the clay fraction (<2 µm e.s.d.); and (3) determining the amount of hydroxy-interlayering in the bulk soil. However, the mixture of minerals makes it impossible to identify the amount of hydroxy-interlayers in each specific mineral. In this thesis the amount of hydroxy-interlayering in the clay fraction (2) was estimated.

Methods 1 and 2: Prior indices based on Mg and K saturation

Methods 1 and 2 showed similarities when ranking the amount of hydroxy-interlayering. Both methods identified the Ekebo sample as that with the highest amount of hydroxy-interlayering, followed by Lilla Böslid and Bjertorp. The Vreta Kloster sample had the lowest amount of hydroxy-interlayering. The samples from Fors and Kungsängen also had low amount of hydroxy-interlayering. For the samples from Fjärdingslöv and 20E, the results differed depending on the method. A summary of the results obtained is presented in Table 6.

Method 3: COG after K saturation and heating

Method 3 was clearly correlated with Method 1 ($r = 0.81^{**}$), but not with Method 2. An even better match ($r = 0.96^*$) was found with Method 1 when only the samples containing obvious amounts of hydroxy-interlayers were considered (Ekebo, Lilla Böslid, Bjertorp and Fjärdingslöv).

Method 4: Change in COG upon in situ heating

This new method based on measurement of COG at 450 °C *in situ* heating showed a clear correlation with Method 3 ($r = 0.94^{***}$ for clay samples originating from soil). There was also a strong correlation with Method 1

($r = -0.78^*$) and Method 2 ($r = 0.84^{***}$). According to Method 4, all clay fractions had a higher amount of hydroxy-interlayering than the pure vermiculite. The samples were therefore grouped into: (1) clay fractions containing obvious amounts of hydroxy-interlayers (Ekebo, Lilla Böslid, Bjertorp and Fjärdingslöv); and (2) clay fractions with only small amounts of hydroxy-interlayers (Vreta Kloster, Fors, Kungsängen, 20E and Bornsjön) (Figure 6). A relationship between the amount of hydroxy-interlayering and soil pH was also found, with more hydroxy-interlayering at lower pH ($r = -0.75^*$ with the calcareous Fors soil with $\text{pH} > 7.7$ excluded). Thermostability was defined as the difference between the COG at 450 and 250 °C. For the clay fractions originating from the soil samples, a strong correlation was found between the stability and the amount of hydroxy-interlayering ($r = 0.97^{***}$). For synthetic samples with different amount of hydroxy-interlayering, there were no clear differences in stability between the samples. However, some soils with a lower amount of hydroxy-interlayering also had higher stability, which may be related to the proportions of hydroxy-interlayers that contract rather than to the properties of the hydroxy-polymers.

Table 6. *Determination of the amount of hydroxy-interlayering with different methods*

	Method 1: Esser (1990) ¹	Method 2: Matsue & Wada (1988) ²	Method 3: COG as K- sat and heated to 300 °C (Å)	Method 4a: Amount as COG at 450 °C ³ (Å)	Method 4b: Stability as change in COG ⁴ (Å)
HIV + 0 mol Al kg ⁻¹			11.24	0.15	0.15
HIV + 0.032 mol Al kg ⁻¹			11.43	0.93	0.59
HIV + 0.67 mol Al kg ⁻¹			11.93	1.52	0.67
Fors	22.9	0.35	10.51	0.23	0.09
Kungsängen	24.5	0.30	10.48	0.16	0.07
Vreta Kloster	35.1	0.12	10.54	0.37	0.17
Bjertorp	21.6	0.52	10.47	0.65	0.41
Ekebo	9.9	0.91	11.26	1.97	0.88
Fjärdingslöv	25.7	0.67	10.63	0.63	0.46
Bornsjön	27.7	0.29	10.45	0.10	0.04
Lilla Böslid	15.6	0.54	10.60	0.90	0.48
20E	21.4	0.15	10.41	0.37	0.27

¹See equation 1: A low value of the Esser index indicates a large amount of hydroxy-interlayering, whereas for the other methods, low values indicate a small amount of hydroxy-interlayering

²See equation 2

³See equation 3.

⁴See equation 4: A low value of the change in stability indicates a large thermostability of the hydroxy-interlayers.




Figure 6. Hydroxy-interlayered vermiculite (HIV) index (Method 4) for pure vermiculite, synthetic HIV and clay fractions from soil samples.

Oxalate extraction of the clay fractions from the soils originating from the unfertilised treatments in the long-term fertility experiments was used as a complementary method for determining the amount of hydroxy-interlayering. A strong correlation ($r = 0.95^{***}$) was found between the amount of hydroxy-interlayering (Method 4a) and oxalate-extractable Al (ranging between 2.18 to 7.49 mg kg⁻¹ clay fraction). However, the extraction did not affect the samples to an extent that was visible by XRD.

5.2 The effect of weathering and soil development, an example from Lanna (Paper III)

5.2.1 Mineralogy affected by weathering

The Lanna soil consisted of 60 – 70 % quartz, plagioclase and K-feldspar, and the mineral composition was largely consistent throughout the profile (Table 7). There was also a significant proportion of clay minerals (between 21 and 33 %). The profile had a higher content of amphibole and iron oxides in the subsoil than the topsoil. The clay content and pH also increased with depth.

When evaluating the clay fraction, the topsoil was found to contain a relatively large proportion of mica/illite, whereas enrichment of expandable minerals was observed in the subsoil. Kaolinite was found in small amounts in the profile, and no chlorite was detected. The minerals in the topsoil showed somewhat higher resistance to collapse upon heating, which is an indication of more extensive hydroxy-interlayering (Table 8). However, the clay fraction of the topsoil was also rich in quartz and feldspar.

Table 7. *Bulk soil mineralogy of the Lanna soil as determined by XRD after spray drying*

Depth	Quartz	Plagioclase	K-feldspar	Amphibole	Iron oxides	Clay minerals ¹
	(% of bulk soil)					
0-10 cm	31	22	20	1.9	2.3	23
10-30 cm	30	22	20	1.8	2.4	24
30-50 cm	21	22	19	1.9	3.0	33
50-70 cm	21	23	20	2.3	3.1	31
70-100 cm	22	23	19	2.5	2.7	31

¹The value in this column includes patterns of chlorite, muscovite, biotite, hydrobiotite, vermiculite, illite, mixed layer minerals with illite/smectite and kaolinite

Table 8. Clay mineralogy and index for the amount of hydroxy-interlayering (Method 3) in samples from the Lanna soil profile

Depth	Kaolinite	Chlorite (% of clay fraction)	Mica/illite	Expandable minerals	COG K-sat 300°C (Å)
0-10 cm	2	<i>n.d.</i>	23	75	10.5
10-30 cm	3	<i>n.d.</i>	21	76	10.4
30-50 cm	2	<i>n.d.</i>	17	81	10.2
50-70 cm	3	<i>n.d.</i>	14	83	10.2
70-100 cm	3	<i>n.d.</i>	16	81	10.2

5.2.2 Phosphorus speciation affected by soil development

The P speciation was determined by LCF analysis of the XANES spectra. The results shown in Figure 7 present the best fits for each sample. There was a clear trend for increasing amount of apatite with increasing depth in the soil profile, from 15 % in the topsoil to 86 % in the 70-100 cm layer. By contrast, a decreasing percentage of PO₄ adsorbed to Al hydroxides was found, with 55 % and 14 % of the P in the topsoil and 70-100 cm layer, respectively.

Organically bound P was found in the topsoil. The standard that best described organic P was lecithin, a diester organic P. The P speciation fitting results for the clay fraction indicated that the topsoil was enriched in Al-hydroxide-bound P and organic P relative to the bulk soil. In contrast, fits for the subsoil showed relative depletion of apatite in the clay fraction compared with the bulk soil, which may indicate preferential weathering of apatite from the finer fractions.




Figure 7. Phosphorus speciation in: a) the bulk soil (mg P kg⁻¹ HCl-digestible P) and b) the clay fraction (% of total P) in the Lanna soil profile.

5.2.3 Phosphorus solubility patterns affected by soil development

The batch experiments showed that the solubility of PO_4 in the Lanna soil increased with decreasing pH in samples from all horizons. The lowest dissolved PO_4 concentrations were observed between pH 5.5 and pH 7, depending on the horizon. The topsoil had a higher PO_4 solubility and a relatively low pH at which the minimum P solubility was observed (Figure 8).

Furthermore, the solubility of Ca, Al and Fe increased with decreasing pH, but for the topsoil the solubility of Fe and Al increased above pH 5 and 6, respectively. This may be explained by an increase in dissolved organic matter (DOM) containing dissolved Fe(III)- and Al-organic complexes at higher pH (Gustafsson & van Schaik, 2003). Using the Visual MINTEQ software (Gustafsson, 2014), it was shown that the samples from the 50-70 and 70-100 cm horizons were supersaturated with respect to hydroxyapatite at natural pH (no acid or base added) and also at higher pH, whereas the samples from the surface horizons were undersaturated (Figure 9). This suggests that the apatite in the surface horizons is thermodynamically unstable and likely to dissolve.




Figure 8. Solubility patterns of: a) PO_4 , b) Ca c) Fe and d) Al as influenced by pH for the Lanna soil.




Figure 9. Stability diagram for the equilibrated suspensions of the pH-dependent sorption experiment. The points represent the calculated solution activities in Visual MINTEQ. The solubility line for hydroxyapatite was calculated from the solubility constant given in Visual MINTEQ ($\log *K_s = -44.3$ at 25°C). Data points above the line indicate supersaturation.

5.3 Phosphorus speciation affected by long-term fertilisation (Papers II and IV)

The overall XANES signal was found to be most closely related to the HCl-extractable P in the soil ($\hat{P}\text{-HCl}$; Paper IV). Therefore Figure 10a and 11a present the different P species relative to this fraction.

5.3.1 Phosphorus speciation in soils and clay fractions with no added phosphorus

The LCF analysis of the unfertilised samples showed that the P speciation differed between soil samples from different field sites (Figure 10). All bulk soils contained organic P, but only three of the clay fractions indicated appreciable amounts of organic P and no significant correlation was found between organic carbon and organic P ($r=0.77$ n.s.). However, the three soils that were relatively high in organic C also had organic P in the clay fraction. All bulk soils except the sample from Ekebo contained apatite or some other form of calcium phosphate. For the former, a strong positive relationship was observed between the fitted proportions of calcium phosphates (including apatite) and pH ($r=0.97^{***}$). With LCF, calcium phosphates were identified also in the clay fractions except for Fjärdingslöv. Aluminium or Fe phosphate

mineral phases were not detected by LCF in any of the samples. In the bulk soils, most of the PO₄ adsorbed to metal oxides had been adsorbed to Al hydroxides, and a minor part to Fe (hydr)oxides. It was more difficult to differentiate between adsorbed P to the different metal (hydr)oxides in the clay fractions, which may be a result of the noisier data obtained at NSLS. In the clay fractions from Kungsängen, Bjertorp and Ekebo, fitting indicated P adsorbed to Fe oxides. In soils with more than 35 mmol kg⁻¹ oxalate-extractable Fe, a major proportion of P was adsorbed to Fe (hydr)oxides in the clay fraction. A strong correlation was observed between the ratio of oxalate-extractable Al to Fe and the amount of P adsorbed to Al hydroxides in the clay fraction ($r=0.92^{**}$). For all bulk soils except Ekebo, there was a significant contribution of P adsorbed to Fe oxides. A correlation was found between the amount of P adsorbed to Al- and Fe (hydr)oxides and the suggested index for hydroxy-interlayering (Method 1a, clay content; $r=0.84^{*}$). For the clay fractions, there was also a correlation between PsTotP and the amount of calcium phosphates ($r=0.89^{*}$).

5.3.2 Phosphorus speciation in soils and clay fractions after long-term fertilisation

In both bulk soil and clay fractions, K-edge XANES fitting results indicated that Al-bound P was important for sorption of P added by fertilisation (Figure 11). However, in two of the soils (Vreta Kloster and Fjärdingslöv), part of the added P appeared to be precipitated as apatite, and in three soils (Kungsängen, Vreta Kloster and Bjertorp) adsorption to iron oxides was also important.




Figure 10. Speciation of P in: a) bulk soils (mg P kg⁻¹ HCl-digestible P) and b) clay fractions (%) from the unfertilised treatment (A3) of the six long-term fertility experiment sites.

5.4 Dissolved phosphate as affected by long-term fertilisation (Paper IV)

The pH dependence of the PO₄ solubility was different for P-depleted and P-fertilised soils (Figures 12-14). In the unfertilised soils, there was a clear increase in the PO₄ solubility with decreasing pH (Figure 13). In the fertilised soils, a greater dissolved P concentration was found. In some cases (Bjertorp, Kungsängen, Ekebo), the effect of pH on PO₄ solubility was small and/or inconsistent. The inconsistency agrees with observations by Gustafsson *et al.* (2012), who attributed it to different pH-dependent behaviour for desorption of native soil PO₄ (increasing with decreasing pH) and desorption of recently added PO₄ (decreasing with decreasing pH).

However, in the Fjärdingslöv and Vreta Kloster soils, a similar or stronger trend of dissolved P increasing with decreasing pH after fertilisation was found, with minimum dissolved P occurring at pH ~7 (Figure 14). At neutral soil pH, most of the soils were undersaturated with respect to hydroxyapatite. The exception was Fors, for which both the unfertilised and fertilised soils were supersaturated (Figure 12). For the fertilised soils from Vreta Kloster and Fjärdingslöv, the degree of undersaturation was relatively small, which may indicate apatite dissolution.




Figure 11. Speciation of P in: a) bulk soils (mg P kg⁻¹ HCl-digestible P) and b) clay fractions (%) from the fertilised treatment (D3) of the six long-term fertility experiment sites. A3 is the P content from the unfertilised treatment.




Figure 12. Stability diagrams for the equilibrated suspensions of the pH-dependence experiment for a) Fors, b) Kungsängen, c) Vreta Kloster, d) Bjertorp, e) Ekebo and f) Fjärdingslöv. 'A3' represents the unfertilised treatment, 'C3' and 'D3' fertilised treatments with addition of surplus P + 15/20 kg and 30 kg, respectively. The points represent the calculated solution activities in Visual MINTEQ. The solubility line for hydroxyapatite was calculated from the solubility constant given in Visual MINTEQ ($\log *K_s = -44.3$ at 25°C). Data points above the line indicate supersaturation.

The XANES results (Figure 11) also showed that apatite had accumulated in these two soils as a result of fertilisation. In other words, the increased solubility of the Vreta Kloster and Fjärdingslöv soils could result from apatite formation in response to high P fertilisation (Figures 13 and 14). As the apatite formed is unstable at low pH, it may have dissolved (although not at equilibrium), creating the pH-dependent solubility behaviour shown in Figure 15. However, the possibility cannot be entirely excluded that the results are attributable to weathering of apatite originally present in the P-depleted (unfertilised) treatment, whereas the apatite was more resistant to solubility in the fertilised treatment.




Figure 13. pH-dependent solubility of $\text{PO}_4\text{-P}$ in the unfertilised soils from the long-term fertility experiments.




Figure 14. pH dependent-solubility of $\text{PO}_4\text{-P}$ in the fertilised soils from the long-term fertility experiments.




Figure 15. pH-dependent solubility of: a) Ca, b) Fe and c) Al in soils from the unfertilized treatments of the long-term fertility experiments.

5.5 Phosphate sorption affected by long-term fertilisation (Paper IV)

The PO₄ sorption varied between different soils (see Table 9). In most cases there was stronger PO₄ sorption in the unfertilised soil than in the fertilised counterpart. If the Freundlich exponent m is kept constant, the Freundlich coefficient K_F can be seen as an estimate of the sorption strength. By stepwise regression, a multivariate linear relationship between K_F and oxalate-extractable Al and Fe was found (adjusted $R^2 = 70\%$):

$$K_F = 0.0529 + 0.000918 \times Al_{ox} + 0.000239 \times Fe_{ox} \quad (7)$$

where Al_{ox} and Fe_{ox} denote oxalate-extractable Al and Fe, respectively (in $mmol\ kg^{-1}$). For fertilised soils, an even stronger relationship was found when competition from organic acids (using soil organic C as a surrogate) was considered in addition to oxalate-extractable Al (adjusted $R^2 = 79\%$)

$$K_F = 0.120 + 0.000809 \times Al_{ox} - 0.00280 \times \frac{OrgC}{Al_{ox} + Fe_{ox}} \quad (8)$$

where OrgC is the soil organic C content (%). Moreover, there was a correlation between the sorption of P (as indicated by K_F) and the amount of hydroxy-interlayering in the clay fraction ($r = 0.95^*$).

The initial adsorbed PO₄ (n_{init}) was related to the P extracted by the P-AL extraction ($r = 0.72^{***}$). However, it is clear that P-AL overestimated adsorbed PO₄ in the soils with reactive apatite, as the acid ammonium lactate solution is likely to cause partial dissolution of the calcium phosphate minerals (Figure 16). A relationship was also found between the amount of adsorbed PO₄ in the XANES analysis of the sorption experiments and the hydroxy-interlayering index (Method 4a \times Clay content; $r = 0.97^{***}$).




Figure 16. Relationship between the initial adsorbed PO_4 n_{init} , determined from the sorption equations and the extracted P in ammonium lactate.

Table 9. pH range, K_F , n_{init} and coefficient of determination (R^2) for the Freundlich equation calculated for each sample from the long-term fertility experiments

Soil	Range of pH values after equilibration	K_F	n_{init}	R^2
Fors A3	7.55-7.77	0.085	1.05	0.989
Fors C3	7.50-7.67	0.091	2.54	0.989
Fors D3	7.49-7.64	0.090	2.94	0.994
Kungsängen A3	6.01-6.11	0.165	1.37	0.992
Kungsängen C3	6.06-6.12	0.148	2.67	0.987
Kungsängen D3	6.13-6.16	0.165	3.25	0.992
Vreta Kloster A3	6.12-6.17	0.092	0.94	0.994
Vreta Kloster C3	6.22-6.31	0.131	5.06	0.976
Vreta Kloster D3	6.20-6.27	0.142	5.84	0.990
Bjertorp A3	6.07-6.12	0.094	0.99	0.990
Bjertorp C3	6.22-6.31	0.105	4.76	0.978
Bjertorp D3	6.20-6.28	0.140	5.78	0.988
Ekebo A3	6.07-6.12	0.135	1.15	0.999
Ekebo C3	6.17-6.23	0.140	3.50	0.993
Ekebo D3	6.23-6.41	0.154	4.85	0.987
Fjärdingslöv A3	6.29-6.38	0.088	0.78	0.997
Fjärdingslöv C3	6.27-6.34	0.089	2.48	0.996
Fjärdingslöv D3	6.31-6.40	0.097	4.10	0.991

6 Discussion

6.1 Characterising hydroxy-interlayering in clay fractions and soils (Paper I)

There are different ways in which the amount of hydroxy-interlayering can be defined: (1) The amount of hydroxy-interlayers in a specific mineral; (2) the amount of hydroxy-interlayering in the clay fraction; and (3) the amount of hydroxy-interlayering of the whole soil. There is also a difference between the *amount* of hydroxy-interlayers in the samples and the *degree* of hydroxy-interlayering. The amount of hydroxy-interlayers describes the total content of hydroxy-interlayers in the sample, whereas the degree refers to the fraction of the interlayer space that consists of hydroxy-interlayers. The methods used in this thesis estimate the amount of hydroxy-interlayering in the clay fraction. All methods investigated consistently identified the same soils as having the highest amount of hydroxy-interlayering.

A feature in common for all methods is that they involve the term *stability*. In general, stability can be conceptualised as the resistance to collapse caused by the displacement of cations, dissolving reagents, heating or ageing. The methods suggested by Esser (1990) and Matsue & Wada (1988) involve collapse of the interlayer space caused by K-saturation, using the Mg-saturated sample as a reference. However, this method does not provide a direct measure of the amount of hydroxy-interlayering, but instead it describes a property that is dependent on the amount of hydroxy-interlayering. Another issue with this type of method is the risk of incomplete collapse as a result of K-saturation (Sayegh *et al.*, 1965), especially of expandable minerals. In Method 3 the samples were heated to 300 °C before measurement to decrease rehydration and re-expansion. This procedure seems to result in total collapse for samples low in hydroxy-interlayers, but overestimates the collapse for samples with a higher amount of hydroxy-interlayering. A general impression of all methods

outlined is that they all seem to work well for ranking soils with high amounts of interlayering, but for soils that are low in hydroxy-interlayers, the measurements seem more inconsistent.

The new method (Method 4) quantifies the resistance to collapse upon *in situ* heating. This method involves a more direct measurement of the amount of hydroxy-interlayering than the methods of Esser (1990) and Matsue and Wada (1988). At 250 °C there is complete dehydration of the interlayers of hydroxy-interlayered vermiculite, causing a decrease in the *d*-spacing (*i.e.* contraction) of the mineral. Further increases in the temperature result in dehydroxylation of the hydroxy-Al or hydroxy-Fe polymers in the interlayer, which causes a further decrease of the *d*-spacing (Harris *et al.*, 1992). Hence this method includes contributions from both dehydration and dehydroxylation processes, but only the latter are directly related to the amount of hydroxy-interlayering.

The idea behind the new method (referred to here as Method 4b) is to obtain a more specific estimate of the thermostability of the hydroxy-interlayers, by quantifying the change in the *d*-spacing (as indicated by COG) between 250 and 450 °C. Ideally, this would reflect only the dehydroxylation of the hydroxy-interlayers, and thereby the amount of hydroxy-Al-interlayering.

For the set of soils studied, it was found that the soils with the highest amount of hydroxy-interlayers also had the lowest stability. Soils with low pH commonly also had higher amount of hydroxy-interlayers in the clay fraction, but at the same time they had lower clay content. This probably reflects the fact that the weathering process tends to slow up at higher pH and consequently there is less formation of these hydroxy-interlayers (Stevens & Bayard, 1994).

At present, the new method cannot be claimed to work 'better' than any of the existing methods. However, it can be claimed that it represents a step forward in the determination of hydroxy-interlayering, as it is based on fundamental properties of hydroxy-interlayered minerals.

6.2 How does soil development affect P speciation? (Papers I and III)

6.2.1 Formation of hydroxy-interlayers and (hydr)oxides

Hydroxy-interlayers are commonly found in dioctahedral minerals as a weathering product of dioctahedral mica (Farmer *et al.*, 1988; Brown, 1953). However, trioctahedral minerals with hydroxy-interlayers can also be found, as a result of weathering of biotite (Wilson, 1999). Hydroxy-interlayered minerals are commonly formed by weathering of the edges, forming hydroxy-Al or hydroxy-Fe polymers. Weathering also produces secondary Al and Fe(III) oxide-type phases, which may adsorb PO₄ (Prietzal *et al.*, 2013). For the Lanna

soil, amphiboles were suggested as sources of Al and Fe in secondary oxides, as indicated by the parallelism of the gradients of the decreasing amount of PO₄ adsorbed to Al hydroxides and the increasing amount of amphiboles with increasing depth. The high amount of expandable minerals may also be a result of vermiculitisation, which may be a further source of Fe for forming Fe oxide-type phases (Farmer *et al.*, 1971) and of Al.

For the clay fraction samples from the long-term fertility experiments, XRD patterns suggest that hydroxy-interlayer material is apparently unaffected oxalate extraction. There was, however, a clear relationship between the amount of extracted Al and the amount of hydroxy-interlayers in the clay fraction. These observations can be interpreted in two different ways: (1) a large part of the oxalate-extracted Al consists of interlayer Al, but the latter is only a small fraction of the amount of hydroxy-interlayers (small enough not to be seen by XRD), and/or (2) the correlation between oxalate-extractable Al and hydroxy-interlayered Al represent two different pools that both have a common source of formation. With our data, it is not possible to tell which of these interpretations that is the most likely one. As a consequence, it is uncertain to what extent the P-adsorbing Al-hydroxide represents a precipitated mineral phase (such as Al hydroxide, gibbsite or allophane), or a hydroxy-Al-interlayer with similar P binding properties

6.2.2 Weathering of apatite

For the Lanna soil, it was observed that the amount of apatite increased with depth. According to Walker & Syers (1976), during soil weathering, total P and apatite contents can be expected to decrease, whereas the relative amount of other P species increases. For the Lanna soil, depletion of apatite was observed in the clay fraction compared with the bulk soil. This is probably because fine-grained apatite dissolves faster due to its higher surface area. This assumption was supported by the observation that apatite was thermodynamically unstable in the upper 50 cm of the pedon. The low content of apatite in the topsoil may be caused by long-term weathering, plant uptake of cations and accumulation of organic material, all of which would contribute to lower pH and increasing dissolution of apatite.

In the fertilised samples from Fjärdingslöv and Vreta Kloster, the apatite content was higher than in the unfertilised samples. There are at least two possible reasons for this: (1) The unfertilised samples may have had a higher apatite dissolution rate due to the P-depleted conditions; or (2) Ca phosphates may have formed after fertilisation. The results available do not allow an assessment of which, if either, of these explanations may be correct.

6.2.3 Organic P and PO₄ adsorbed to Fe (hydr)oxides in the topsoil

In the Lanna soil, the XANES speciation results only indicated the presence of organic P in the topsoil, along with PO₄ adsorbed to Fe (hydr)oxides. It has been pointed out in several reviews (e.g. Colombo *et al.*, 2014; Zhu *et al.*, 2014; Borch *et al.*, 2010) that there is a strong relationship between the C and Fe cycles. One of the reasons for this relationship is that plants produce organic ligands, e.g. oxalate and citrate (Colombo *et al.*, 2014; Schwertmann, 1991), which may release Fe(III) that form Fe(III) (hydr)oxides.

According to the XANES results, the best fit for organic P was with lecithin, which is a phosphate diester compound. However, ³¹P-NMR analysis by other workers showed that monoesters dominate in the organic P fraction of Swedish agricultural soils (Ahlgren *et al.*, 2013). The reason for this discrepancy is not clear. One explanation may be degradation or lyophilisation of diesters forming monoesters during the extraction in sodium hydroxide which is performed prior to NMR analysis (Ahlgren *et al.*, 2013). Another possible reason is that K-edge XANES spectroscopy is not sensitive enough to distinguish between different organic P compounds, due to their relatively featureless XANES spectra on the K edge. To obtain more conclusive evidence on the true nature of organic P, the use of P L_{2,3}-edge XANES spectroscopy may be preferable (Kruse *et al.*, 2009).

6.3 Phosphorus speciation, which P is available? (Papers II, III and IV)

A major proportion of P in all soils was present as calcium phosphates, PO₄ bound to Al hydroxides and organic P forms. Furthermore, accumulation of PO₄ adsorbed to Al hydroxides after fertilization was suggested for both bulk soil and clay fraction.

The issue of which extraction agent best reflects available P in soil is an ever-going (and possibly never-ending) discussion due to the inherent lack of specificity and operational nature of destructive chemical fractionations. In this study, it was concluded that the P extracted by ammonium lactate (P-AL) is fairly well related to the initial adsorbed P in the sample, as shown by sorption isotherm experiments with 7 d equilibration (Figure 16). However, this is not true for soils that contain reactive apatite. For the samples from the Swedish long-term fertility experiment, Bergström *et al.* (2015) observed that during the last 50 years the P-AL pool in the unfertilized treatments slowly decreased with time. This finding indicates that adsorbed PO₄ is potentially available to plants, and therefore the (hydr)oxide-adsorbed P content will decrease during P depletion.

Concerning the pH-dependent solubility of PO_4 , it was observed that the solubility increased at lower pH, which can be explained by various processes: (1) Dissolution of the Al hydroxide sorbents responsible for P sorption; and (2) dissolution of calcium phosphates or apatite. There is reason to believe that both of these processes are involved. An example is the Lanna soil, where both the 0-10 cm and 70-100 cm horizons displayed increased P dissolution at lower pH, despite the former containing little apatite, while the P speciation of the latter sample was dominated by apatite. Gustafsson *et al.* (2012) argued that the pH dependence may also be affected by the texture in the soil. In general, the soils included in this thesis all showed the same trend (Figure 13). However for the unfertilised Ekebo and Bjertorp samples, $\text{PO}_4\text{-P}$ did not increase until pH decreased below 5, which represents a slightly different behaviour than that of the other clay soils, and of the clay soils examined by Gustafsson *et al.* (2012). This may be explained at least in part by the lower content of apatite in the Ekebo and Bjertorp soils (Figure 9). Thus, the relatively high solubility of PO_4 at low pH often found in Swedish clay soils may be explained by relatively high accumulation of apatite and hydroxy-Al compounds that are unstable under low pH conditions. Moreover, because the chemical conditions in the soil generally do not allow neoformation of Ca phosphates (except possibly after heavy fertilisation in some soils), (hydr)oxide surfaces are likely to be the most important for P sorption.

At pH >7, an increase in the P solubility can be observed. This is most probably caused by the pH-dependent adsorption/desorption behaviour of PO_4 on Fe and Al (hydr)oxide sorbents, caused in turn by *e.g.* an increased negative charge of the sorbent at higher pH.

6.4 Is it possible to develop a standard procedure for XANES data treatment?

There is no unequivocal answer that can be given to the question of what set of parameters should be used for background subtraction and normalisation of P K-edge XANES spectra. At the same time, small changes in these parameters can make a large difference to *e.g.* the height of the white line, and this will affect any interpretation concerning *e.g.* quantitative estimates of different phases using LCF, or any similar method.

One example of the above, which is important in the context of this thesis, is the relative proportions of PO_4 adsorbed to Al and Fe (hydr)oxides determined in XANES fitting analyses. The XANES spectra of these phases showed great similarities: the main difference was that Fe-bound PO_4 had a small pre-edge feature, while there were also small differences in the post-edge

region. However, in the LCF analysis of soil samples that only contained low concentrations of P, these two phases may be difficult to differentiate. When analysing such small differences, good data quality (*i.e.* high signal-to-noise ratio) is very important. It is possible that the noise may explain the difference observed between the P speciation of the bulk soil and of the clay fractions from Kungsängen (Figure 10). The bulk soil samples, the spectra of which were collected at SLRI, had much better data quality, and therefore these results (showing predominance of P sorbed to Al over P sorbed to Fe) can be deemed more reliable. Consequently, in the conclusions of this thesis, more weight is placed on the XANES results from SLRI (Papers III and IV) than on the NSLS results (Paper II), although for the most part the trends in both cases were in qualitative agreement.

A certain pre-edge and post-edge normalisation range is commonly used to get a consistent method for XANES data interpretation. Eveborn *et al.* (2009) used a wide normalisation range with a quadratic equation, whereas others have suggested the use of smaller ranges for environmental samples, *e.g.* Werner & Prietzel (2015). The former alternative could be preferred in cases in which the absorbance is strong enough for clear EXAFS oscillations to occur. Normalisation has been shown to be the most important factor affecting the outcome and performance of LCF and principal component analysis (PCA) (Werner & Prietzel, 2015; Beauchemin *et al.*, 2003).

Based on the experience gained during the work presented in this thesis, I suggest a given set of normalisation parameters to start with for background subtraction and normalisation (for P K-edge XANES spectra from SLRI: -30 to -10 eV for the pre-edge range and +30 to +45 eV for the normalisation range). However, theoretically the background subtraction and normalisation equations would ideally result in parallel lines across the XANES spectrum, and this is not always observed with the starting parameters. Therefore I suggest small changes in the normalisation range to get the lines parallel or, if this is not possible, nearly parallel. Another way to get more reliable results when differences or similarities are assessed is to use one of the samples as a standard to fit to the other as was done for the speciation of the additional P derived from fertilizer application. It is sometimes also important to limit the number of standards before performing LCF to get an acceptable fit. In this study, PCA was used for this purpose (suggested by Beauchemin *et al.*, 2003). Furthermore, as mentioned by Werner & Prietzel (2015), it is important to verify the result with other methods. After all, LCF analysis is a statistical method with its own limitations.

6.5 Evaluation of hypotheses

1. *Soils with a large proportion of hydroxy-Al-interlayered clay minerals have higher P sorption capacity than those with a small proportion.*

Yes, but it remains uncertain if this is a result of sorption to hydroxy-interlayers. The strong relationship between the amount of hydroxy-interlayering and P sorption may also be explained by a common source of formation of hydroxy-Al-interlayers and P-sorbing Al hydroxides, as there is a strong relationship between the latter.

2. *a) In clay soils with low P status, P is mainly sorbed to Al hydroxide phases.*

Yes, as observed in the XANES analyses of the unfertilised soil samples from the long-term fertility experiments

b) In clay soils with high P status, P is bound to both Al and Fe (hydr)oxide phases.

In three of the six soils from the fertilised treatment in the long-term fertility experiments, sorption to Fe (hydr)oxides was observed by XANES analysis. These soils commonly have a high content of oxalate-extractable Fe. Therefore this hypothesis can be confirmed for soils with a high Fe content.

c) Ca-bound P as apatite is only reactive in calcareous soils.

No, reactive apatite was observed after fertilisation by XANES analysis in two non-calcareous soils. This observation was supported by pH-dependent solubility experiments.

3. *Apatite increases with depth in the soil profile as a result of its dissolution in the upper horizons during pedogenic weathering.*

Yes, as observed by XANES analysis in the Lanna soil profile.

4. *Solubility of phosphate (PO_4) in Swedish agricultural clay soils is lowest at near neutral pH and increases with both increasing and decreasing pH.*

This was true for most unfertilized soils (exception: Ekebo, for which the minimum solubility of PO_4 occurred at pH 4.5). After fertilization, the pH dependence became smaller except in cases where reactive apatite was present.

7 Conclusions

- A new hydroxy-Al-interlayering index was developed, in which the shift in the centre of gravity of the XRD pattern between 250 and 450 °C was determined. The amount of hydroxy-interlayering was found to be different in different soils from the Swedish long-term soil fertility experiments.
- There is a strong relationship between the amount of hydroxy-interlayers and the concentration of Al hydroxides, as evidenced by oxalate extractions. This makes it difficult to evaluate whether the Al-hydroxide sorbent that retains P is present as a mineral phase or in an hydroxy-interlayer.
- For a soil profile from a Swedish agricultural soil, a more or less gradual shift of P speciation was observed with increasing soil depth. In the topsoil, P speciation was dominated by PO₄ sorbed to Al/Fe (hydr)oxides and organic P, whereas in the subsoil most P was present as apatite.
- Apatite, organic P and PO₄-P sorbed to Al hydroxides are common P constituents in topsoils. Fertilisation brings about an increase in the pool of P adsorbed to oxide surfaces, mainly Al hydroxides. In unfertilized soils, the lowest solubility of P was generally found at neutral pH (6.5-7.5). The P solubility increased after long-term excess P fertilisation.

8 Implications and future perspectives

This thesis showed that PO_4 in clay soils usually has its lowest solubility at neutral pH and that its solubility increases at decreasing pH. The ammonium lactate extraction used in Sweden as soil test P (P-AL; e.g. Eriksson *et al.*, 2013) seems to be a good indicator of the adsorbed P, except for soils dominated by calcium phosphates, where ‘available P’ seems to be overestimated by the extraction. Soil mapping in Sweden is recommended every 10 years, which for P with its slow processes seems a relevant time period. However, more research is needed on ways to identify areas with apatite-rich soils, and also on ways to implement the value of P extracted in ammonium lactate in these soils.

In this thesis, it is suggested that oxalate-extractable Al and Fe could be used as an indication of the sorption properties of soil samples. However, for the fertilised soils studied an even stronger relationship was found between oxalate-extractable Al minus the ratio of organic C to the sum of oxalate-extractable Al and Fe. The latter could be seen as competition by organic acids for the sorbing surfaces, but Al surfaces seem to be the most important surface available. More research is needed to elucidate the P sorption processes and how they are affected by competition from other anions and organic acids. In Sweden, it has been suggested that measurements of Al and Fe extractable in ammonium lactate should be used to evaluate the properties of sorbing surfaces (Ulén, 2006). However, Eriksson *et al.* (2013) found a correlation between oxalate-extractable and ammonium lactate-extractable Al, but not as clear a correlation as for Fe. More research is needed to define a simple way to characterise the sorption processes in soils.

Svanbäck *et al.* (2013) found that soils with higher amount of oxalate-extractable Al and Fe also commonly had lower P losses from the topsoil, even after fertilisation. This thesis confirmed for these soils that most of the PO_4 was adsorbed to Al- and Fe (hydr)oxides. However, a combination of chemical and

physical parameters, *e.g.* infiltration capacity and structure, is needed to further understand the processes affecting the P losses (Djordjic *et al.*, 2004).

More research is also needed to really understand ongoing sorption/desorption processes in soil, and how they may be affected by soil pH. Such results may help understand the processes determining PO_4 availability in soils, and thereby decrease the need for fertilisation of apatite minerals and mitigation methods in order to decrease the P losses. However, the values obtained are just for the whole soil system, and may not represent what is available for the crops. More research is needed on how plants use this pH-dependent desorption behaviour and the effects on field scale.

References

- Ahlberg, A., Olsson, I. & Simkevicius, P. (2003). Triassic-Jurassic weathering and clay mineral dispersal in basement areas and sedimentary basins of southern Sweden. *Sedimentary Geology*, 161, pp. 15-29.
- Ahlgren, J., Djodjic, F., Börjesson, G. & Mattsson, L. (2013). Identification and quantification of organic phosphorus forms in soils from fertility experiments. *Soil Use and Management*, 29, pp. 24-35.
- Andersson, H., Bergström, L., Ulén, B., Djodjic, F. & Kirchmann, H. (2015). The role of subsoil as a source or sink for phosphorus leaching. *Journal of Environmental Quality*, 44, pp. 535-544.
- Andersson, H., Bergström, L., Djodjic, F., Ulén, B. & Kirchmann, H. (2013). Topsoil and subsoil properties influence phosphorus leaching from four agricultural soils. *Journal of Environmental Quality*, 42, pp. 455-463.
- Antelo, J., Fiol, S., Perez, C., Marino, S., Arce, F., Gondar, D. & Lopez, R. (2010). Analysis of phosphate adsorption onto ferrihydrite using the CD-MUSIC model. *Journal of Colloid and Interface Science*, 347, pp. 112-119.
- Aronsson, H., Stenberg, M. & Ulén, B. (2011). Leaching of N, P and glyphosate from two soils after herbicide treatment and incorporation of a ryegrass catch crop. *Soil Use and Management*, 27, pp. 54-68.
- Barnhisel, R.I. & Bertsch, P.M. (1989). Chlorites and hydroxy-interlayered vermiculite and smectite. *Minerals in Soil Environments, SSSA Book Series, no. 1.*, pp. 729-788.
- Barrow, N.J. (2008). The description of sorption curves. *European Journal of Soil Science*, 59(5), pp. 900-910.
- Barrow, N.J., Bowden, J.W., Posner, A.M. & Quirk, J.P. (1980). Describing the effects of electrolyte on adsorption of phosphate by a variable charge surface. *Australian Journal of Soil Research*, 18, pp. 395-404.
- Beauchemin, S., Hesterberg, D., Chou, J., Beauchemin, M., Simard, R.R. & Sayers, D.E. (2003). Speciation of phosphorus in phosphorus-enriched agricultural soils using X-ray absorption near-edge structure spectroscopy and chemical fractionation. *Journal of Environmental Quality*, 32(5), pp. 1809-1819.

- Bergström, L., Kirchmann, H., Djodjic, F., Kyllmar, K., Ulén, B., Liu, J., Andersson, H., Aronsson, H., Börjesson, G., Kynkänniemi, P., Svanbäck, A. & Villa, A. (2015). Turnover and losses of phosphorus in Swedish agricultural soils: long-term changes, leaching trends, and mitigation measures. *Journal of Environmental Quality*, 44, pp. 512-523.
- Bergström, L., Jarvis, N. & Stenström, J. (1994). Pesticide leaching data to validate simulation models for registration purposes. *Journal of Environmental Science and Health Part A-Environmental Science and Engineering & Toxic and Hazardous Substance Control*, 29, pp. 1073-1104.
- Boesch, D., Hechy, R., O'Melia, C., Schindler, D. & Seitzinger, S. (2006). *Eutrophication of Swedish Seas*. Report 5509, Stockholm: Swedish Environmental Protection Agency.
- Borch, T., Kretzschmar, R., Kappler, A., van Cappellen, P., Ginder-Vogel, M., Voegelin, A. & Campbell, K. (2010). Biogeochemical redox processes and their impact in contaminant dynamics. *Environmental Science & Technology*, 44, pp. 15-23.
- Brady, N.C. & Weil, R.R. (1999). Soil phosphorus and potassium. In: *The Nature and Properties of Soils*. 12th. ed. New Jersey: Prentice-Hall, Inc., pp. 540-584.
- Brindley, G.W. & Brown, G. (1980) Crystal structures of clay minerals and their X-ray identification. Mineralogical Society, Monograph No. 5. London
- Brown, G. (1953). The dioctahedral analogue of vermiculite. *Clay Miner. Bull.*, 2, pp. 64-70.
- Bruker, A. (2005). *EVA DIFFRAC^{plus}*. (Version: 11, 0, 0, 3) [Computer Program]. Germany.
- Carlgren, K. & Mattsson, L. (2001). Swedish soil fertility experiments. *Acta Agriculturae Scandinavica Section B-Soil and Plant Science*, 51, pp. 49-78.
- Carstea, D.D. (1968). Formation of hydroxy-Al and -Fe interlayers in montmorillonite and vermiculite -influence of particle size and temperature. *Clays and Clay Minerals*, 16, pp. 231-238.
- Chang, S.C. & Jackson, M.L. (1957). Fractionation of soil phosphorus. *Soil Science*, 84, pp. 133-144.
- Colombo, C., Palumbo, G., He, J.-H., Pinton, R. & Cesco, S. (2014). Review on iron availability in soil: interaction of Fe minerals, plants, and microbes. *Journal of Soils and Sediments*, 14, pp. 538-548.
- Cordell, D., Drangert, J.O. & White, S. (2009). The story of phosphorus: Global food security and food for thought. *Global Environmental Change - Human and Policy Dimensions*, 19, pp. 292-305.
- Cui, Y.S. & Weng, L.P. (2013). Arsenate and Phosphate Adsorption in Relation to Oxides Composition in Soils: LCD Modeling. *Environmental Science & Technology*, 47, pp. 7269-7276.

- Devau, N., Hinsinger, P., Le Cadre, E., Colomb, B. & Gérard, F. (2011). Fertilization and pH effects on processes and mechanisms controlling dissolved inorganic phosphorus in soils. *Geochimica et Cosmochimica Acta*, 75, pp. 2980-2996.
- Djordjic, F., Börling, K. & Bergström, L. (2004). Phosphorus leaching in relation to soil type and soil phosphorus content. *Journal of Environmental Quality*, 33, pp. 678-684.
- Egnér, H., Riehm, H. & Domingo, W.R. (1960). Investigations on chemical soil analysis as the basis for estimating the nutrient status of soils. II. Chemical methods of extraction for phosphorus and potassium determinations. *Kungliga Lantbrukshögskolans Annaler*, 26, pp. 199-215.
- Eriksson, A.K., Ulén, B., Berzina, L., Iital, A., Janssons, V., Sileika, A.S. & Toomsoo, A. (2013). Phosphorus in agricultural soils around the Baltic Sea - comparison of laboratory methods as indices for phosphorus leaching to waters. *Soil Use and Management*, 29, pp. 5-14.
- Esser, K.B. (1990). X-ray-diffraction indexes for relative quantification of interlayering in phyllosilicates. *Soil Science Society of America Journal*, 54, pp. 923-926.
- Eveborn, D., Gustafsson, J.P., Hesterberg, D. & Hillier, S. (2009). XANES speciation of P in environmental samples: an assessment of filter media for on-site wastewater treatment. *Environmental Science & Technology*, 43, pp. 6515-6521.
- Farmer, V.C., Russell, J.D., McHardy, W.J., Newman, A.C.D., Ahlrichs, J.L. & Rimsaite, J.Y. (1971). Evidence for loss of protons and octahedral iron from oxidized biotites and vermiculites. *Mineralogical Magazine*, 38, pp. 121-137.
- Farmer, V.C., Smith, B.F.L., Wilson, M.J., Loveland, P.J. & Payton, R.W. (1988). Readily-extractable hydroxyaluminium interlayers in clay-sized and silt-sized vermiculite. *Clay Minerals*, 23, pp. 271-277.
- Franke, R. & Hormes, J. (1995). The P K-near edge absorption spectra of phosphates. *Physica B*, 216, pp. 85-95.
- Gieseking, J.E. (1975) Soil components. Volume 2. Inorganic components. Soil components. Volume 2. Inorganic components. Springer-Verlag., Berlin, and New York, , German Federal Republic.
- Gustafsson, J.P. (2014). *Visual MINTEQ, version 3.1*. [Computer Program]. Stockholm: KTH Royal Institute of Technology. Web: <http://vminteq.lwr.kth.se>.
- Gustafsson, J.P., Mwamila, L.B. & Kergoat, K. (2012). The pH dependence of phosphate sorption and desorption in Swedish agricultural soils. *Geoderma*, 189/190, pp. 304-311.
- Gustafsson, J.P. & van Schaik, J.W.J. (2003). Cation binding in a mor layer: batch experiments and modelling. *European Journal of Soil Science*, 54, pp. 295-310.

- Hansen, H.C.B., Hansen, P.E. & Magid, J. (1999). Empirical modelling of the kinetics of phosphate sorption to macropore materials in aggregated subsoils. *European Journal of Soil Science*, 50, pp. 317-327.
- Harris, W.G., Hollien, K.A., Bates, S.R. & Acree, W.A. (1992). Dehydration of hydroxy-interlayered vermiculite as a function of time and temperature. *Clays and Clay Minerals*, 40, pp. 335-340.
- Hedley, M.J., Stewart, J.W.B. & Chauhan, B.S. (1982). Changes in inorganic and organic soil-phosphorus fractions induced by cultivation practices and by laboratory incubation. *Soil Science Society of America Journal*, 46, pp. 970-976.
- Hesterberg, D., Zhou, W.Q., Hutchison, K.J., Beauchemin, S. & Sayers, D.E. (1999). XAFS study of adsorbed and mineral forms of phosphate. *Journal of Synchrotron Radiation*, 6, pp. 636-638.
- Hillier, S. (1999). Use of an air brush to spray dry samples for X-ray powder diffraction. *Clay Minerals*, 34, pp. 127-135.
- Hillier, S. (2003). Quantitative analysis of clay and other minerals in sandstones by X-ray powder diffraction (XRPD). *Int. Assoc. Sedimentol. Spec. Publ.*, 34, pp. 213-251.
- Ingall, E.D., Brandes, J.A., Diaz, J.M., de Jonge, M.D., Paterson, D., McNulty, I., Elliott, W.C. & Northrup, P. (2011). Phosphorus K-edge XANES spectroscopy of mineral standards. *Journal of Synchrotron Radiation*, 18, pp. 189-197.
- ISO 11277 (2009). Soil quality - Determination of particle size distribution in mineral soil material - Method by sieving and sedimentation. International Organisation for Standardization, Geneva, Switzerland.
- ISO 11466 (1995). Soil quality - Extraction of trace elements soluble in aqua regia. International Organisation for Standardization, Geneva, Switzerland.
- Janssen, R.P.T., Bruggenwert, M.G.M. & van Riemsdijk, W.H. (1997). Interactions between citrate and montmorillonite Al hydroxide polymer systems. *European Journal of Soil Science*, 48, pp. 463-472.
- Johansson, S. (1944). Soil and water on the Lanna experimental farm. *Sveriges Geologiska Undersökningar, Årsbok*, 38, pp. 41.
- Kaiser, K. & Zech, W. (1996). Defects in estimation of aluminum in humus complexes of podzolic soils by pyrophosphate extraction. *Soil Science*, 161(7), pp. 452-458.
- Kar, G., Hundal, L.S., Schoenau, J.J. & Peak, D. (2011). Direct chemical speciation of P in sequential chemical extraction residues using P K-edge X-ray absorption near-edge structure spectroscopy. *Soil Science*, 176, pp. 589-595.
- Kelly, S.D., Hesterberg, D. & Ravel, B. (2008). Analysis of soils and minerals using X-ray absorption spectroscopy. In: Ulery, A.L. & Drees, L.R. (eds) *Methods of soil analysis, Mineralogical methods*. (SSSA Book Series, SSSA, pp. 387-463.

- Kirchmann, H. (1991). Properties and classification of soils of the Swedish long-term fertility experiments. 1. Sites at Fors and Kungsängen. *Acta Agriculturae Scandinavica*, 41, pp. 227-242.
- Kirchmann, H., Eriksson, J. & Snäll, S. (1999). Properties and classification of soils of the Swedish long-term fertility experiments - IV. Sites at Ekebo and Fjärdingslöv. *Acta Agriculturae Scandinavica Section B-Soil and Plant Science*, 49, pp. 25-38.
- Kirchmann, H., Snäll, S., Eriksson, J. & Mattsson, L. (2005). Properties and classification of soils of the Swedish long-term fertility experiments: V. Sites at Vreta Kloster and Högåsa. *Acta Agriculturae Scandinavica Section B-Soil and Plant Science*, 55, pp. 98-110.
- Kizewski, F., Liu, Y.T., Morris, A. & Hesterberg, D. (2011). Spectroscopic approaches for phosphorus speciation in soils and other environmental systems. *Journal of Environmental Quality*, 40(3), pp. 751-766.
- KLS (1965). *Kungliga Lantbruksstyrelsens kungörelse med bestämmelser för undersökning av jord vid statens lantbrukskemiska kontrollanstalt och lantbrukskemisk kontrollstation och lantbrukskemisk station med statens fastställda stadgar. (The announcement of the Royal Agriculture Administration for soil analysis at the Agricultural Chemistry National Institute and control stations and agricultural stations ruled by governmental regulations.) [in Swedish.]* Kungliga Lantbruksstyrelsens kungörelser.
- Kruse, J., Leinweber, P., Eckhardt, K.U., Godlinski, F., Hu, Y.F. & Zuin, L. (2009). Phosphorus L(2,3)-edge XANES: overview of reference compounds. *Journal of Synchrotron Radiation*, 16, pp. 247-259.
- Kynkäänniemi, P., Ulén, B., Torstensson, G. & Tonderski, K.S. (2013). Phosphorus retention in a newly constructed wetland receiving agricultural tile drainage water. *Journal of Environmental Quality*, 42, pp. 596-605.
- Leinweber, P., Haumaier, L. & Zech, W. (1997). Sequential extractions and P-31-NMR spectroscopy of phosphorus forms in animal manures, whole soils and particle-size separates from a densely populated livestock area in northwest Germany. *Biology and Fertility of Soils*, 25, pp. 89-94.
- Leri, A.C., Hay, M.B., Lanzirotti, A., Rao, W. & Myneni, S.C.B. (2006). Quantitative determination of absolute organohalogen concentrations in environmental samples by X-ray absorption spectroscopy. *Analytical Chemistry*, 78, pp. 5711-5718.
- Mareschal, L., Ranger, J. & Turpault, M.P. (2009). Stoichiometry of a dissolution reaction of a trioctahedral vermiculite at pH 2.7. *Geochimica et Cosmochimica Acta*, 73, pp. 307-319.
- Marwa, E.M.M., Hillier, S., Rice, C.M. & Meharg, A.A. (2009). Mineralogical and chemical characterization of some vermiculites from the Mozambique Belt of Tanzania for agricultural use. *Clay Minerals*, 44, pp. 1-17.

- Matsue, N. & Wada, K. (1988). Interlayer minerals of partially interlayered vermiculites in dystrochrepts derived from tertiary sediments. *Journal of Soil Science*, 39, pp. 155-162.
- Mattson, S., Williams, E.G., Koutler-Andersson, E. & et al. (1950). Phosphate relationships of soil and plant. 5. Forms of P in the Lanna soil. *Kungliga Lantbrukshögskolans Annaler*, 17, pp. 130-140.
- McLaughlin, M.J., McBeath, T.M., Smernik, R., Stacey, S.P., Ajiboye, B. & Guppy, C. (2011). The chemical nature of P accumulation in agricultural soils-implications for fertiliser management and design: an Australian perspective. *Plant and Soil*, 349, pp. 69-87.
- Moore, D.M. & Reynolds, R.C. (1997). *X-Ray diffraction and the identification and analysis of clay minerals*. Second edition. Oxford University Press.
- Murphy, J. & Riley, J.P. (1962) A modified single solution method for the determination of phosphate in natural waters. *Analytica Chimica Acta*, 27, pp. 31–36.
- Negassa, W. & Leinweber, P. (2009). How does the Hedley sequential phosphorus fractionation reflect impacts of land use and management on soil phosphorus: A review. *Journal of Plant Nutrition and Soil Science-Zeitschrift Fur Pflanzenernahrung Und Bodenkunde*, 172, pp. 305-325.
- Neumann, A., Torstensson, G. & Aronsson, H. (2011). Losses of nitrogen and phosphorus via the drainage system from organic crop rotations with and without livestock on a clay soil in southwest Sweden. *Organic Agriculture*, 1(4), pp. 217-229.
- Oburger, E., Jones, D.L. & Wenzel, W.W. (2011). Phosphorus saturation and pH differentially regulate the efficiency of organic acid anion-mediated P solubilization mechanisms in soil. *Plant and Soil*, 341, pp. 363-382.
- Omoso, O., McCarty, D.K., Hillier, S. & Kleeberg, R. (2006). Some successful approaches to quantitative mineral analysis as revealed by the 3rd Reynolds Cup contest. *Clays and Clay Minerals*, 54, pp. 748-760.
- Penn, C.J., Mullins, G.L., Zelazny, L.W. (2005) Mineralogy in relation to phosphorus sorption and dissolved phosphorus losses in runoff. *Soil Science Society of America Journal*, 69, pp. 1532-1540.
- Pierzynski, G.M., Logan, T.J., Traina, S.J. & Bigham, J.M. (1990). Phosphorus chemistry and mineralogy in excessively fertilized soils - quantitative analysis of phosphorus rich particles. *Soil Science Society of America Journal*, 54, pp. 1576-1583.
- Pierzynski, G.M., McDowell, R.W. & Sims, J.T. (2005). *Chemistry, cycling, and potential movement of inorganic phosphorus in soils*. (Phosphorus: agriculture and the environment. Madison,: American Society of Agronomy.
- Prietzl, J., Dümig, A., Wu, Y., Zhou, J. & Klysubun, W. (2013). Synchrotron-based P K-edge XANES spectroscopy reveals rapid changes of phosphorus speciation in the topsoil of two glacier foreland chronosequences. *Geochimica et Cosmochimica Acta*, 108, pp. 154-171.

- Ravel, B. & Newville, M. (2005). ATHENA, ARTEMIS, HEPHAESTUS: data analysis for X-ray absorption spectroscopy using IFEFFIT. *Journal of Synchrotron Radiation*, 12, pp. 537-541.
- Reynolds, R.C., Jr. (1985). *NEWMOD (C) a computer program for the calculation of one-dimensional diffraction patterns of mixed-layered clays*. [Computer Program]. Hanover, NH, USA.
- Rich, I.C. (1960). Aluminum in Interlayers of vermiculite. *Soil Science Society of America Journal*, 24, pp. 26-32.
- Sawhney, B.L. (1960). Aluminium interlayers in clay minerals, montmorillonite and vermiculite - laboratory synthesis. *Nature*, 187, pp. 261-262.
- Sayegh, A.H., Harward, M.E. & Knox, E.G. (1965). Humidity and temperature interactions with respect to K-saturated expanding clay minerals. *American Mineralogist*, 50(3-4), pp. 490-495.
- Schwertmann, U. (1991). Solubility and dissolution of iron oxides. *Plant and Soil*, 130, pp. 1-25.
- Slaughter, M. & Milne, I.H. (1960). The formation of chlorite-like structures from montmorillonite. *Clays and Clay Minerals*, 7, pp. 114-124.
- Stevens, R.L. & Bayard, E. (1994). Clay mineralogy of agricultural soils (Ap horizon) in Västergötland, SW Sweden. *GFF*, 116, pp. 87-91.
- Stevenson, F.J. (1994). Organic phosphorus and sulfur compounds. In: *Humus Chemistry : genesis, composition, reactions* (2nd edition), John Wiley & Sons, Inc., pp. 112-140.
- Strauss, R., Brümmer, G.W. & Barrow, N.J. (1997). Effects of crystallinity of goethite: II. Rates of sorption and desorption of phosphate. *European Journal of Soil Science*, 48, pp. 101-114.
- Svanbäck, A., Ulén, B. & Etana, A. (2014). Mitigation of phosphorus leaching losses via subsurface drains from a cracking marine clay soil. *Agriculture Ecosystems & Environment*, 184, pp. 124-134.
- Svanbäck, A., Ulén, B., Etana, A., Bergström, L., Kleinman, P.J.A. & Mattsson, L. (2013). Influence of soil phosphorus and manure on phosphorus leaching in Swedish topsoils. *Nutrient Cycling in Agroecosystems*, 96, pp. 133-147.
- Sø, H.U., Postma, D., Jakobsen, R. & Larsen, F. (2011). Sorption of phosphate onto calcite; results from batch experiments and surface complexation modeling. *Geochimica et Cosmochimica Acta*, 75, pp. 2911-2923.
- Tolner, L. & Füleky, G. (1995). Determination of the originally adsorbed soil phosphorus by modified Freundlich isotherm. *Communications in Soil Science and Plant Analysis*, 26, pp. 1213-1231.
- Tolpeshta, II, Sokolova, T.A., Bonifacio, E. & Falcone, G. (2010). Pedogenic chlorites in podzolic soils with different intensities of hydromorphism: origin, properties, and conditions of their formation. *Eurasian Soil Science*, 43, pp. 777-787.
- Ulen, B. (2006). A simplified risk assessment for losses of dissolved reactive phosphorus through drainage pipes from agricultural soils. *Acta Agriculturae Scandinavica Section B-Soil and Plant Science*, 56, pp. 307-314.

- Ulén, B. & Persson, K. (1999). Field-scale phosphorus losses from a drained clay soil in Sweden. *Hydrological Processes*, 13, pp. 2801-2812.
- Ulén, B., von Brömssen, C., Johansson, G., Torstensson, G. & Forsberg, L.S. (2012). Trends in nutrient concentrations in drainage water from single fields under ordinary cultivation. *Agriculture Ecosystems & Environment*, 151, pp. 61-69.
- Ulén, B.M., Larsbo, M., Kreuger, J.K. & Svanbäck, A. (2014). Spatial variation in herbicide leaching from a marine clay soil via subsurface drains. *Pest Management Science*, 70(3), pp. 405-414.
- Walker, T.W. & Syers, J.K. (1976). Fate of phosphorus during pedogenesis. *Geoderma*, 15, pp. 1-19.
- Van Reeuwijk, L.P. (1995). *Procedures for Soil Analysis*. Wageningen, Netherlands: International Soil Reference and Information Centre.
- Weber, K.A., Achenbach, L.A. & Coates, J.D. (2006). Microorganisms pumping iron: anaerobic microbial iron oxidation and reduction. *Nature*, 4, pp. 752-764.
- Weng, L.P., Vega, F.A. & van Riemsdijk, W.H. (2011). Competitive and synergistic effects in pH-dependent phosphate adsorption in soils: LCD modeling. *Environmental Science & Technology*, 45, pp. 8420-8428.
- Werner, F. & Prietzel, J. (2015). Standard protocol and quality assessment of soil phosphorus speciation by P K-edge XANES spectroscopy. *Environmental Science & Technology*, 49, pp. 10521-10528.
- Wiklander, L. (1950). Mineralogical composition of quaternary Swedish clays. *Nature*, 166, pp. 276-277.
- Willett, J.R., Chartres, C.J. & Nguyen, T.T. (1988). Migration of phosphate into aggregated particles of ferrihydrite. *Journal of Soil Science*, 39, pp. 275-282.
- Williams, J.D.H., Syers, J.K. & Walker T, W. (1967). Fractionation of soil inorganic phosphate by a modification of Chang and Jackson's procedure. *Proceedings. Soil Science Society of America*, 31, pp. 736-739.
- Wilson, M.J. (1999). The origin and formation of clay minerals in soils: past, present and future perspectives. *Clay Minerals*, 34, pp. 7-25.
- Wolf, A.M. & Baker, D.E. (1990). Colorimetric method for phosphorus measurements in ammonium oxalate soil extracts. *Communications in Soil Science and Plant Analysis*, 21, pp. 2257-2263.
- Zhu, Y., Duan, G., Chen, B., Peng, X., Chen, Z. & Sun, G. (2014). Mineral weathering and element cycling in soil-microorganism-plant system. *Science China; Earth sciences*, 57, pp. 888-896.

Acknowledgements

The Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning (Formas) is acknowledged for financial support (contract no. 2010-1677). This research was partly carried out at the National Synchrotron Light Source (NSLS) at Brookhaven National Laboratory (BNL), which is supported by the U.S. Department of Energy; Synchrotron Light Research Institute (SLRI), Nakhon Ratchasima, Thailand; and the James Hutton Institute, Aberdeen, Scotland.

Thanks to all the people that have helped me during these years. I would like to express special thanks to:

My main supervisor *Jon Petter Gustafsson*, for giving me the opportunity to work within this subject. I am very grateful that he has shared his deep knowledge of soil chemistry. Special thanks for all help, support and travel companionship during these years.

My co-supervisor *Stephen Hillier*, for welcoming me to the mineralogy lab at the James Hutton Institute and for sharing his deep knowledge of mineralogy.

My co-supervisor *Dean Hesterberg*, for introducing me to XANES analysis, for his encouragement and for sharing his deep knowledge within soil science.

My co-supervisor *Magnus Simonsson*, for critical discussion and for sharing his extensive knowledge about... everything.

My co-supervisor *Barbro Ulén*, for already in my Master's days introducing me to the subject of phosphorus and for sharing her wide knowledge about practical aspects of phosphorus management.

Former and present laboratory staff at the Department of Soil and Environment; especially *Roger Lindberg* for ICP analyses; *Lena Ek* for ICP analyses; *Inger Juremalm* for ICP and LECO analyses and different P

extractions; and *Christina Öhman* for soil texture analyses. Thanks also to *Mirsada Kulenovic* and *Gunilla Bergvall* for always being available for discussions.

Laboratory staff at the James Hutton Institute for helping with the mineralogy analyses. Special thanks to *Helen Pendrowski* and *Ian Phillips* for help in the lab during my visits to Aberdeen.

Paul Northrup, beamline scientist at X15B, NSLS; *Wantana Klushubun* beamline scientist at BL8, SLRI; thanks for help and support during data collection. Special thanks to *Panidtha Sombunchoo*, *Weeraya Wongtepa*, *Supanan Lapboonrueng* and *Chanakarn Cholsuk* for taking care of us during the beam times, and of course all other staff at BL8, SLRI.

Bertil Nilsson at KTH, for help with PO₄-P analyses.

Helena Andersson and *Anders Lindsjö*, for sharing soil samples.

Aidin Geranmayeh, for help with some of the pH-dependent P dissolution experiments.

David Eveborn at JTI, for cooperation with the XANES analyses.

Magnus Simonsson and *Dan Berggren Kleja*, for commenting on the thesis, and *Mary McAfee*, for language editing.

Ragnar Persson and *Håkan Karlsson*, for help with computers.

Pia Edfeldt, *Anne Olsson* and *Anna Wennberg*, for administrative help.

Former and present PhD students at the Department of Soil and Environment. Special thanks to the “soil chemistry students” *Maja*, *Charlotta*, *Sabina*, *Åsa* and *Marguerite*. Thanks also to the phosphorus PhDs *Jian*, *Masud*, *Helena*, *Annika*, *Pia*, *Julia*, *Ana*, *Frank* and *Matthew*. Great thanks to some really good friends at work, including those mentioned earlier and *Linnea*, *Martin*, *Johan*, *Joris*, *Veera*, *Kristin*, *Carina*, *Carin*, *Sabine*, *Abdul* and *Maria*. Thanks *Athefeh* for taking care of me in the Cunningham Building during my first trip to Aberdeen.

Former and present colleagues at the Department of Soil and Environment.

All students I have met in different courses, for giving inspiration and asking questions about the project, forcing me to think in a different way. Thanks for asking general questions about soil and environmental science, forcing me to gain deeper knowledge about these topics. Great thanks to *Bernt Andersson* at the teaching lab, for giving not only suggestions for the teaching, but also valuable comments on the project.

All friends for help and support during these years. *Axel*, “I Hope You Know That You're Invaluable”.

My mother *Gun-Britt* and father *Jan-Erik*, for always taking care of me as your little daughter.

And last, but not least... Thanks to *Alice, Valentina, William, Sofie, Amanda* and all the other children in this world, who I hope can use this work to create a more sustainable world.