

# Spatial patterns and broad-scale weather cues of beech mast seeding in Europe

Giorgio Vacchiano<sup>1,11\*</sup>, Andrew Hacket-Pain<sup>2,3\*</sup>, Marco Turco<sup>4,5</sup>, Renzo Motta<sup>1</sup>, Janet Maringer<sup>6,7</sup>, Marco Conedera<sup>7</sup>, Igor Drobyshv<sup>8,9</sup> and Davide Ascoli<sup>10</sup>

<sup>1</sup>DISAFA, University of Turin, Largo Braccini 2, 10095 Grugliasco (TO), Italy; <sup>2</sup>St Catherine's College, Manor Road, Oxford, OX1 3UJ, UK; <sup>3</sup>Fitzwilliam College, Storeys Way, Cambridge CB3 0DG UK; <sup>4</sup>Department of Applied Physics, University of Barcelona, Av. Diagonal 647, 08028 Barcelona, Spain; <sup>5</sup>Barcelona Supercomputing Center (BSC), c/Jordi Girona 29, 08034 Barcelona, Spain; <sup>6</sup>Institute for Landscape Planning and Ecology, University of Stuttgart, Keplerstr. 11 70174 Stuttgart, Germany; <sup>7</sup>Swiss Federal Institute for Forest, Snow, and Landscape Research WSL, a Ramél 18, CH-6953 Cadenazzo, Switzerland; <sup>8</sup>Southern Swedish Forest Research Centre, Swedish University of Agricultural Sciences, PO Box 49, 230 53 Alnarp, Sweden; <sup>9</sup>Chaire industrielle CRSNG-UQAT-UQAM en aménagement forestier durable, Université du Québec en Abitibi-Témiscamingue, 445 Boulevard de l'Université, Rouyn-Noranda, QC J9X 5E4 Canada; <sup>10</sup>Dipartimento di Agraria, University of Naples Federico II, via Università 100, 80055 Portici (NA), Italy; <sup>11</sup>Present address: European Commission, Joint Research Centre, D1 Bio-economy, Via Fermi 2749, Ispira (VA), Italy

## Summary

Author for correspondence:  
Giorgio Vacchiano  
Tel: +39 329 6497188  
Email: giorgio.vacchiano@unito.it

Received: 15 November 2016  
Accepted: 9 April 2017

*New Phytologist* (2017) **215**: 595–608  
doi: 10.1111/nph.14600

**Key words:** *Fagus sylvatica* (beech), mast seeding, Moran effect, population ecology, seed production, synchronisation, weather cues.

- Mast seeding is a crucial population process in many tree species, but its spatio-temporal patterns and drivers at the continental scale remain unknown.
- Using a large dataset (8000 masting observations across Europe for years 1950–2014) we analysed the spatial pattern of masting across the entire geographical range of European beech, how it is influenced by precipitation, temperature and drought, and the temporal and spatial stability of masting–weather correlations.
- Beech masting exhibited a general distance-dependent synchronicity and a pattern structured in three broad geographical groups consistent with continental climate regimes. Spearman's correlations and logistic regression revealed a general pattern of beech masting correlating negatively with temperature in the summer 2 yr before masting, and positively with summer temperature 1 yr before masting (i.e. 2T model). The temperature difference between the two previous summers (DeltaT model) was also a good predictor. Moving correlation analysis applied to the longest eight chronologies (74–114 yr) revealed stable correlations between temperature and masting, confirming consistency in weather cues across space and time.
- These results confirm widespread dependency of masting on temperature and lend robustness to the attempts to reconstruct and predict mast years using temperature data.

## Introduction

Mast seeding (or masting) is the synchronous and highly variable production of fruits and seeds (Pearse *et al.*, 2016), and is a crucial population process in many grass, shrub and tree species (Kelly & Sork, 2002). As a form of information-mediated interaction, masting synchrony has important implications for broader ecological patterns emerging at the community and ecosystem levels (Mescher & Pearse, 2016). The synchrony of masting varies across species (Norden *et al.*, 2007), time (Drobyshv *et al.*, 2010) and space (Suzuki *et al.*, 2005), with cascading effects on plant regeneration (Ascoli *et al.*, 2015), community composition (Lichti *et al.*, 2014), nutrient fluxes (Zackrisson *et al.*, 1999), carbon allocation (Müller-Haubold *et al.*, 2013) and trophic cascades (Blackwell *et al.*, 2001), including those that involve organisms that carry human infectious diseases (e.g.

Hantaviruses: Clement *et al.*, 2010; Lyme disease: Ostfeld & Keesing, 2000).

Describing and predicting masting is therefore critical to better understand population dynamics, assess present and future ecosystem resilience, and design adaptive forest management strategies (Wagner *et al.*, 2010). In recent decades, the temporal pattern of masting has been described for several species in boreal, temperate and tropical biomes (Koenig & Knops, 2000). A growing body of research has elucidated some of the environmental and physiological cues of masting (e.g. Kelly, 1994; Kelly & Sork, 2002; Kelly *et al.*, 2013; Miyazaki *et al.*, 2014; Pearse *et al.*, 2016), and suggested several mechanisms responsible for the synchronisation of masting in individual species from the stand to the regional scale (Satake & Iwasa, 2000; Koenig & Knops, 2013; Koenig *et al.*, 2015).

The oldest and simplest hypothesis for masting states that seed crops vary in response to weather variation (Büsgen *et al.*, 1929; Kelly, 1994). In particular, temperature and

\*These authors contributed equally to this work.


precipitation in the years previous to seed production have been consistently related to masting (e.g. in grasses: Schaubert *et al.*, 2002; shrubs: Meyer & Pendleton, 2015; trees: Bisi *et al.*, 2016). The nature of such relationships can be either correlative, that is weather is a ‘cue’ that triggers other processes and that plants are able to sense (e.g. Kelly & Sork, 2002; Kelly *et al.*, 2013), or causal, in the case that weather directly influences resources and other proximate causes of masting (Monks *et al.*, 2016; Pearse *et al.*, 2016) (correlative analysis on its own cannot distinguish these models, and we hereafter use the word ‘cue’ in a broad sense without addressing underlying mechanisms). Several studies have assessed masting at the scale of a plant’s distribution range (e.g. Kelly *et al.*, 2000; Schaubert *et al.*, 2002; Masaki *et al.*, 2008), but few have tested whether such cues are constant in space and time across the range. In other words, do spatial and temporal variations in masting emerge from (1) variations in weather, or (2) differences in local plant response to the same weather patterns, e.g. as a consequence of inter-population differences and adaptations in genes that regulate flowering (Tan & Swain, 2006)? Addressing this question will provide important information on predicting masting both in the short and in the long term, such as in the case of masting responses to climate change and the consequences on large-scale ecosystem processes.

The aims of this paper are to: (1) describe the spatial pattern of masting across the entire distribution of European beech (*Fagus sylvatica* L.), one of the most important European forest species (Fig. 1a); (2) measure the strength of the relationship between beech masting and weather, using alternative models built with variables previously found relevant for masting in plants, and beech in particular, that is precipitation, temperature and drought, either using monthly values, or aggregating both by season and to express 2-yr differences (‘delta’ models); and (3) assess the stability of masting–weather correlations through space (i.e. whether the strength, timing and relative importance of weather cues vary across geographical space) and time.

## Materials and Methods

### Beech masting data

To address such questions, we used a recently available, long-term, large-scale database of masting for European tree species (MASTREE: Ascoli *et al.*, 2017). Each observation was characterised by the following measurements: the date of inclusion in the database, the masting proxy considered (e.g. seed, pollen), the scale of measurement of the original data (continuous or ordinal), the year of measurement, the NUTS-1 (Nomenclature of Territorial Units for Statistics version 2013, level 1, that is European administrative subdivisions corresponding to macro-regional aggregations) (data source: GISCO – Eurostat (European Commission)) where the observation was recorded (Fig. 1b), the start and end year of the series, and the length of the continuous



**Fig. 1** (a) Current distribution of beech in Europe at 1-km resolution (Casalegno *et al.*, 2011, filtered for cell cover  $\geq 5\%$ ); (b) number of beech masting data series in each NUTS-1 for the period 1950–2014.

recording series to which each observation belongs (range: 1–191 yr). Non-EU countries where beech masting data were recorded (i.e. Ukraine, Serbia, Switzerland, Bosnia and Herzegovina) were also included in the database with dummy NUTS-1 codes. Observations where NUTS-1 location was uncertain were excluded from further analysis.

All database records belonging to series with length  $\geq 5$  yr were associated to a five-class ordinal measurement (ORDmast) from (1) very poor masting to (5) very abundant masting (Ascoli *et al.*, 2017). To build our target variable we extracted values of ORDmast for beech in the period 1950–2014 (for spatial pattern analysis and Spearman’s correlations) or 1901–2014 (for ordinal regression and moving correlation analysis), because antecedent observations were sparse and unevenly spread across the continent. Pollen and flowering data were excluded, as pooling different masting proxies may introduce some noise, such as would happen should flowers’ crops not mature into fruits owing to environmental constraints. A total of 769 individual series covering 7983 annually resolved observations from 22 European countries was selected for further analyses.

## Data treatment

To obtain long masting series with a minimum amount of missing data, individual masting series were aggregated into 61 NUTS-1 chronologies (NCs) by using the most frequently observed masting class for each year (Supporting Information Table S1). The highest masting class was used in the case of multi-modality, but the impact of a different choice would be negligible (mean Spearman's correlation between NC calculated using highest and lowest masting class in case of multi-modality = 0.91, range = 0.68–1.00). All forests within a NUTS-1 should have homogenous climatic and ecological characteristics; the assumption was tested by computing the mean Spearman's correlation coefficient between individual weather series (see 'Weather cues analysis', period 1901–2014) and between individual masting series in each NUTS-1 (henceforth 'intra-NUTS correlations') with  $\geq 7$  yr of observation in common. This sample size was chosen as a trade-off between the need for robustness (critical value for Spearman's correlation with  $n = 7$  and  $P = 0.05$ :  $\rho = 0.79$ ) and data inclusion (i.e. keeping at least 60% of all NUTS-1 in the analysis). Possible inflation of cross-correlation values due to temporally autocorrelated series was corrected by calculating adjusted degrees of freedom according to Dutilleul *et al.* (1993). Correlation coefficients computed on  $< 30$  observation pairs were corrected according to Hotelling's (1953) second-order transformation. NUTS-1 indeed showed high internal consistency of both weather series ( $\rho > 0.9$ , except for precipitation) and masting series (of 36 NUTS-1 with at least two masting series sharing seven or more elements, 30 had a significantly positive intra-NUTS correlation, with an average value of 0.66) (Fig. S1; Table S2).

The analyses were carried out according to the following scheme: we used Mantel (1967) tests and hierarchical clustering to analyse spatial patterns, Spearman's correlations and ordinal logistic regression to measure the relationship between masting and weather, and moving correlation analysis to assess the temporal stability of such relationship.

## Spatial analysis of masting patterns

To address our first objective, we analysed the spatial structure of masting synchrony at the continental scale by running a Mantel test on NC. This test measures the correlation between two dissimilarity matrices containing measures of geographical and ecological distance. Here, it addresses the hypothesis that masting chronologies farther apart would be less similar to one another (de-synchronised) than closer ones.

Time series of seed production may exhibit lagged negative temporal autocorrelation (e.g. Koenig & Knops, 2000) and non-stationarity (i.e. temporal trends; Allen *et al.*, 2014). Both may alter cross-correlation analyses. Therefore, all NCs were checked for temporal autocorrelation (max. order = 1, excluding segments with  $< 7$  consecutive years of observation which may bias the autocorrelation function; Sork *et al.*, 1993) by fitting a cumulative link regression model (CLM: Agresti, 2002) (probit link with 2000 iterations) against NCs of the previous year (NC<sub>-1</sub>). Of 47

NUTS-1 chronologies with  $\geq 7$  consecutive years of observation, 21 had a significantly negative autocorrelation at lag 1 (Table S2). NCs that exhibited significant temporal trends (i.e. slope of linear regression against year of observation significant at  $P \leq 0.05$ ) were detrended by extracting regression residuals (negative trend: ITI, SI0; positive trend: DEA, DEB, DEC, PL1, PL2, PL5, PL6).

We used coordinates of NUTS-1 centres to compute the geographical distance matrix for the Mantel test, and the index of Suzuki *et al.* (2005), a metric derived from Spearman's correlation coefficient, for the masting dissimilarity matrix. Only NC pairs with an overlap of  $\geq 7$  consecutive years of observation were included in the latter. Alongside the Mantel correlation coefficient, we computed a Mantel correlogram (Oden & Sokal, 1986) by binning distances and calculating the mean pairwise correlation between NCs for pairs in each distance class. The significance of each Mantel correlation coefficient was computed by a randomisation approach (2000 permutations) in which the rows and/or columns of the distance matrices are randomly rearranged (Mantel, 1967), and corrected for multiple comparison according to Holm (1979) in the case of Mantel correlograms. Furthermore, we ran two unidirectional Mantel tests and correlograms as a function of distance along longitude (Dlon) and latitude (Dlat) only, by fixing the other coordinate to its mean value across all NUTS-1, to scrutinise the structure of masting along the two orthogonal geographical directions. Mantel tests and correlograms were run with package NCF v.1.1 (Bjornstad, 2015) for the R statistical framework (R Core Team, 2016).

Second, we assessed the geographical pattern of beech masting in Europe by running a hierarchical cluster analysis on NCs using Ward's minimum variance method (Murtagh & Legendre, 2014), which minimises within-cluster distances relative to between-cluster distances (Ward & Hook, 1963). Only NC pairs sharing  $\geq 7$  consecutive years of observation were included in the dissimilarity matrix. NUTS-1 not satisfying such condition when paired against every other NUTS-1 were filtered out. Dissimilarities between individual NC pairs with an insufficient number of observations (4.9% of all NC pairs) were simulated by a linear model of the form  $a_0 + a_1 \text{Dlon} + a_2 \text{Dlat}$  ( $a_0 = 18.41$ ,  $a_1 = 1.61 \times 10^{-5}$ ,  $a_2 = 1.54 \times 10^{-5}$ , adjusted  $R^2 = 0.35$ ;  $F$ -statistic = 709.9 on 2 and 2672 degrees of freedom,  $P < 0.001$ ). We determined the optimal number of clusters by maximising the index according to Dunn (1974) with the R package NBCLUST (Charrad *et al.*, 2014), and computed cluster stability by nonparametric bootstrap with the R package FPC (Hennig, 2015). The validity of each cluster was also assessed by checking that the mean Spearman correlation between all NC pairs in each cluster was higher than the mean correlation between all pairs from two different clusters.

## Weather cues analysis

To measure the strength of weather cues of masting, we calculated Spearman's correlations between each NC (filtered on  $\geq 7$  yr of observation, after detrending if needed: see above) and the following variables: mean monthly temperature (MEAN),

monthly mean of daily maximum temperature (MAX), monthly mean of daily minimum temperature (MIN), monthly precipitation (PRE), 3-month standardised precipitation index (SPI3; McKee *et al.*, 1993) and 3-month standardised precipitation and evaporation index (SPEI3; Vicente-Serrano *et al.*, 2010). Weather series were obtained by averaging monthly data across all cells included in each NUTS-1 from the gridded database CRU TS 3.23 (0.5° resolution; years 1901–2014) (Harris *et al.*, 2014), and detrended before all subsequent analysis by running a linear filter on each individual monthly variable for the timespan selected (1901–2014 or 1950–2014). SPI3 was calculated using the nonparametric approach described by Hao *et al.* (2014), in which the probability distributions are calculated empirically (Gringorten, 1963), rather than by fitting a parametric distribution function. SPEI3 was calculated from the difference between available water (i.e. 3-month sum of PRE) and the potential evapotranspiration, which is based on the FAO-56 Penman–Monteith estimation (Allen *et al.*, 1998) and directly gridded by the CRU. The difference was fit to a log-logistic probability distribution to transform the original values to standardised units (Vicente-Serrano *et al.*, 2010). SPEI3 measures the climatic water balance and therefore provides a more reliable and spatially comparable measure of drought severity than precipitation alone (Vicente-Serrano *et al.*, 2013). Data extraction and calculation of drought indices were performed with the R packages *cruts* (Taylor & Parida, 2016) and *SPEI* (Beguería & Vicente-Serrano, 2013). Correlations (years 1950–2014) were computed for all 36 months of a 3-yr period, including the calendar year of seed production and the 2 yr prior (lag –1 and –2). After preliminary scrutiny of the most significant correlations, we also ran correlations against aggregated summer (June–July) weather variables of 1 and 2 yr before masting, and against the difference (Delta) between values of each weather variables measured 1 and 2 yr prior (e.g. Delta Temperature, Kelly *et al.*, 2013). The absence of nonlinear relationships was visually checked before running all correlations. For each correlation, significance was tested at the 95% confidence level, with a simple Bonferroni correction, that is adjusting the required alpha value according to the number of comparisons (0.05/36), to account for multiple comparisons. Finally, to assess the similarity of weather spatial patterns to those of masting, we applied hierarchical cluster analysis to each weather variable, using the same settings and period (1950–2014) as the analysis run on masting, and compared the consistency of masting vs weather clusters by computing the overall proportion of NUTS-1 that were classified into the same cluster under both analyses.

### Spatio-temporal stability of weather cues

To test for spatial stability of masting–weather relationships, we fitted a linear model of Spearman's correlation coefficient between masting and MAX of June, July and August of the 1 and 2 yr prior, and latitude. Subsequently, we modelled the eight longest NCs (DE1, DE2, DE9, DEF, DK0, NL1, SE2, UKJ – including 74–115 yearly observations in the period 1901–2014) as a function of detrended weather variables, using ordinal

logistic regression within the R package *rms* (Harrell, 2015). In this analysis, NC<sub>-1</sub> was used as an additional independent variable, to account for potential temporal autocorrelation resulting, for example, from resource depletion (Davis, 1957) or resource switching (Kelly & Sork, 2002). All models were fitted with 44–65 observations (years 1950–2014), and validated using both a new prediction interval (years 1901–1949, 30–41 observations for each NC except NL1 with 10), and a bootstrapped leave-one-out cross-validation run on the calibration time period. Weather variables (i.e. MAX and PRE in June and July –1 and –2, hereafter MAX<sub>JUN-1</sub>, MAX<sub>JUL-1</sub>, MAX<sub>JUN-2</sub>, MAX<sub>JUL-2</sub>, PRE<sub>JUN-1</sub>, PRE<sub>JUL-1</sub>, PRE<sub>JUN-2</sub>, PRE<sub>JUL-2</sub>) were selected based on the previous correlation analysis and evidence from literature on beech masting (e.g. Piovesan & Adams, 2001; Drobyshev *et al.*, 2010). The absence of nonlinear univariate relationships was visually checked before running the models. All independent variables were z-transformed to ensure comparability of effect sizes within models; to account for collinearity among weather variables, optimal models were selected using backward stepwise selection based on the Akaike information criterion (AIC). Nagelkerke  $R^2$  was used to compare models for different NCs.



To test for temporal stability of masting–weather relationships, each of the eight longest NCs was fitted against the four most important weather variables selected by logistic models and correlation analysis (i.e. MAX<sub>JUN-1</sub>, MAX<sub>JUL-1</sub>, MAX<sub>JUN-2</sub> and MAX<sub>JUL-2</sub>), using year as an interaction factor. If the year × MAX interaction is significant, that will suggest a temporal change in masting sensitivity to maximum summer temperatures. Moreover, to test also for nonlinear trends in correlation values, we additionally ran a moving correlation analysis (MCA) between MAX<sub>JUN-1</sub>, MAX<sub>JUL-1</sub>, MAX<sub>JUN-2</sub> and MAX<sub>JUL-2</sub>, and the same eight NCs. MCA was conducted on detrended weather variables using Spearman's rank correlation and a window size of 28 yr, that is the largest window giving four independent intervals for the period 1901–2014. Most series had some missing values, but a minimum of 15 values (i.e. > 50% of years observed) was required for any window.

The R code used for analyses is provided in Notes S1.

## Results

Graphical analysis of mapped NUTS-1 chronologies (Fig. S2) suggested a certain degree of spatial structuring, except when most of the continent exhibited high seed production (e.g. in 1995). The existence of spatial aggregation in masting was confirmed by significantly positive ( $P < 0.01$ ) Mantel correlation coefficients ( $M = 0.53$ , 0.31 and 0.42 for the isotropic, latitude-only and longitude-only tests, respectively) and correlograms (Figs 2, S3). Hierarchical clustering of NCs produced three relatively stable clusters broadly corresponding to southern (SO, cluster stability = 56%), northern (NO, 68%), and eastern Europe (EA, 71%) (Figs 3, S4). Further dendrogram subdivisions suggested differences between Romania, Poland and all other NUTS-1 in EA, between Mediterranean (central Italy) and all other NUTS-1 in SO, and between Atlantic (France, UK) and central NUTS-1 in NO.

**Fig. 2** (a) Mantel correlograms for NUTS-1 masting chronologies for beech (1950–2014). Distance in 500-km wide bins. Black dots indicate significant ( $P \leq 0.05$ ) correlations, sequentially corrected for multiple testing using Holm's procedure. (b) Pairwise Spearman's correlations between NUTS-1 masting chronologies (1950–2014) against raw distance (km); black line, local polynomial regression smoother.




**Fig. 3** Hierarchical clustering of NUTS-1 masting chronologies for beech (1950–2014) based on Suzuki's dissimilarity index (red, eastern cluster; green, northern; blue, southern; grey, no data within beech distribution for the study period). © EuroGeographics for the administrative boundaries. Output clipped on European beech distribution (Fig. 1a). Asterisks indicate NUTS-1 used for analysis of long masting chronologies.

Correlation analysis revealed consistently positive correlations between NC and the temperature of the prior summers at the NUTS-1 level across the species distribution (and in all three clusters), especially when using seasonal summer weather or 2-yr differences (Fig. 4). Correlations were generally strongest for MAX and MEAN (Fig. S5), and to a lesser degree MIN (Fig. S6) (mean correlation across all NUTS-1: 0.38, 0.36, 0.39,  $-0.21$ ,

and 0.28, 0.24, 0.28,  $-0.13$ , respectively, against MAX, MIN, MEAN and PRE in June–July of 2 yr prior and 1 yr prior). MAX<sub>AUG-1</sub> was not a consistent signal across Europe. One-third of NUTS-1 did not have significant correlations ( $R \geq 0.35$  with a sample size of  $n = 61$ ) either with MAX<sub>JUN-1</sub>, MAX<sub>JUL-1</sub> or MAX<sub>AUG-1</sub>, especially those in the Netherlands, Italy and the Carpathian region (the last of these were based on shorter records). Temperature in the summer 2 yr before masting was negatively correlated with NC across the species distribution (Fig. 4), and particularly in cluster NO. Consequently, DeltaT usually produced significant correlations with masting. Weaker (and rarely significant) correlations were found for the autumn and early winter 2 yr before masting (negative MAX<sub>NOV-2</sub> in Austria, Czech Republic, Poland and Germany, positive MAX<sub>DEC-2</sub> in Mediterranean France) and for the late winter and spring of the year before masting (negative MAX<sub>FEB-1</sub> in Belgium and UK, positive MAX<sub>MAR-1</sub> in Austria, Poland and Croatia, negative MEAN<sub>APR-1</sub> in Italy and France). No consistent pattern of correlations was found between NC and temperature in the year of masting, although some regional patterns during spring were found (e.g. positive correlations with MAX<sub>FEB</sub> in Poland and UK, or with MAX<sub>MAY</sub> in Poland).

Correlations between NC and PRE were weaker and much less consistent than with temperature (Fig. S7). Significantly positive correlations with PRE in two summers prior and negative in one summer prior emerged locally (e.g. in Germany, UK, France and Switzerland), although a clear distinction between clusters was not evident. Correlations with summer<sub>2</sub> were on average stronger than with summer<sub>1</sub>. SPI3 and SPEI3 were similar to MAX, with strong and significant correlations in summer<sub>2</sub> and, less strongly, summer<sub>1</sub> (Figs S8, S9), albeit on a more restricted geographical extent (Germany, Denmark, UK, Belgium, Sweden). Spring water balance (PRE, SPEI3) was generally uncorrelated to masting in beech (except a positive correlation of PRE<sub>APR-1</sub> in France and PRE<sub>APR0</sub> in Croatia).

Overall, most NCs conformed to a general pattern of negative correlations with temperature in the summer 2 yr before masting, and positive correlations with temperature in the summer 1 yr before masting (Table 1), with no substantial differences in the response of masting to weather among geographical clusters. Precipitation and drought indices were less strongly and consistently correlated with NC than temperature. Additionally, neither



**Fig. 4** Spearman's correlation between monthly maximum temperature (1950–2014) and NUTS-1 masting chronologies for beech. NUTS-1 are ordered and coloured according to the cluster to which they belong (red, eastern cluster; green, northern; blue, southern; grey, no data within beech distribution for the study period; black, excluded from clustering due to insufficient chronology length). The three bottom lines show correlation against seasonal summer weather (June–July) and the Delta variable (difference between weather variable in year.<sub>2</sub> and year.<sub>1</sub>). The sample size (number of years on record) is reported on the secondary x-axis. (.) Significant at  $P \leq 0.05$ , (\*) significant at  $P \leq 0.00139$  (Bonferroni-corrected). MEAN, the mean correlation for the corresponding month across the study area. JJ.2, June–July, 2 yr prior; JJ.1, June–July, 1 yr prior; delta, JJ.2 – JJ.1.

**Table 1** Common weather cues for beech masting across the species distribution range relative to the year of seed production (summary of correlation analyses)

	Year-2	Year-1	Year 0
Main signal	COOL summer	WARM summer	
Secondary signal	WET summer	DRY summer	
Regional signals		COOL February and April, WARM March, DRY February and autumn	WARM February and May, WET spring

temperature nor precipitation in the year of masting was consistently correlated with NC, except for a positive influence of early spring temperature in northern Europe. The geographical pattern of weather variables in the period 1950–2014 was very similar to that of masting, with rates of agreement between masting and weather clusters ranging from 62% (TMN) to 87% (PRE) (Table 2).

Latitude was not a significant driver of the correlation between masting and MAX (Fig. S10). Ordinal logistic models fitted to the eight longest NCs had medium to high explanatory power

(range of Nagelkerke  $R^2$ : 0.33–0.72, mean = 0.57). Stepwise AIC selection isolated between three and six independent variables (Fig. 5), which corroborated the results of weather correlations computed for NUTS-1. The most commonly selected terms were MAX<sub>JUL-1</sub> (selected in all models) and MAX<sub>JUL-2</sub> (all but one). MAX<sub>JUN-1</sub> or MAX<sub>JUN-2</sub> were additionally selected in five and six models, respectively. NC<sub>-1</sub>, with a negative coefficient, was selected in five models. Precipitation terms were selected less frequently than temperature, and only half of the models included any precipitation term. Standardised coefficients for precipitation were generally lower than those for temperature, indicating a smaller effect on masting. Model validation produced a mean Nagelkerke  $R^2$  of 0.46 after bootstrapped leave-one-out cross-validation (range = 0.53–0.65), and 0.40 after extrapolation to 1901–1950 (0.21–0.55), indicating that models were partially able to reproduce masting variation outside of the training dataset (Table 3). Clusters SO and EA were represented by only one model each, with the latter showing a lower explanatory power and weaker summer<sub>-1</sub> effects.

Temporal trends in masting were significant in NL1, DE1 (negative trend), DE2 and DE9 (positive); however, the interaction between time and summer MAX was significant only in three out of 32 cases (Table 3). MCA applied to the eight longest chronologies revealed generally stable correlations

**Table 2** Proportion of matches from the assignment of NUTS-1 to masting and weather clusters


Weather variable	Weather cluster	Masting-EA	Masting-NO	Masting-SO
MEAN	EA	14	0	2
	NO	4	21	4
	SO	0	0	2
	Accuracy	78.7%		
MIN	EA	11	0	0
	NO	4	21	1
	SO	3	0	7
	Accuracy	61.9%		
MAX	EA	14	0	5
	NO	4	15	1
	SO	0	6	2
	Accuracy	66.0%		
PRE	EA	16	2	1
	NO	2	19	1
	SO	0	0	6
	Accuracy	87.2%		
SPI3	EA	14	0	1
	NO	4	21	2
	SO	0	0	5
	Accuracy	85.1%		
SPEI3	EA	11	0	1
	NO	7	21	2
	SO	0	0	5
	Accuracy	78.7%		

Study species: beech. Accuracy: rate of total matches (masting cluster = weather cluster) to total number of NUTS-1. Cluster identification: EA, Eastern Europe; NO, Northern; SO, Southern. Monthly weather variables: MEAN, mean temperature; MIN, minimum temperature; MAX, maximum temperature; PRE, precipitation; SPI3, standardised precipitation index; SPEI3, standardised precipitation and evaporation index.

between temperature and masting, particularly for MAX<sub>JUL-1</sub> and MAX<sub>JUL-2</sub> (Figs 6, S11), except for DK0 and DE9 (increasingly stronger correlation through time). Most NCs showed decade-long periods when correlations with MAX were nonsignificant, although the timing of these periods was not synchronised across Europe. Some NCs showed evidence of ‘switching’ between July and June temperature (i.e. periods of reduced correlations with MAX<sub>JUL</sub> corresponded to increased correlations with MAX<sub>JUN</sub>, e.g. DK0), but in others the temporal variations in the strength of all four correlations were homogenous (e.g. UKJ). Some NUTS-1 showed ‘parallel’ correlation trends with year-1 and year-2, that is a decreasing importance of positive MAX<sub>1</sub> corresponding to an increasing importance of negative MAX<sub>2</sub>, especially in DE1 and DE2; however, the dataset is too sparse to strongly generalise such evidence.

## Discussion

Using a distribution-wide dataset with *c.* 8000 individual observations, we have shown that a strong spatial structure exists in masting patterns of *F. sylvatica* across its distribution range. Synchrony was higher between neighbouring populations (Fig. S1),



**Fig. 5** Ordinal logistic models of beech masting (eight longest NUTS-1 chronologies) as a function of weather predictors: (a) model statistics for calibration (1950–2014) and validation (1901–1949). Rsq\_1950\_2014 is the  $R^2$  of the calibration dataset, Rsq\_bootstrap is the bootstrapped  $R^2$  from leave-one out cross-validation (1000 re-samples) and Rsq\_1901\_1949 is the  $R^2$  of the predicted values for 1901–1949 vs observed (validation dataset). (b) Standardised model coefficients. Only significant predictors are filled in the table, with the colour depending on the coefficient. Monthly weather variables: MAX, maximum temperature; PRE, precipitation. NC-1, masting in the previous year.

particularly in northern Europe, and strongly declined with distance (Fig. 2), consistent with previous findings for other temperate species (Koenig & Knops, 1998; Garrison *et al.*, 2008; Gallego Zamorano *et al.*, 2016). While synchrony generally declined with distance (e.g. ‘typical’ years with partial masting at the continental scale such as 2002 or 2009; Fig. S2), continental-scale mast years occurred on several occasions, e.g. twice in the last 40 yr (1976 and 1995, with less comprehensive but still widespread events in 1992, 2006 and 2011; Fig. S2). This is consistent with what has been previously reported for beech at both continental (Nussbaumer *et al.*, 2016) and regional scale (e.g. Hilton & Packham, 2003), and it is based on an unprecedented sample size. These distribution-wide mast events may have important implications for large-scale, long-distance ecological processes, such as forest regeneration after large disturbances (Peters *et al.*, 2005; Ascoli *et al.*, 2015; Funk *et al.*, 2015), pollen- and seed-related gene flow (Kremer *et al.*, 2012), bird migration (Koenig & Knops, 2001), predator–prey population dynamics (Blackwell *et al.*, 2001), pest and disease diffusion (Liebhold *et al.*, 2000), biological invasions (Harper, 2005), forest species range shift (Takenaka, 2005) and climate resilience (Mustin, 2013). Even if masting synchrony had little impact over and

**Table 3** Coefficients and statistics of ordinal logistic regression models for beech masting as a function of multiple weather variables in the eight longest NUTS-1 chronologies (backwards stepwise selection by AIC; ns, nonsignificant at  $P > 0.05$ )

Coefficients	DE1	DE2	DE9	DEF	DK0	NL1	SE2	UKJ
NC <sub>-1</sub>	-1.05	ns	-0.79	ns	-1.18	ns	-1.25	-1.00
PRE <sub>JUL-1</sub>	ns							
PRE <sub>JUL-2</sub>	ns	ns	0.61	ns	1.29	ns	ns	ns
PRE <sub>JUN-1</sub>	ns	0.59	ns	ns	ns	ns	ns	ns
PRE <sub>JUN-2</sub>	ns	ns	ns	ns	ns	ns	0.55	ns
MAX <sub>JUL-1</sub>	0.94	0.52	1.49	1.03	1.34	0.49	1.27	1.42
MAX <sub>JUL-2</sub>	-1.06	-0.78	-1.16	-1.14	ns	-1.31	-1.047	-1.16
MAX <sub>JUN-1</sub>	0.88	0.98	0.57	ns	ns	ns	0.75	0.59
MAX <sub>JUN-2</sub>	ns	-0.73	-0.68	-0.84	-1.61	-0.73	ns	-0.65
Model statistics								
Observations	58	65	57	44	65	56	55	65
<i>P</i>	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
<i>R</i> <sup>2</sup> (calibration)	0.54	0.33	0.70	0.46	0.66	0.47	0.72	0.63
<i>R</i> <sup>2</sup> (leave one-out)	0.49	0.27	0.62	0.41	0.61	0.43	0.65	0.58
<i>R</i> <sup>2</sup> (validation)	0.51	0.21	0.32	0.43	0.21	0.40	0.54	0.55
Year in MAX <sub>JUL-1</sub>	-0.19	ns	0.41	ns	ns	-0.69	ns	ns
Year in MAX <sub>JUL-2</sub>	ns	ns	0.47	ns	ns	-0.69	ns	ns
Year in MAX <sub>JUN-1</sub>	ns							
Year in MAX <sub>JUN-2</sub>	ns	0.40	ns	ns	ns	-0.74	ns	ns
Year × MAX <sub>JUL-1</sub>	-0.68	ns						
Year × MAX <sub>JUL-2</sub>	ns							
Year × MAX <sub>JUN-1</sub>	ns	ns	ns	ns	-0.53	ns	ns	ns
Year × MAX <sub>JUN-2</sub>	ns	0.58	ns	ns	ns	ns	ns	ns
Cluster	SO	EA	NO	NO	NO	NO	NO	NO


Monthly weather variables: MAX, maximum temperature; PRE, precipitation. Year and year × MAX were computed using bivariate models with one interaction term. Cluster identification: EA, Eastern Europe; NO, Northern; SO, Southern.

above the local effects through predator satiation and/or enhanced pollination, and is simply a result of the weather cues used locally, when the latter co-vary across large areas the ecosystem consequences may be far-reaching.

Furthermore, the temporal variability of masting in beech showed a distinct spatial structure during the last 65 yr, with three clusters located in northern, southern and eastern Europe (Fig. 3). These clusters correspond closely to weather patterns (Fig. S12), and broadly to regions of Europe influenced by different climate regimes: the northern cluster corresponds to the region of western Europe associated with an oceanic climate strongly influenced by the Atlantic (Cfb according to Peel *et al.*, 2007), the southern cluster overlaps with the Mediterranean region (Csa), while the eastern cluster is the most continental one and is less influenced by Atlantic weather (roughly corresponding to Dfb). Indeed, the longitude-based Mantel test showed a lower correlation coefficient than the latitude-based test, which may be a consequence of weather events characterised by a longitude-based spatial pattern prompting synchronised masting (Fernández-Martínez *et al.*, 2016). Similar spatial structuring in beech has been found by local masting studies, which explained it by an increasing influence of spring frost in more continental areas (Gross, 1934), but also by tree-ring studies, which linked contrasting growth patterns to the different influence of climate teleconnections, such as between the eastern and western Mediterranean basin (Chen *et al.*, 2015; Seim *et al.*, 2015).

Numerous studies have demonstrated that mast years in many tree species are associated with specific weather conditions

(‘weather cues’) before mast events (Koenig & Knops, 2014; Roland *et al.*, 2014), and particularly with summer temperatures 1 and 2 yr before masting (Schauber *et al.*, 2002; Kelly *et al.*, 2013). We found similar results in this study, showing that a small number of weather variables act as strong cues for masting in almost all European beech populations, despite large climatic, genetic and environmental differences. Indeed, individual NCs in which this typical set of cues was not detected were often based on a limited number of observations. We found no substantial differences in these weather cues of masting among regions or clusters using either correlation analysis (Figs 4, S10) or regression models (Fig. 5), nor any significant effect of latitude (Fig. S10). This demonstrated that, across the distribution, the cues for masting are highly spatially consistent, with positive correlations for MAX<sub>JUL-1</sub> (and to a lesser degree MAX<sub>JUN-1</sub>), and negative for MAX<sub>JUL-2</sub> and MAX<sub>JUN-2</sub>, with some local specificities. Combining June and July clearly improved the consistency of strong (and significant) correlations, as did using DeltaT as a synthetic index of temperature differences from year to year (Kelly *et al.*, 2013). In some cases, the seasonal analysis accounted for regional differences in the strongest individual month; AT1–3 were good examples, as they responded more strongly to June temperatures than to July temperatures (in contrast to most other chronologies). In particular, DeltaT led to improved correlations in cases where correlations with MAX<sub>summer-1</sub> and MAX<sub>summer-2</sub> had the expected signal, but were both relatively weak (e.g. DE2, PL2, PL4), or where one individual correlation was much stronger than the other (e.g. DE1, DE2, DEE, SE2, FR6). In the



**Fig. 6** Moving Spearman's correlation (lines: 28-yr timesteps) between the eight longest NUTS-1 beech masting chronologies and maximum temperature (MAX) (1901–2014). Thick lines represent significant ( $P \leq 0.05$ ) correlations.

regression models for the NUTS-1 with the most data, a large proportion of the variance was explained by summer temperature in years  $-1$  and  $-2$ , suggesting that other signals are not very important.

Therefore, we suggest that the observed spatial organisation of masting is more dependent on weather variation across space, rather than on different sensitivities of beech population to the weather cues, in contrast to what Masaki *et al.* (2008) found for *Fagus crenata*. In other words, traits related to masting seem to be the same across the whole beech distribution range, with the exception of small regional differences – for example, a shift of the most important summer month along a latitudinal gradient, or an increased role of temperatures in the months associated with flowering, pollination and seed maturation (Hase, 1964) in northern Europe.

While the well-known relationship between general summer weather and masting in beech was well supported by our results, we were also able to disentangle the relative importance of temperature and precipitation as the dominant cue of masting. Wachter (1964) and Piovesan & Adams (2001) suggested that

summer precipitation or drought, along with or rather than temperature, was the main cue of masting in beech, while Drobyshev *et al.* (2010) found no relationship between masting and summer precipitation or drought (but did find a strong temperature signal). We have shown that summer precipitation in the 2 yr before masting was an important predictor of mast events in some regions (Figs 4, 5), and that summer drought was correlated with NC in some regions (Fig. S9), but that both precipitation and drought were clearly of secondary importance to temperature as a cue of masting. An additional analysis of the relationship between summer MAX and summer SPEI across Europe showed correlations ranging from  $-0.3$  to  $-0.5$  (Fig. S13), suggesting that drought could be more effective in predicting masting in certain locals than in others, and hence the contrasting evidence for the previous year's drought effects in the literature. The effect of spring precipitation appeared generally negligible, contrary to findings in more Mediterranean species (Fernández-Martínez *et al.*, 2015). Additionally, the importance of precipitation did not appear to vary systematically with latitude, for example in northern vs southern regions where summer drought stress may

be limiting (average correlation between latitude and Spearman's coefficient for  $MAX_{JUN-1}$ ,  $MAX_{JUL-1}$  and  $MAX_{AUG-1} = -0.13$ ;  $MAX_{JUN-2}$ ,  $MAX_{JUL-2}$  and  $MAX_{AUG-2} = 0.05$ ;  $PRE_{JUN-1}$ ,  $PRE_{JUL-1}$  and  $PRE_{AUG-1} = 0.07$ ;  $PRE_{JUN-2}$ ,  $PRE_{JUL-2}$  and  $PRE_{AUG-2} = 0.09$ ). Instead, summer temperatures in the previous 2 yr, particularly in July, were always the main cue of masting, with mast years associated with a cool summer 2 yr before masting, and warm conditions in the summer before masting. This is highly consistent with previous findings on the sensitivity to summer temperatures in both *Fagus* and *Nothofagus* (2 yr prior: Gruber, 2003; Richardson *et al.*, 2005; Smaill *et al.*, 2011; Kelly *et al.*, 2013; 1 yr prior: Hase, 1964; Wachter, 1964; Schaubert *et al.*, 2002; Suzuki *et al.*, 2005; Övergaard *et al.*, 2007; Masaki *et al.*, 2008). Recent analyses of the climate sensitivity of beech diameter increment have also showed that cool, moist summers have a positive effect on the growth of the same year, favouring a resource accumulation hypothesis (Dorado Liñan *et al.*, 2017), while high summer temperatures have a negative effect on growth of the following year throughout the whole geographical distribution, including in northern and central regions (Hackett-Pain *et al.*, 2016), which could be interpreted as a growth vs reproduction tradeoff if masting was triggered in those years (Monks & Kelly, 2006; Hackett-Pain *et al.*, 2015).

In addition to weather cues, we also found that masting was strongly affected by negative temporal autocorrelation, that is masting category in the previous year ( $NC_{-1}$ ). Ordinal logistic regression models were consistently able to predict mast years with accuracy (mean  $R^2 = 0.57$ ) using summer temperature (and in some case precipitation) in the two previous years, plus information on the previous year's masting. Negative temporal autocorrelation with a lag of 1 or 2 yr is one of the defining characteristics of masting time-series (Davis, 1957; Sork *et al.*, 1993; Selås *et al.*, 2002; Koenig *et al.*, 2003), and is the mathematical expression of the rarity of consecutive mast years (category 4 or 5 in our dataset; consecutive years of low masting category were instead common). The existence of negative autocorrelation in masting time series has been traditionally interpreted as evidence for resource depletion, that is trees deplete most resources in the mast year, which limits reproduction in the following year and makes consecutive heavy seed crops very rare (Davis, 1957; Sork *et al.*, 1993; Kelly & Sork, 2002). However, recent studies have shown that negative temporal autocorrelation would also emerge if masting were controlled by  $\Delta T$  only (Kelly *et al.*, 2013; Kon & Saito, 2015; but see also Koenig *et al.*, 2015 for criticism of such model).

The strong correlations between masting and weather found by this study do not provide any conclusive evidence to the debate on whether temperature is a 'cue' for trees to trigger high seed crops or whether it acts instead through intermediate steps indicative of a direct mechanistic connection to seed production (Pearse *et al.*, 2014). Koenig & Knops (2000) found that spatial autocorrelation in seed production of northern hemisphere tree species occurred at the same spatial scale as autocorrelation in rainfall and temperature, consistent with the underlying effect of climatic factors on masting. However, they also found that seed production had much higher variability than the weather factors,

implying the existence of nonlinearities in weather effects, or of drivers for masting, which remain unaccounted for.

While strong climate differences exist across the distribution of beech, the majority of populations analysed herein responded similarly to weather (e.g. negative response to temperature and positive to precipitation 2 yr before masting; Table 1). The negative correlation with  $MAX_{JUL-2}$  could be related to resource accumulation in cooler years ('priming' the trees to respond to increased temperature 1 yr later, *sensu* Richardson *et al.*, 2005), an interpretation that is consistent with a model of masting that includes an element of carbon and/or nitrogen limitation (Sala *et al.*, 2012; Müller-Haubold *et al.*, 2015; Monks *et al.*, 2016; Abe *et al.*, 2016; Pearse *et al.*, 2016; Fernandez-Martinez *et al.*, 2017). Indeed, a higher soil moisture due to more precipitation and lower summer temperatures has been shown to increase litter mass loss and N mineralisation and uptake (Gessler *et al.*, 2005; Smaill *et al.*, 2011), which favours masting in beech (Han *et al.*, 2014; Miyazaki *et al.*, 2014).

High temperatures in the summer before masting ( $MAX_{JUL-1}$ ) have been linked to flower primordia differentiation (Wachter, 1964; Gruber, 2003; see also Merkle *et al.*, 1980 for oaks, Allen *et al.*, 2014; Miyazaki *et al.*, 2014), in particular via an increase in endogenous gibberellins (Turnbull, 2011; Pearse *et al.*, 2016). Following this reasoning, we might expect the phenology of primordia differentiation to vary with latitude, creating a geographical gradient in the timing of the previous summer cue similar to the pattern we found in some southern European NUTS-1 (Fig. 4). Additionally, we also found correlations with weather during the periods associated with other known processes that influence flowering phenology, pollen production (Kasprzyk *et al.*, 2014; Pearse *et al.*, 2015) and seed maturation in the year of masting, such as late winter frost (Matthews, 1955; Wachter, 1964), at least in northern Europe. The resource priming in year  $t_2$  can therefore interact with the MAX cue in summer $_{t-1}$  via a resource pulse that boosts an already favourable flower initiation.

Finally, the analysis of some of the longest series available showed that the sensitivity of beech masting to the most important weather cues ( $MAX_{JUL-1}$  and  $MAX_{JUL-2}$ ) was substantially consistent through time in the last century (Fig. 6), with one possible exception (DE9 with the strongest MCA trend of masting, and logistic model with poorest predictive power). While many studies have reported associations between weather cues and mast years, very few had the length of record required to test whether these cues are consistent through time. Additionally, regression models fitted using data from the period 1950–2014 successfully described mast years in the first half of the 20<sup>th</sup> century (Fig. 5) – although we did not switch the periods due to insufficient sample size for model calibration. This is an important result, as there is little existing information on whether climate change affects the sensitivity of masting to weather cues, or whether the timing of cues shifts seasonally as a response to changing temperatures, as it has been demonstrated for leaf and flower phenology (Menzel *et al.*, 2006). Assessing the effects of changing climate on the frequency and timing of mast years is challenging (McKone *et al.*, 1998; Drobyshev *et al.*, 2014). Despite the pre-eminent role of summer MAX, our analysis did not provide any strong evidence

to suggest that the relationships between weather and masting were sensitive to 20<sup>th</sup> century warming (contrary to Övergaard *et al.*, 2007), as predicted by the theoretical model of Kelly *et al.* (2013). This lends robustness to the attempts to reconstruct and predict mast years using temperature data (e.g. Drobyshev *et al.*, 2014). However, this should be tested more thoroughly. In particular, it is still unclear whether both gradual and abrupt (e.g. extreme events) components of climate change influence masting frequency and spatial synchrony within and across species or phylogenetic groups (Koenig *et al.*, 2016), for example through changes in resource levels (Miyazaki, 2013; Allen *et al.*, 2014), pollen availability (Koenig *et al.*, 2015), coexistence of species with different biomass allocation strategies (Pérez-Ramos *et al.*, 2015), and in the interactions between the processes of resource accumulation and flower induction (Monks *et al.*, 2016).

## Acknowledgements

The authors acknowledge Laura McGarty for language editing, and Dr Dave Kelly and an anonymous reviewer for their insightful comments to the preliminary version of the manuscript.

## Author contributions

G.V. and A.H-P. analysed the data and wrote the manuscript; D.A. designed the research, provided and analysed masting data, and wrote parts of the manuscript; M.T. provided and interpreted weather data; I.D., M.C., J.M. and R.M. contributed to research design and data interpretation.

## References

- Abe T, Tachiki Y, Kon H, Nagasaka A, Onodera K, Minamino K, Han Q, Satake A. 2016. Parameterisation and validation of a resource budget model for masting using spatiotemporal flowering data of individual trees. *Ecology Letters* 19: 1129–1139.
- Agresti A. 2002. *Categorical data*. New York, USA: John Wiley & Sons.
- Allen RB, Hurst JM, Portier J, Richardson SJ. 2014. Elevation-dependent responses of tree mast seeding to climate change over 45 years. *Ecology and Evolution* 4: 3525–3537.
- Allen RG, Pereira LS, Raes D, Smith M. 1998. *Crop evapotranspiration. Guidelines for computing crop water requirements*. FAO Irrigation and drainage paper 56. Rome, Italy: FAO.
- Ascoli D, Maringer J, Hackett-Pain A, Conedera M, Drobyshev I, Motta R, Cirolli M, Kantorowicz W, Zang C, Schueler S *et al.* 2017. Two centuries of masting data for European beech and Norway spruce across the European subcontinent. *Ecology*, In press doi: 10.1002/ecy.1785.
- Ascoli D, Vacchiano G, Maringer J, Bovio G, Conedera M. 2015. The synchronicity of masting and intermediate severity fire effects favors beech recruitment. *Forest Ecology and Management* 353: 126–135.
- Beguieria S, Vicente-Serrano SM. 2013. *SPEI: Calculation of the Standardised Precipitation-Evapotranspiration Index*. R package version 1.6. [WWW document] URL <https://CRAN.R-project.org/package=SPEI> [accessed 1 September 2016].
- Bisi F, von Hardenberg J, Bertolino S, Wauters LA, Imperio S, Preatoni DG, Provenzale A, Mazzamuto MV, Martinoli A. 2016. Current and future conifer seed production in the Alps: testing weather factors as cues behind masting. *European Journal of Forest Research* 135: 743–754.
- Bjornstad ON. 2015. *nfc: Spatial Nonparametric Covariance Functions*. R package version 1.1-6. [WWW document] URL <https://CRAN.R-project.org/package=nfc> [accessed 1 September 2016].
- Blackwell GL, Potter MA, Minot EO. 2001. Rodent and predator population dynamics in an eruptive system. *Ecological Modelling* 142: 227–245.
- Büsgen M, Münch E, Thomson T. 1929. *The structure and life of forest trees*. London, UK: Chapman & Hall.
- Casalegno S, Amatulli G, Bastrup-Birk A, Durrant TH, Pekkarinen A. 2011. Modelling and mapping the suitability of European forest formations at 1-km resolution. *European Journal of Forest Research* 130: 971–981.
- Charrad M, Ghazzali N, Boiteau V, Niknafs A. 2014. NbClust: an R package for determining the relevant number of clusters in a data set. *Journal of Statistical Software* 61: 1–36.
- Chen K, Dorado-Liñán I, Akhmetzyanov L, Gea-Izquierdo G, Zlatanov T, Menzel A. 2015. Influence of climate drivers and the North Atlantic Oscillation on beech growth at marginal sites across the Mediterranean. *Climate Research* 66: 229–242.
- Clement J, Maes P, de Strihou CV, van der Groen G, Barrios JM, Verstraeten WW, van Ranst M. 2010. Beechnuts and outbreaks of *Nephropathia epidemica* (NE): of mast, mice and men. *Nephrology Dialysis Transplantation* 25: 1740–1746.
- Davis LD. 1957. Flowering and alternate bearing. *Proceedings of the American Society of Horticultural Science* 70: 545–556.
- Dorado Liñan I, Zorita E, Martínez-Sancho E, Gea-Izquierdo G, Di Filippo A, Gutiérrez E, Levanic T, Piovesan G, Vacchiano G, Zang C *et al.* 2017. Large-scale atmospheric circulation enhances the Mediterranean East-West tree growth contrast at rear-edge deciduous forests. *Agricultural and Forest Meteorology* 239: 86–95.
- Drobyshev I, Niklasson M, Mazerolle MJ, Bergeron Y. 2014. Reconstruction of a 253-year long mast record of European beech reveals its association with large scale temperature variability and no long-term trend in mast frequencies. *Agricultural and Forest Meteorology* 192: 9–17.
- Drobyshev I, Overgaard R, Saygin I, Niklasson M, Hickler T, Karlsson M, Sykes MT. 2010. Masting behaviour and dendrochronology of European beech (*Fagus sylvatica* L.) in southern Sweden. *Forest Ecology and Management* 259: 2160–2171.
- Dunn JC. 1974. Well-separated clusters and optimal fuzzy partitions. *Journal of Cybernetics* 4: 95–104.
- Dutilleul P, Clifford P, Richardson S, Hemon D. 1993. Modifying the t test for assessing the correlation between two spatial processes. *Biometrics* 49: 305–314.
- Fernández-Martínez M, Garbulsky M, Peñuelas J, Peguero G, Espelta JM. 2015. Temporal trends in the enhanced vegetation index and spring weather predict seed production in Mediterranean oaks. *Plant Ecology* 216: 1061–1072.
- Fernández-Martínez M, Vicca S, Janssens IA, Espelta JM, Peñuelas J. 2016. The North Atlantic Oscillation synchronises fruit production in western European forests. *Ecography*. doi: 10.1111/ecog.02296.
- Fernández-Martínez M, Vicca S, Janssens IA, Espelta JM, Peñuelas J. 2017. The role of nutrients, productivity and climate in determining tree fruit production in European forests. *New Phytologist* 213: 669–679.
- Funk KA, Koenig WD, Knops JM. 2015. Fire effects on acorn production are consistent with the stored resource hypothesis for masting behavior. *Canadian Journal of Forest Research* 46: 20–24.
- Gallego Zamorano J, Hokkanen T, Lehikoinen A. 2016. Climate driven synchrony in seed production of masting deciduous and conifer tree species. *Journal of Plant Ecology*, in press. doi: 10.1093/jpe/rtw117.
- Garrison BA, Koenig WD, Knops JM. 2008. Spatial synchrony and temporal patterns in acorn production of California black oaks. In: Merenlender A, McCreary D, Purcell KL, eds. *Proceedings of the sixth symposium on oak woodlands: today's challenges, tomorrow's opportunities*. Albany, CA, USA: USDA Forest Service, Pacific SW Forest and Range Experiment Station, General Technical Report PSW-GTR-217, 343–356.
- Gessler A, Jung K, Gasche R, Papen H, Heidenfelder A, Börner E, Metzler B, Augustin S, Hildebrand E, Rennenberg H. 2005. Climate and forest management influence nitrogen balance of European beech forests: microbial N transformations and inorganic N net uptake capacity of mycorrhizal roots. *European Journal of Forest Research* 124: 95–111.
- Gringorten II. 1963. A plotting rule for extreme probability paper. *Journal of Geophysical Research* 68: 813–814.

- Gross H. 1934. Die Rotbuche in Ostpreußen. *Zeitschrift für Forst- und Jagdwesen* 66: 622–651.
- Gruber R. 2003. Control and forecasting of the fructification of European beech (*Fagus sylvatica* L.) for the stand Zierenberg 38A and the level I stand in Hessen by climate factors. *Allgemeine Forst und Jagdzeitung* 174: 67–79.
- Hackett-Pain AJ, Cavin L, Friend AD, Jump AS. 2016. Consistent limitation of growth by high temperature and low precipitation from range core to southern edge of European beech indicates widespread vulnerability to changing climate. *European Journal of Forest Research* 135: 897–909.
- Hackett-Pain AJ, Friend AD, Lagueard JGA, Thomas PA. 2015. The influence of masting phenomenon on growth–climate relationships in trees: explaining the influence of previous summers' climate on ring width. *Tree Physiology* 35: 319–330.
- Han Q, Kabeya D, Iio A, Inagaki Y, Kakubari Y. 2014. Nitrogen storage dynamics are affected by masting events in *Fagus crenata*. *Oecologia* 174: 679–687.
- Hao Z, AghaKouchak A, Nakhjiri N, Farahmand A. 2014. Global integrated drought monitoring and prediction system. *Scientific Data* 1: 1–10.
- Harper GA. 2005. Heavy rimu (*Dacrydium cupressinum*) mast seeding and rat (*Rattus* spp.) population eruptions on Stewart Island/Rakiura. *New Zealand Journal of Zoology* 32: 155–162.
- Harrell FE. 2015. *rms: Regression Modeling Strategies*. R package version 4.3-1. [WWW document] URL <https://CRAN.R-project.org/package=rms>. [accessed 1 September 2016].
- Harris IP, Jones PD, Osborn TJ, Lister DH. 2014. Updated high-resolution grids of monthly climatic observations—the CRU TS3.10 dataset. *International Journal of Climatology* 34: 623–642.
- Hase W. 1964. Die Buchenmast in Schleswig-Holstein und ihre Abhängigkeit von der Witterung. *Mitteilungen des Deutschen Wetterdienstes* 31: 1–52.
- Hennig C. 2015. *fpc: Flexible Procedures for Clustering*. R package version 2.1-10, 2015. [WWW document] URL <https://CRAN.R-project.org/package=fpc> [accessed 1 September 2016].
- Hilton GM, Packham JR. 2003. Variation in the masting of common beech (*Fagus sylvatica* L.) in northern Europe over two centuries (1800–2001). *Forestry* 76: 319–328.
- Holm S. 1979. A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics* 6: 65–70.
- Hotelling H. 1953. New light on the correlation coefficient and its transforms. *Journal of the Royal Statistical Society Series B (Methodological)* 15: 193–232.
- Kasprzyk I, Ortyl B, Dulcka-Jeż A. 2014. Relationships among weather parameters, airborne pollen and seed crops of *Fagus* and *Quercus* in Poland. *Agricultural and Forest Meteorology* 197: 111–122.
- Kelly D. 1994. The evolutionary ecology of mast seeding. *Trends in Ecology & Evolution* 9: 465–470.
- Kelly D, Geldenhuis A, James A, Penelope Holland E, Plank MJ, Brockie RE, Cowan PE, Harper GA, Lee WG, Maitland MJ *et al.* 2013. Of mast and mean: differential-temperature cue makes mast seeding insensitive to climate change. *Ecology Letters* 16: 90–98.
- Kelly D, Harrison AL, Lee WG, Payton IJ, Wilson PR, Schaubert EM. 2000. Predator satiation and extreme mast seeding in 11 species of *Chionochloa* (*Poaceae*). *Oikos* 90: 477–488.
- Kelly D, Sork VL. 2002. Mast seeding in perennial plants: why, how, where? *Annual Review of Ecology and Systematics* 1: 427–447.
- Koenig WD, Alejano R, Carbonero MD, Fernández-Rebollo P, Knops JM, Marañón T, Padilla-Díaz CM, Pearse IS, Pérez-Ramos IM, Vázquez-Piqué J *et al.* 2016. Is the relationship between mast-seeding and weather in oaks related to their life-history or phylogeny? *Ecology* 97: 2603–2615.
- Koenig WD, Kelly D, Sork VL, Duncan RP, Elkinton JS, Peltonen MS, Westfall RD. 2003. Dissecting components of population-level variation in seed production and the evolution of masting behavior. *Oikos* 102: 581–591.
- Koenig WD, Knops JM. 1998. Scale of mast-seeding and tree-ring growth. *Nature* 396: 225–226.
- Koenig WD, Knops JM. 2000. Patterns of annual seed production by northern hemisphere trees: a global perspective. *American Naturalist* 155: 59–69.
- Koenig WD, Knops JM. 2001. Seed-crop size and eruptions of North American boreal seed-eating birds. *Journal of Animal Ecology* 70: 609–620.
- Koenig WD, Knops JM. 2013. Large-scale spatial synchrony and cross-synchrony in acorn production by two California oaks. *Ecology* 94: 83–93.
- Koenig WD, Knops JM. 2014. Environmental correlates of acorn production by four species of Minnesota oaks. *Population Ecology* 56: 63–71.
- Koenig WD, Knops JM, Carmen WJ, Pearse IS. 2015. What drives masting? The phenological synchrony hypothesis. *Ecology* 96: 184–192.
- Kon H, Saito H. 2015. Test of the temperature difference model predicting masting behavior. *Canadian Journal of Forest Research* 45: 1835–1844.
- Kremer A, Ronce O, Robledo-Arnuncio JJ, Guillaume F, Bohrer G, Nathan R, Bridle JR, Gomulkiewicz R, Klein EK, Ritland K *et al.* 2012. Long-distance gene flow and adaptation of forest trees to rapid climate change. *Ecology* 15: 378–392.
- Lichti NI, Steele MA, Zhang H, Swihart RK. 2014. Mast species composition alters seed fate in North American rodent-dispersed hardwoods. *Ecology* 95: 1746–1758.
- Liebold A, Elkinton J, Williams D, Muzika RM. 2000. What causes outbreaks of the gypsy moth in North America? *Population Ecology* 42: 257–266.
- Mantel N. 1967. The detection of disease clustering and a generalized regression approach. *Cancer Research* 27: 209–220.
- Masaki T, Oka T, Osumi K, Suzuki W. 2008. Geographical variation in climatic cues for mast seeding of *Fagus crenata*. *Population Ecology* 50: 357–366.
- Matthews JD. 1955. The influence of weather on the frequency of beech mast years in England. *Forestry* 28: 107–116.
- McKee TB, Doesken NJ, Kleist J. 1993. The relationship of drought frequency and duration to time scales. In: Proceedings of the 8th Conference on Applied Climatology. Boston, MA, USA: American Meteorological Society, 179–183.
- McKone MJ, Kelly D, Lee WG. 1998. Effect of climate change on mast-seeding species: frequency of mass flowering and escape from specialist insect seed predators. *Global Change Biology* 4: 591–596.
- Menzel A, Sparks T, Estrella N, Koch E, Aasa A, Ahas R, Alm-Kübler K, Bissolli P, Braslavská O, Briede A *et al.* 2006. European phenological response to climate change matches the warming pattern. *Global Change Biology* 12: 1969–1976.
- Merkle SA, Feret PP, Croxdale JG, Sharik TL. 1980. Development of floral primordia in white oak. *Forest Science* 26: 238–250.
- Mescher MC, Pearse IS. 2016. Communicative interactions involving plants: information, evolution, and ecology. *Current Opinion in Plant Biology* 32: 69–76.
- Meyer SE, Pendleton BK. 2015. Seedling establishment in a masting desert shrub parallels the pattern for forest trees. *Acta Oecologica* 65: 1–10.
- Miyazaki Y. 2013. Dynamics of internal carbon resources during masting behavior in trees. *Ecological Research* 28: 143–150.
- Miyazaki Y, Maruyama Y, Chiba Y, Kobayashi MJ, Joseph B, Shimizu KK, Mochida K, Hiura T, Kon H, Satake A. 2014. Nitrogen as a key regulator of flowering in *Fagus crenata*: understanding the physiological mechanism of masting by gene expression analysis. *Ecology Letters* 17: 1299–1309.
- Monks A, Kelly D. 2006. Testing the resource matching hypothesis in the mast seeding tree *Nothofagus truncata* (*Fagaceae*). *Austral Ecology* 31: 366–375.
- Monks A, Monks JM, Tanentzap AJ. 2016. Resource limitation underlying multiple masting models makes mast seeding sensitive to future climate change. *New Phytologist* 210: 419–430.
- Müller-Haubold H, Hertel D, Leuschner C. 2015. Climatic drivers of mast fruiting in European beech and resulting C and N allocation shifts. *Ecosystems* 18: 1083–1100.
- Müller-Haubold H, Hertel D, Seidel D, Knutzen F, Leuschner C. 2013. Climate responses of aboveground productivity and allocation in *Fagus sylvatica*: a transect study in mature forests. *Ecosystems* 16: 1498–1516.
- Murtagg F, Legendre P. 2014. Ward's hierarchical agglomerative clustering method: which algorithms implement Ward's criterion? *Journal of Classification* 31: 274–295.
- Mustin K. 2013. Red noise increases extinction risk during rapid climate change. *Diversity and Distributions* 19: 815–824.
- Norden N, Chave J, Belbenoit P, Caubère A, Châtelet P, Forget PM, Thébaud C. 2007. Mast fruiting is a frequent strategy in woody species of Eastern South America. *PLoS ONE* 2: e1079.
- Nussbaumer A, Waldne P, Etzold S, Gessler A, Benham S, Thomsen IM, Jørgensen BB, Timmermann V, Verstraeten A, Sioen G, *et al.* 2016. Patterns

- of mast fruiting of common beech, sessile and common oak, Norway spruce and Scots pine in Central and Northern Europe. *Forest Ecology and Management* 363: 237–251.
- Oden NL, Sokal RR. 1986. Directional autocorrelation: an extension of spatial correlograms to two dimensions. *Systematic Zoology* 35: 608–617.
- Ostfeld RS, Keasing F. 2000. Pulsed resources and community dynamics of consumers in terrestrial ecosystems. *Trends in Ecology & Evolution* 15: 232–237.
- Övergaard R, Gemmel P, Karlsson M. 2007. Effects of weather conditions on mast year frequency in beech (*Fagus sylvatica* L.) in Sweden. *Forestry* 80: 555–565.
- Pearse IS, Koenig WD, Funk KA, Pesendorfer MB. 2015. Pollen limitation and flower abortion in a wind-pollinated, masting tree. *Ecology* 96: 587–593.
- Pearse IS, Koenig WD, Kelly D. 2016. Mechanisms of mast seeding: resources, weather, cues, and selection. *New Phytologist* 212: 546–562.
- Pearse IS, Koenig WD, Knops JM. 2014. Cues versus proximate drivers: testing the mechanism behind masting behavior. *Oikos* 123: 179–184.
- Peel MC, Finlayson BL, McMahon TA. 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences* 11: 1633–1644.
- Pérez-Ramos IM, Padilla-Díaz CM, Koenig WD, Maranon T. 2015. Environmental drivers of mast-seeding in Mediterranean oak species: does leaf habit matter? *Journal of Ecology* 103: 691–700.
- Peters VS, Macdonald SE, Dale MR. 2005. The interaction between masting and fire is key to white spruce regeneration. *Ecology* 86: 1744–1750.
- Piovesan G, Adams JM. 2001. Masting behaviour in beech: linking reproduction and climatic variation. *Canadian Journal of Botany* 79: 1039–1047.
- R Core Team. 2016. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. [WWW document] URL <https://www.R-project.org> [accessed 1 November 2016].
- Richardson SJ, Allen RB, Whitehead D, Carswell FE, Ruscoe WA, Platt KH. 2005. Climate and net carbon availability determine temporal patterns of seed production by *Nothofagus*. *Ecology* 86: 972981.
- Roland CA, Schmidt JH, Johnstone JF. 2014. Climate sensitivity of reproduction in a mast-seeding boreal conifer across its distributional range from lowland to treeline forests. *Oecologia* 174: 665–677.
- Sala A, Hopping K, McIntire EJ, Delzon S, Crone EE. 2012. Masting in whitebark pine (*Pinus albicaulis*) depletes stored nutrients. *New Phytologist* 196: 189–199.
- Satake A, Iwasa YO. 2000. Pollen coupling of forest trees: forming synchronized and periodic reproduction out of chaos. *Journal of Theoretical Biology* 203: 63–84.
- Schauber EM, Kelly D, Turchin P, Simon C, Lee WG, Allen RB, Payton IJ, Wilson PR, Cowan PE, Brockie RE. 2002. Masting by eighteen New Zealand plant species: the role of temperature as a synchronizing cue. *Ecology* 83: 1214–1225.
- Seim A, Treydte K, Trouet V, Frank D, Fonti P, Tegel W, Panayotov M, Fernández-Donado L, Krusic P, Büntgen U. 2015. Climate sensitivity of Mediterranean pine growth reveals distinct east–west dipole. *International Journal of Climatology* 35: 2503–2513.
- Selås V, Piovesan G, Adams JM, Bernabei M. 2002. Climatic factors controlling reproduction and growth of Norway spruce in southern Norway. *Canadian Journal of Forest Research* 32: 217–225.
- Smaill SJ, Clinton PW, Allen RB, Davis MR. 2011. Climate cues and resources interact to determine seed production by a masting species. *Journal of Ecology* 99: 870–877.
- Sork VL, Bramble J, Sexton O. 1993. Ecology of mast-fruiting in three species of North American deciduous oaks. *Ecology* 74: 528–541.
- Suzuki W, Osumi K, Masaki T. 2005. Mast seeding and its spatial scale in *Fagus crenata* in northern Japan. *Forest Ecology and Management* 205: 105–116.
- Takenaka A. 2005. Local coexistence of tree species and the dynamics of global distribution pattern along an environmental gradient: a simulation study. *Ecological Research* 20: 297–304.
- Tan FC, Swain SM. 2006. Genetics of flower initiation and development in annual and perennial plants. *Physiologia Plantarum* 128: 8–17.
- Taylor BM, Parida B. 2016. *cruts: Interface to climatic research unit time-series version 3.21 data*. R package version 0.3. [WWW document] URL <https://CRAN.R-project.org/package=cruts> [accessed 1 September 2016].
- Turnbull C. 2011. Long-distance regulation of flowering time. *Journal of Experimental Botany* 62: 4399–4413.
- Vicente-Serrano SM, Beguería S, López-Moreno JI. 2010. A multiscalar drought index sensitive to global warming: the standardized precipitation evapotranspiration index. *Journal of Climate* 23: 1696–1718.
- Vicente-Serrano SM, Gouveia C, Camarero JJ, Beguería S, Trigo R, López-Moreno JI, Azorín-Molina C, Pasho E, Lorenzo-Lacruz J, Revuelto J *et al.* 2013. Response of vegetation to drought time-scales across global land biomes. *Proceedings of the National Academy of Sciences USA* 110: 52–57.
- Wachter H. 1964. Über die Beziehung zwischen Witterung und Buchenmastjahren. *Forstarchiv* 35: 69–78.
- Wagner S, Collet C, Madsen P, Nakashizuka T, Nyland RD, Sagheb-Talebi K. 2010. Beech regeneration research: from ecological to silvicultural aspects. *Forest Ecology and Management* 259: 2172–2182.
- Ward JH, Hook ME. 1963. Application of an hierarchical grouping procedure to a problem of grouping profiles. *Educational and Psychological Measurement* 23: 69–82.
- Zackrisson O, Nilsson MC, Jäderlund A, Wardle DA. 1999. Nutritional effects of seed fall during mast years in boreal forest. *Oikos* 84: 17–26.

## Supporting Information

Additional Supporting Information may be found online in the Supporting Information tab for this article:

**Fig. S1** Mean Spearman's rank correlation of masting series within each NUTS-1.

**Fig. S2** NUTS-1 masting chronologies from year 1976 to 2014.

**Fig. S3** Mantel correlograms for NUTS-1 masting chronologies (1950–2014) across latitude and longitude only.

**Fig. S4** Dendrogram for the hierarchical clustering of NUTS-1 masting chronologies (1950–2014).

**Fig. S5** Spearman's correlation between monthly mean temperature (1950–2014) and NUTS-1 masting chronologies.

**Fig. S6** Spearman's correlation between monthly minimum temperature (1950–2014) and NUTS-1 masting chronologies.

**Fig. S7** Spearman's correlation between monthly precipitation (1950–2014) and NUTS-1 masting chronologies.

**Fig. S8** Spearman's correlation between monthly SPI3 (1950–2014) and NUTS-1 masting chronologies.

**Fig. S9** Spearman's correlation between monthly SPEI3 (1950–2014) and NUTS-1 masting chronologies.

**Fig. S10** Linear models of Spearman's correlation between masting and MAX in June, July or August of the 1 and 2 yr prior vs latitude.

**Fig. S11** Summary of moving Spearman's correlation (1901–2014, window size: 28 yr) between the eight longest NCs and MAX<sub>JUL-1</sub> and MAX<sub>JUL-2</sub>.

**Fig. S12** Hierarchical clustering of NUTS-1 weather variables (1950–2014) based on Suzuki's dissimilarity index.

**Fig. S13** Spearman's correlation between MAX in June–July and SPEI3 across the study area, period 1901–2014.

**Table S1** NUTS-1 chronologies of masting from 1901 to 2016 on an ordinal scale of 1 (very poor) to 5 (very abundant); dash, no data

**Table S2** Intra-NUTS correlation of masting series and temporal autocorrelation in NCs

**Notes S1** R code for the analysis carried out in the present paper.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the New Phytologist Central Office.



## About *New Phytologist*

- *New Phytologist* is an electronic (online-only) journal owned by the New Phytologist Trust, a **not-for-profit organization** dedicated to the promotion of plant science, facilitating projects from symposia to free access for our Tansley reviews.
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as ready' via *Early View* – our average time to decision is <28 days. There are **no page or colour charges** and a PDF version will be provided for each article.
- The journal is available online at Wiley Online Library. Visit **www.newphytologist.com** to search the articles and register for table of contents email alerts.
- If you have any questions, do get in touch with Central Office (np-centraloffice@lancaster.ac.uk) or, if it is more convenient, our USA Office (np-usaoffice@lancaster.ac.uk)
- For submission instructions, subscription and all the latest information visit **www.newphytologist.com**