




Carbon sequestration potential and the multiple functions of Nordic grasslands

Ann Norderhaug¹ · Karina E. Clemmensen² · Paul Kardol^{2,3} ·
Anna Gudrun Thorhallsdottir⁴ · Iulie Aslaksen⁵ 

Received: 8 November 2022 / Accepted: 15 April 2023 / Published online: 4 May 2023
© The Author(s) 2023

Abstract

Grasslands are important carbon sinks, but the underlying processes for their soil carbon sequestration potential are still not well understood, despite much attention given to this topic. In Europe, grasslands, especially semi-natural grasslands, are also important for promoting biodiversity. Moreover, recent global reports have highlighted the importance of biodiversity in supporting climate actions. In boreal and alpine regions in the Nordic countries, grasslands also play an important role in milk and meat production and food security. Certain grassland features and management practices may enhance their soil carbon sequestration potential. Semi-natural grasslands maintained by optimized livestock grazing are vital for aboveground biodiversity and show promise for belowground biodiversity and carbon sequestration potential. It is essential to assess the multiple functions of grasslands, particularly semi-natural grasslands, to facilitate the optimization of policy measures across policy areas. Climate and biodiversity policies should not counteract each other, as some do today. This essay addresses the multiple functions of grasslands and calls for more knowledge about carbon sequestration in Nordic grasslands. This will enable the management of these ecosystems to align with climate mitigation, maintain biodiversity, and satisfy the global need for increased food supply.

Keywords Grasslands · Semi-natural grasslands · Soil carbon · Climate change · Grazing · Greenhouse gases · Biodiversity · Food security

✉ Iulie Aslaksen
iulie.aslaksen@ssb.no

¹ Ann Norderhaug Kulturlandskapskonsulent, Nøtterøy, Norway

² Swedish University of Agricultural Sciences (SLU), Uppsala, Sweden

³ Swedish University of Agricultural Sciences (SLU), Umeå, Sweden

⁴ Hólar University, Hólar, Iceland

⁵ Research Department, Statistics Norway, P. O. Box 2633 St. Hanshaugen, 0131 Oslo, Norway

1 Introduction

The potential of grassland soil carbon sequestration has gained increasing attention (Lorenz and Lal 2018). A recent review by Bai and Cotrufo (2022) estimated that grasslands store approximately one-third of the world's terrestrial carbon stocks. Several European grassland studies also support the idea that grasslands can serve as important carbon sinks (Chang et al. 2015, Chang et al. 2016). The Food and Agriculture Organization of the United Nations (FAO) (2017) likewise emphasizes the substantial potential of grasslands to provide large and stable soil carbon sinks.

The carbon sequestration potential of grasslands, however, remains a subject of debate. Rööß et al. (2017) claimed that this potential is limited while Godde et al. (2020) questioned “overly optimistic expectations” and stressed the high context dependency of soil carbon sequestration rates in grasslands. Factors such as climate, vegetation type, soil type and quality, the composition of soil biological communities, and management practices all influence carbon sequestration potential (Garnett et al. 2017). A Mediterranean study emphasized the need for region-specific data on grassland soil carbon to reflect differences in climatic and agricultural conditions and called for a strategic plan to extend on-site field measurements (Aguilera et al. 2021). Their analysis highlights the necessity for refining carbon footprint estimation methods in life cycle assessments of agricultural products, including region-specific data on relevant processes, such as soil carbon sequestration.

Grasslands have, however, multiple functions, including providing habitats for biodiversity. The global report from the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES 2019) highlights the importance of biodiversity conservation as loss of biodiversity is as crucial as climate change. The land report from the Intergovernmental Panel for Climate Change (IPCC 2019), the global IPCC 2022 report (IPCC 2022), and the UN Biodiversity Conference (COP 15) in December 2022 have also pointed to the connections between the climate and biodiversity crises, highlighting the importance of biodiversity and other ecosystem services to counteract and adapt to climate changes, and the need for alignment of climate policy measures to support biodiversity. Loss of biodiversity is intensifying, with potential negative consequences for ecosystem services, including the capacity of ecosystems for climate mitigation and adaptation to climate changes. Newbold et al. (2016) assessed that land use pressures have reduced grassland biodiversity beyond the planetary boundary suggested as a safe limit, indicating that continuing biodiversity loss will undermine efforts toward long-term sustainable development. The IPBES and IPCC reports call for the development of a mix of policy instruments to motivate private and public decision-makers to implement both climate mitigation and biodiversity conservation measures while maintaining high levels of other ecosystem services.

In addition to their importance for biodiversity and for carbon sequestration, grasslands can also play an important role in food security, especially in the Nordic countries, where cultivation is constrained by climatic factors. While the potential for carbon sequestration in grasslands worldwide largely lies in the restoration of degraded grasslands, large areas of Nordic grasslands remain, although they are under pressure from forest regrowth due to lack of grazing to maintain the open grasslands. In this context, this essay draws attention to the importance of grasslands for soil carbon sequestration and biodiversity conservation, as well as for other ecosystem services, such as grazing resources. Specifically, we discuss whether grassland management for biodiversity and

food production may have the co-benefit of climate mitigation through increased soil carbon sequestration. Since regional differences are important for these complex interactions, we will focus the discussion on Europe and especially Nordic grasslands.

2 Grasslands in Europe

Grasslands are herbaceous vegetation types mostly dominated by grasses (*Poaceae*) or other graminoids (*Cyperaceae*, *Juncaceae*) and with a relatively dense vegetation cover (>25 % surface cover) (Dengler et al. 2014). They are often transitional ecosystems maintained at a stable successional stage by disturbances such as grazing, mowing, or fire (Norderhaug et al. 1999; Fuhlendorf et al. 2006; Ratajczak et al. 2014; Scasta et al. 2016). In Europe, grasslands can be classified as natural, semi-natural, or cultivated. Natural grasslands occur in areas where forest development is not possible due to dry or cold conditions or other factors that prevent tree growth. However, such areas are rare in Europe today, with small pockets of “genuine” steppe found in Ukraine and Romania (Emanuelsson et al. 2009). On the other hand, semi-natural and cultivated grasslands have been developed by human management in naturally forested areas. These grasslands require human intervention to prevent overgrowth by trees. Semi-natural grasslands, which were created through low-intensity grazing or hay production, are often species-rich. Some of the semi-dry and base-rich semi-natural grasslands found in Europe hold the world record for vascular plant-species richness on small scale (e. g. 89 species on 1m²) (Wilson et al. 2012; Habel et al. 2013). However, both natural and semi-natural grasslands in Europe have increasingly been converted to cultivated grasslands that are fertilized and intensively managed for grazing or for hay production, resulting in a loss of biodiversity (Bignal and McCracken 1996; Pykälä 2007).

3 Characteristics of grasslands with high potential for soil carbon sequestration

Certain grassland characteristics and management strategies may be particularly important for promoting or reducing soil carbon sequestration potentials. Here, we present some key points that summarize these characteristics and strategies, with a call for more research.

3.1 Highly developed root systems and high belowground biomass allocation

Grassland plant species are adapted to grazing and mowing through a low apical growth point, high allocation of biomass belowground, and a root/shoot ratio that is about 10 times higher than in forests (Jackson et al. 1996). The depth distribution of roots is a critical factor in the storage and stabilization of root carbon in the soil (Poirier et al. 2018). In fact, grassland soil carbon is often largely composed of root-derived carbon (Martinez et al. 2016; Yang et al. 2021). As the mean residence time of root carbon increases with soil depth (Rasse et al. 2005), deeper-rooted plants may be particularly effective at sequestering carbon in the soil (Sosa-Hernández et al. 2019). In addition, the organic matter in the subsoil contains more microbial-derived compounds than topsoil organic matter, which can be integrated into mineral-associated soil organic fractions with especially long mean residence (Rumpel and Kögel-Knabner 2011; Sosa-Hernández et al. 2019).

3.2 High aboveground and belowground biodiversity

Increasing plant species diversity can increase root biomass and root exudates, potentially promoting microbial necromass contribution to soil organic storage (Bai and Cotrufo 2022). In a 4-year managed grassland experiment with a biodiversity gradient ranging from one to 60 species, it was shown that carbon storage increased with sown species richness (Steinbeiss et al. 2008). Additionally, manipulation of species richness in a 19-year grassland experiment showed that increasing plant species richness enhanced carbon storage (Pastore et al. 2021). Aboveground biodiversity influences belowground diversity (Chen et al. 2018), and vice versa. For example, arbuscular mycorrhizal fungal diversity can increase plant diversity in grasslands, although the mycorrhizal community composition seems to be more important than the richness per se (Kozioł and Bever 2019). However, the knowledge about the relations between aboveground and belowground biodiversity remains limited (De Deyn et al. 2005). Belowground biodiversity is however of high importance for carbon and nutrient cycling processes, and diversity across groups of organisms plays a decisive role in fully maintaining these processes (de Graaff et al. 2015).

3.3 Mycorrhiza

Most grassland plants live in symbiosis with arbuscular mycorrhizal fungi, which depend on recent plant photosynthates for their growth and activity. Arbuscular mycorrhizal fungi colonize plant roots and form soil mycelia that scavenge for nutrients in larger soil volumes and smaller soil pores than roots (Sosa-Hernández et al. 2019). Some of the nutrients, particularly nitrogen, phosphorus, and sulfur, taken up by arbuscular mycorrhizal fungi are transported to the roots and may stimulate plant growth. Meanwhile, some nutrients are immobilized, along with carbon, in living and dead soil mycelium (Giovannetti et al. 2017). Therefore, the mycorrhizal mycelium provides a direct pathway for recently fixed carbon into the soil matrix. This is in contrast to free-living saprotrophic organisms that rely on carbon from decomposing organic material for their metabolism and thus elicit a net loss of soil carbon. Arbuscular mycorrhizal mycelium can also promote aggregate stability, leading to improved soil structure and carbon sequestration (Lehmann et al. 2017). However, even though arbuscular mycorrhizal fungi lack the ability to decompose complex organic matter (Tisserant et al. 2013), they may still enhance decomposition and nutrient mineralization by stimulating saprotrophic organisms (Jansa et al. 2019). Therefore, the net effects of arbuscular mycorrhiza on soil carbon stocks in different types of grasslands are not yet well understood, although they likely have a positive long-term effect on soil carbon stocks, particularly in the subsoil (Sosa-Hernández et al. 2019, Lehmann et al. 2017).

3.4 Optimal grazing pressure

Grazing stimulates the accumulation of soil organic carbon by promoting root growth and turnover and by defoliation, which provides large inputs of organic matter to the soil (Ziter and MacDougall 2013). For example, Allard et al. (2007) found that 3 years of grazing in a French upland semi-natural grassland promoted the carbon sink, regardless of grazing intensity and fertilization. However, a meta-analysis in 17 grasslands

worldwide showed that the grazing effect on carbon storage depends on grazing intensity, environmental conditions, such as precipitation and soil texture, and the types of grasses (classified as grasslands dominated by C_3 grasses, C_4 grasses, or a mixture of both) (McSherry and Ritchie 2013). C_3 and C_4 grasses have different photosynthetic pathways, and grazing impacts their root structure differently. European grasslands are predominantly C_3 grasslands (de Deus et al. 2021; Still et al. 2003). In general, soil carbon sequestration is positively correlated with grazing intensity in C_4 grasslands and negatively in C_3 grasslands (Garnett et al. 2017). Exploring interactions between grass type and grazing McSherry and Ritchie (2013) found that for C_3 grasses, grazing had a positive effect on soil organic carbon only at light grazing intensities, and this effect became negative at moderate to heavy grazing intensities.

While light to moderate intensity grazing is more likely to maintain soil carbon stocks and has greater potential to enhance sequestration, heavy grazing reduces plant growth, destroys the plant cover, and causes soil carbon losses (Garnett et al. 2017). Studies indicate that moderate-intensity grazing promotes soil carbon sequestration more than no grazing or high-intensity grazing (Ziter and MacDougall 2013; Chang et al. 2016). A nationwide survey of permanent grasslands in England confirmed that moderate management (grazing and fertilizing) resulted in the greatest soil carbon stocks (Ward et al. 2016). Some studies even conclude that carbon sequestration in grazed grasslands partly, or under specific conditions, entirely offsets greenhouse gas (GHG) emissions from livestock (Bellarby et al. 2013; Godde et al. 2020). Recent studies indicate that optimal grazing practices, such as optimal timing and numbers of livestock and moving ruminants between pastures to avoid overgrazing, may facilitate soil carbon sequestration (Stanley et al. 2018; Teague and Kreuter 2020). Moving grazing animals between pastures is a key aspect of sustainable management. Without fencing, grazing animals naturally move between pastures. Observations of dairy cows on mountain summer pastures in Norway indicate that they cover large areas during grazing, and traditional cow breeds moved longer distances (Hessle et al. 2014); this prevents overgrazing. The contrasting results on grazing and soil carbon sequestration potential illustrate the importance of considering the specific agro-ecological conditions (Garnett et al. 2017).

3.5 Sensitivity to physical soil disturbance and fertilization

Ploughing and tilling practices to revitalize grasslands and increase yield typically result in loss of soil carbon. According to Jastrow et al. (2007), reducing soil organic carbon turnover by minimizing soil disturbance through tillage or erosion is crucial for enhancing carbon sequestration. Converting of grasslands into cropland also results in loss of soil carbon. Poeplau et al. (2011) showed that 17 years after grasslands were converted to annual croplands, more than a third of their soil organic carbon stocks had been lost. In contrast, establishing grasslands on former croplands led to continuously increasing soil organic carbon for more than 100 years and a higher gain in soil organic carbon than when converting cropland to forest. This finding was confirmed by a literature review conducted by Conant et al. (2017). Moreover, fertilization can increase plant biomass and may therefore further promote soil carbon sequestration in non-tilled, permanent grasslands (Karlton et al. 2010). In a study of chronic nitrogen fertilization of permanent grasslands, Cenini et al. (2015) found that heavy (organo-mineral) but not light (organic) soil density fractions showed increased soil carbon sequestration. However, inputs of nitrogen and phosphorous to grasslands lead to changes in the taxonomic and functional composition of soil

microbial communities (Leff et al. 2015), which can influence the entire belowground ecosystem. Hence, practices that affect soil organisms, such as soil disturbance and fertilization, are likely to also affect soil organic matter dynamics, since these are tightly linked to the nutrient and carbon needs of the soil organisms (Janzen 2006).

4 Potential for carbon sequestration in grasslands

The characteristics that may support the potential of grasslands for soil carbon sequestration appear to be closely aligned with those of semi-natural grasslands, and to a large extent, natural grasslands. This suggests that semi-natural grasslands may have a good potential for carbon sequestration and potentially a higher soil carbon content than cultivated grasslands, although this has not been clearly documented. Many scientific papers discussing soil carbon sequestration and carbon stocks in grasslands do not specify the type of grassland studied. Permanent grasslands can be either improved, fertilized grasslands or semi-natural grasslands.

Semi-natural grasslands are the result of low-intensity agricultural practices that have been in place for a long time (Dengler et al. 2014). Over this time, they may have developed considerable root systems and built up large soil carbon stocks, while the carbon sequestration rate has diminished successively and approached an equilibrium (Garnett et al. 2017). This equilibrium may be maintained as long as the management remains unchanged (Garnett et al. 2017). On the other hand, the management of cultivated grasslands may vary from year to year, giving less opportunity for root system development. However, permanent improved grasslands may have better developed root systems. Further, most of the plant species in semi-natural grasslands are associated with arbuscular mycorrhizal fungi (Norwegian Biodiversity Information Centre 2019) which probably improve soil carbon sequestration (Sosa-Hernández et al. 2019). In contrast, cultivated grasslands tend to have lower abundance of mycorrhizal fungi (Schnoor et al. 2011; Bowles et al. 2016, House and Bever 2018), although we do not know at which timescales permanent grasslands may reestablish biodiversity and characteristics of semi-natural grasslands.

In Europe, most remaining semi-natural grasslands are grazed. Several studies have shown that overgrazing causes loss of soil carbon stock and that light to moderate grazing could maintain soil carbon stocks and have a greater potential for soil carbon sequestration than heavy or too light grazing (Garnett et al. 2017). A study of sheep grazing on low-alpine grasslands in Norway found that 7 years of grazing had little impact on soil carbon stocks, with high sheep stocking rates causing a small decrease and low stocking rates causing no or a slight increase (Martinsen et al. 2011). Furthermore, another study of lightly grazed grassland, *Empetrum*-dominated heath, and *Salix*-shrub-dominated vegetation in the Norwegian mountains showed that grassland had the largest total ecosystem carbon pool (Sørensen et al. 2018). In alpine regions, semi-natural and natural grasslands may have quite stable and substantial stocks of soil carbon (Castano et al. 2022), but at the same time, may have limited potential for additional, short-term sequestration because soil processes are slow. However, alpine grasslands store more of the soil carbon in particulate form, often forming an organic topsoil, while temperate grasslands store relatively more carbon in mineral-associated organic matter in the subsoil. An interesting avenue for further research thus is to unravel mechanisms of carbon storage in organic topsoils versus mineral subsoils across climatic gradients, since minerals can get saturated with

organic matter, while carbon storage in particulate organic matter has no such limit (Bai and Cotrufo 2022).

Cultivated grasslands are more productive than semi-natural grasslands due to their higher nitrogen and phosphorous content, which allows for increased grazing animals and stimulation of aboveground biomass production and root growth. While fertilization of cultivated grasslands generally increases soil carbon stocks by stimulating plant growth, fertilization of semi-natural grasslands reduces biodiversity (Emanuelsson et al. 2009). As semi-natural grasslands are of high importance for biodiversity, management strategies aimed at increasing soil carbon stocks, such as fertilization, may not be suitable for sustainable management of these ecosystems.

5 Climate policy in synergy with other sustainable development goals: biodiversity and food production

Semi-natural grasslands play an important role in maintaining European biodiversity (Pykälä 2007). However, in Western Europe, the area of this type of ecosystem has significantly decreased due to factors such as intensification of agriculture, abandonment of traditional agricultural practices, and land conversion. This trend has resulted in long red lists of endangered species of plants, fungi, insects, birds, and other animals that depend on semi-natural grasslands. In Norway, for instance, 29% of the threatened species live in semi-natural habitats of grasslands and coastal heathlands (Norwegian Biodiversity Information Centre 2021), while in Sweden, semi-natural habitats are critical for 34% of red-listed species (SLU Swedish Species Information Centre 2020), and in Finland, for 24% (Finland's Environmental Administration 2019). The trends for grassland biodiversity are negative, and more and more species are being red listed (Jakobsson and Pedersen (eds.) 2020; IPBES 2019), emphasizing the need to preserve remaining semi-natural grasslands to prevent further loss of biodiversity.

Furthermore, maintaining semi-natural grasslands can help preserve existing carbon stocks in the ground (Garnett et al. 2017), keep the landscape open, and contribute to a higher albedo effect. Although albedo is not included in the reporting guidelines of the United Nations Framework Convention on Climate Change (UNFCCC), Bonan et al. (2008) emphasized that albedo can have large effects on climate feedbacks at northern latitudes. Compared to grasslands, boreal forests have lower albedo and absorb more solar radiation, contributing to temperature increase in the atmosphere. Therefore, understanding the importance of grasslands *versus* forests for albedo in the context of climate mitigation, especially in the Nordic countries and other boreal regions, is essential.

Semi-natural grasslands need to be recognized for their multiple functions. To balance climate mitigation and grassland biodiversity preservation, Burrascano et al. (2016) suggest aligning decision-making across policy sectors, focusing on a range of ecosystem services and biodiversity issues, and valuing extensively managed (low-intensity) ecosystems for their multiple functions. The current agricultural policy in the EU lacks attention to the management of permanent grasslands, which is crucial for their environmental benefits (Pe'er et al. 2014).

The roles of grasslands differ among regions. In northern and alpine regions of Europe, where crop production is constrained by climatic and ecological conditions, grasslands play an important role in meat and dairy production (Nordic councils of Ministers 2017). Grazing ruminants can utilize grasslands which cannot be cultivated but still can contribute

to food for humans. In the Nordic countries, particularly in mountainous Norway where only 3% of the total land is cultivated, the areas of rangeland with semi-natural and natural grasslands are vital for improving the self-sufficiency and ensuring more reliable food security (Nordic Council of Ministers 2017). The World Resources Institute (2019) emphasizes the responsibility of all countries to utilize their possibilities for food production to secure a sustainable food future.

IPBES (2019) highlights that activities aimed at mitigating climate change carry the risk of negative side effects for biodiversity and food security. Afforestation, an important aspect of EU climate policy, is one such activity that has been criticized for loss of biodiversity, particularly when semi-natural grasslands are afforested (Burrascano et al. 2016; Veldman et al. 2019). In addition, afforestation does not necessarily lead to higher soil carbon stocks (Poepflau et al. 2011). A Norwegian study conducted on partially afforested grazed grasslands found that even 50 years after planting, afforestation did not result in higher carbon sequestration than in the grassland (Strand et al. 2020).

6 Research needs

According to the reporting guidelines of the United Nations Framework Convention on Climate Change (UNFCCC), national inventories covering emissions and removals of GHGs from agriculture, forestry, and other land use (volume 4) should cover livestock and manure management (chapter 10) as well as carbon sequestration in grassland (chapter 6) (IPCC 2006). However, only grasslands classified as “managed land” by being fertilized, tilled, and sown are included in the national inventory reports, excluding semi-natural grasslands, even though these must be grazed or mown to be maintained. This presents a problem as semi-natural grasslands cover vast areas and may store large amounts of soil carbon. Yet grazing ruminants are classified as only contributing to GHG emissions, due to the lack of knowledge of soil carbon sequestration in grasslands, especially semi-natural grasslands. To enhance the knowledge basis for climate policy, more research is needed on grassland soil carbon, paying attention to regional differences in the complex interactions of grassland ecology and agricultural practices.

In a global perspective, it is especially important to acquire more knowledge on soil carbon in semi-natural grasslands, as attention has been drawn to grazing when the United Nations declared 2026 as the International Year of Rangelands and Pastoralists (United Nations 2022). It was then emphasized that keeping the vast areas of rangelands, with their large soil carbon sinks, managed under sustainable grazing practices, is essential for climate change mitigation (United Nations 2022).

Several measurement challenges exist when it comes to grassland soil carbon sequestration. One of these challenges is the spatial variability of soil carbon stocks. To effectively implement policies aimed at increasing soil organic carbon at a large scale and integrating soil organic carbon into national and international climate reporting, a combination of direct measurements and modeling is necessary (Smith et al. 2019). Standard protocols for measuring and monitoring soil carbon stocks would also facilitate comparisons between countries, regions, and land use types. However, it is essential to consider differences in climatic, ecological, and agricultural conditions when making such comparisons.

Assessing the soil carbon sequestration potential of grasslands is challenging due to the complexity of interactions between various factors such as vegetation characteristics, belowground biodiversity, grazing intensity, and management (Teague and Kreuter 2020).

Nonetheless, it is essential to gain more knowledge about the climate mitigation potential of grasslands, particularly semi-natural grasslands, to enhance the knowledge base for climate policy. For instance, do semi-natural grasslands have larger carbon stocks in deeper layers than cultivated grasslands? Can their management be optimized to increase carbon sequestration while maintaining their biodiversity? According to Janzen (2006), a better understanding of carbon flows in the soil and of the roles of soil microbes and fauna is necessary to balance the different ecosystem services expected from grasslands. It is particularly crucial to balance the need for carbon stabilization in the soil (i.e., climate mitigation) with the need for recirculation of dead organic matter to release nutrients to support plant growth (Janzen 2006).

7 Conclusion

We must take a fresh approach to climate policy and consider the potential of grasslands as vital carbon sinks. It is essential to focus on finding a balance between policies for climate mitigation and biodiversity management as the loss of biodiversity is as great a threat as climate change (IPBES 2019). Semi-natural grasslands are particularly important for biodiversity in Europe, and in some regions, they also contribute to food production and food security. Therefore, it is crucial to assess the multiple functions of grasslands, especially semi-natural grasslands, to optimize policy measures across sectors. To achieve this, more research is required to understand the potential of carbon sequestration in grasslands, particularly semi-natural grasslands, in different regions, so we can manage them optimally for climate mitigation and preservation of biodiversity, as well as other critical ecosystem services.

Acknowledgements The authors thank the Research Council of Norway (RCN) for financial support for the project CLIMATE-LAND (RCN-235560), and the work of Iulie Aslaksen received additional support from PLATON (RCN-295789).

Funding Open access funding provided by Statistics Norway

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Aguilera E et al (2021) Greenhouse gas emissions from Mediterranean agriculture: evidence of unbalanced research efforts and knowledge gaps. *Global Env Change* 69:102319
- Allard V, Soussana JF, Falcimagne R, Berbigier P, Bonnefond JM, Ceschia E, D'hour C, Hénault P, Laville P, Martin C, Pinarès-Patino C (2007) The role of grazing management for the net biome productivity and greenhouse gas budget (CO₂, N₂O and CH₄) of semi-natural grassland. *Agric Ecosyst Environ* 121:47–58
- Bai Y, Cotrufo F (2022) Grassland soil carbon sequestration: Current understanding, challenges, and solutions. *Science* 377(6606):603–608. <https://doi.org/10.1126/science.abo2380>

- Bellarby J, Tirado R, Leip A, Weiss F, Lesschen JP, Smith P (2013) Livestock greenhouse gas emissions and mitigation potential in Europe. *Glob Chang Biol* 19:3–18
- Bignal EM, McCracken DI (1996) Low-intensity farming systems in the conservation of the countryside. *J Appl Ecol* 33:413–424
- Bonan GB (2008) Forests and climate change: forcings, feedbacks, and the climate benefits of forests. *Science* 320(5882):1444–1449. <https://doi.org/10.1126/science.1155121>
- Bowles TM, Jackson LE, Loehner M, Cavagnaro TR (2016) Ecological intensification and arbuscular mycorrhizas : a meta-analysis of tillage and cover crop effects. *J Appl Ecol* 54:1785–1793. <https://doi.org/10.1111/1365-2664.12815>
- Burrascano S, Chytr M, Kuemmerle T, Giarrizzo E, Luyssaert S, Sabatini FM, Blasi C (2016) Current European policies are unlikely to jointly foster carbon sequestration and protect biodiversity. *Biol Conserv* 201:370–376
- Castano C, Hallin S, Egelkraut D, Lindahl BD, Olofsson J, Clemmensen KE (2022) Contrasting plant-soil-microbial feedbacks stabilize vegetation types and uncouple topsoil C and N stocks across a subarctic-alpine landscape. *New Phytol*. <https://doi.org/10.1111/nph.18679>
- Cenini VL, Fornara DA, McMullan G, Ternan N, Lajtha K, Crawley MJ (2015) Chronic nitrogen fertilization and carbon sequestration in grassland soils: evidence of a microbial enzyme link. *Biogeochem* 126:301–313. <https://doi.org/10.1007/s10533-015-0157-5>
- Chang J, Ciais P, Viomy N, Vuichard N, Sultan B, Soussana JF (2015) The greenhouse gas balance of European grasslands. *Glob Chang Biol* 21:3748–3761
- Chang J et al (2016) Combining livestock production information. *Biogeosci* 13:3757–3776. <https://doi.org/10.5194/bg-13-3757-2016>
- Chen S, Wang W, Xu W, Wang Y, Wan H, Chen D, Tang Z, Tang X, Zhou G, Xie Z, Zhou D, Shanguan Z, Huang J, He JS, Wang Y, Sheng J, Tang L, Li X, Dong M, Wu Y, Wang Q, Wang Z, Wu J, F. Chapin S III, Bai Y (2018) Plant diversity enhances productivity and soil carbon storage. *PNAS* 115(16): 4027–4032. <https://doi.org/10.1073/pnas.1700298114>
- Conant RT, Cerri CEP, Osborne BB, Paustian K (2017) Grassland management impacts on soil carbon stocks: a new synthesis. *Ecol Appl* 27:662–668. <https://doi.org/10.1002/eap.1473>
- de Deus Vidal J Jr, le Roux PC, Johnson SD, te Beest M, Clark VR (2021) Beyond the tree-line: the C3-C4 “grass-line” can track global change in the world’s grassy mountain systems. *Front Ecol Evol* 9:760118. <https://doi.org/10.3389/fevo.2021.760118>
- De Deyn GB, Van der Putten WH (2005) Linking aboveground and belowground diversity. *Trends Ecol Evol* 20:625–633. <https://doi.org/10.1016/j.tree.2005.08.009>
- de Graaff M-A, Adkins J, Kardol P, Throop HL (2015) A meta-analysis of soil biodiversity impacts on the carbon cycle. *Soil* 1:257–271. www.soil-journal.net/1/257/2015/. <https://doi.org/10.5194/soil-1-257-2015>
- Dengler J, Janišová M, Török P, Wellstein C (2014) Biodiversity of Palearctic grasslands: a synthesis. *Agric Ecosyst Environ* 182:1–14
- Emanuelsson U, Arding M, Petersson M (2009) The rural landscapes of Europe: how man has shaped European nature. *Formas, Stockholm*
- Finland’s Environmental Administration (2019) Suomen lajien uhanalaisuus (Finlands rödlista 2019) - Punainen kirja 2019. <https://www.ymparisto.fi/en/nature-waters-and-seas/natural-diversity/diversity-species/threat-level-assessments-species>. Accessed 28 Mar 2023
- FAO (2017) Soil organic carbon: The hidden potential. Food and Agriculture Organization of the United Nations <https://www.fao.org/documents/card/en/c/ed16dbf7-b777-4d07-8790-798604fd490a/>. Accessed 28 Mar 2023
- Fuhlendorf SD, Harrell WC, Engle MD, Hamilton RG, Davis CA, Leslie jr. DM. (2006) Should heterogeneity be the basis for conservation? Grassland bird response to fire and grazing. *Ecol Appl* 16(5):1706–1716
- Garnett T, Godde C, Muller A, Rösös E, Smith P, de Boer IJM, zu Ermgassen E, Herrero M, van Middelaar C, Schader C, van Zanten H (2017) Grazed and confused? Ruminating on cattle, grazing systems, methane, nitrous oxide, the soil carbon sequestration question – and what it all means for greenhouse gas emissions. Food Climate Research Network (FCRN), Oxford Martin Programme on the Future of Food Environmental Change Institute, University of Oxford. https://www.oxfordmartin.ox.ac.uk/downloads/reports/fcrn_gnc_report.pdf
- Giovannetti M, Volpe V, Salvioli A, Bonfante P (2017) Fungal and plant tools for the uptake of nutrients in arbuscular mycorrhizas: a molecular view. In: Johnson NC, Gehring C, Jansa J (eds) *Mycorrhizal Mediation of Soil: Fertility, Structure and Carbon Storage*. Elsevier, Amsterdam, pp 107–128
- Godde CM, de Boer IJM, zu Ermgassen E et al (2020) Soil carbon sequestration in grazing systems: managing expectations. *Clim Chang* 161:385–391. <https://doi.org/10.1007/s10584-020-02673-x>
- Habel JC, Dengler J, Janišová M (2013) European grassland ecosystems: threatened hotspots of biodiversity. *Biodivers Conserv* 22:2131

- Hessle A, Dahlström F, Bele B, Norderhaug A, Söderström M (2014) Effects of breed on foraging sites and diets in dairy cows on mountain pasture. *International Journal of Biodiversity Science, Ecosystem Services & Management* 10:334–342. <https://doi.org/10.1080/21513732.2014.968805>
- House GL, Bever JD (2018) Disturbance reduces the differentiation of mycorrhizal fungal communities in grasslands along a precipitation gradient. *Ecol Appl* 28:736–748
- IPBES (2019) Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. In: Díaz S, Settele J, Brondizio ES, Ngo HT, Guèze M, Agard J, Arneth A, Balvanera P, Brauman KA, Butchart SHM, Chan KMA, Garibaldi LA, Ichii K, Liu J, Subramanian SM, Midgley GF, Miloslavich P, Molnár Z, Obura D, Pfaff A, Polasky S, Purvis A, Razzaque J, Reyers B, Chowdhury RR, Shin YJ, Visseren-Hamakers IJ, Willis KJ, Zayas CN (eds.) IPBES secretariat, Bonn, Germany. https://www.ipbes.net/sites/default/files/inline/files/ipbes_global_assessment_report_summary_for_policymakers.pdf
- IPCC (2006) 2006 IPCC Guidelines for National Greenhouse Gas Inventories, volume 4. Agriculture, Forestry and Other Land Use. Intergovernmental Panel on Climate Change (IPCC). <https://www.ipcc-nggip.iges.or.jp/public/2006gl/vol4.html>. Accessed 28 Mar 2023
- IPCC (2019) Summary for Policymakers. In: *Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems*. In: Shukla PR, Skea J, Calvo Buendia E, Masson-Delmotte V, Pörtner H-O, Roberts DC, Zhai P, Slade R, Connors S, van Diemen R, Ferrat M, Haughey E, Luz S, Neogi S, Pathak M, Petzold J, Portugal Pereira J, Vyas P, Huntley E, Kissick K, Belkacemi M, Malley J (eds). <https://doi.org/10.1017/9781009157988.001>
- IPCC (2022) Climate change 2022: Impacts, adaptation and vulnerability. Contribution of working group II to the sixth assessment report of the Intergovernmental Panel on Climate Change. In: Pörtner H-O, Roberts DC, Tignor M, Poloczanska ES, Mintenbeck K, Alegría A, Craig M, Langsdorf S, Lösschke S, Möller V, Okem A, Rama B (eds). Cambridge University Press, Cambridge, UK and New York, NY, USA. <https://doi.org/10.1017/9781009325844>
- Jackson RB, Canadell J, Ehleringer JR, Mooney HA, Sala OE, Schulze ED (1996) A global analysis of root distributions for terrestrial biomes. *Oecologia* 108:389–411
- Jakobsson S, Pedersen B (eds) (2020) The Nature Index for Norway 2020. State and trends for Norwegian biodiversity. NINA Report 1886. Norwegian Institute for Nature Research (NINA). Norwegian Institute for Nature Research (NINA), Trondheim, Norway. <https://brage.nina.no/nina-xmlui/handle/11250/2686068>
- Jansa J, Forczek ST, Rozmoš M, Püschel D, Bukovská P, Hršelová H (2019) Arbuscular mycorrhiza and soil organic nitrogen: network of players and interactions. *Chem and BiolTech in Agric* 6:10
- Janzen HH (2006) The soil carbon dilemma: Shall we hoard it or use it? *Soil Biol Biochem* 38:419–424
- Jastrow JD, Amonette JE, Bailey VL (2007) Mechanisms controlling soil carbon turnover and their potential application for enhancing carbon sequestration. *Clim Chang* 80:5–23. <https://doi.org/10.1007/s10584-006-9178-3>
- Karlton E et al (2010) Inlagring av kol i betesmark. Jordbruksverket Rapport 2010:25 (Report in Swedish on carbon sequestration in semi-natural grasslands)
- Kozioł L, Bever JD (2019) Mycorrhizal feedbacks generate positive frequency dependence accelerating grassland succession. *J Ecol* 107:622–632
- Leff JW, Jones SE, Prober SM, Barberan A, Borer ET, Firn JL, Harpole WS, Hobbie SE, Hofmockel KS, Knops JMH et al (2015) Consistent responses of soil microbial communities to elevated nutrient inputs in grasslands across the globe. *PNAS* 112:10967–10972
- Lehmann A, Leifheit EF, Rillig MC (2017) Mycorrhizas and soil aggregation. In: Johnson NC, Gehring C, Jansa J (eds) *Mycorrhizal Mediation of Soil: Fertility, Structure and Carbon Storage*. Elsevier, Amsterdam, pp 241–262. <https://www.sciencedirect.com/science/article/pii/B9780128043127000140?via%3Dihub>. Accessed 28 Mar 2023
- Lorenz K, Lal R (2018) Carbon sequestration in agricultural ecosystems. Springer
- Martinez C, Alberti G, Cotrufo MF, Magnani F, Zanotelli D, Camin F, Gianelle D, Cescatti A, Rodeghiero M (2016) Belowground carbon allocation patterns as determined by the in-growth soil core ¹³C technique across different ecosystem types. *Geoderma* 263:140–150. <https://doi.org/10.1016/j.geoderma.2015.08.043>
- Martinsen V, Mulder J, Austrheim G, Mysterud A (2011) Carbon storage in low-alpine grassland soils: effects of different grazing intensities of sheep. *Eur J Soil Sci* 62:822–833. <https://doi.org/10.1111/j.1365-2389.2011.01393.x>
- McSherry ME, Ritchie ME (2013) Effects of grazing on grassland soil carbon: a global review. *Glob Chang Biol* 19(5):1347–1357
- Newbold T, Hudson LN, Arnell AP, Contu S, De Palma A, Ferrier S, Hill SL, Hoskins AJ, Lysenko I, Phillips HR, Burton VJ, Chng CW, Emerson S, Gao D, Pask-Hale G, Hutton J, Jung M, Sanchez-Ortiz K,

- Simmons BI et al (2016) Has land use pushed terrestrial biodiversity beyond the planetary boundary? A global assessment *Science* 353(6296):288–291. <https://doi.org/10.1126/science.aaf2201>
- Nordic Council of Ministers (2017) Future Nordic Diets. <https://norden.diva-portal.org/smash/get/diva2:1163192/FULLTEXT01.pdf>. Accessed 28 Mar 2023
- Norderhaug A, Austad I, Hauge I, Kvamme M (red.) (1999) Skjøtselsboka for kulturlandskap og gamle norske kulturmarker. Landbruksforlaget. <https://www.miljodirektoratet.no/globalassets/publikasjoner/dirnat2/attachment/636/1.pdf>. Accessed 28 Mar 2023
- Norwegian Biodiversity Information Centre (2021) Hvor finnes de tuende artene? Norsk rødliste for arter 2021. Artsdatabanken, Trondheim, Norway. <https://artsdatabanken.no/rodliseforarter2021/Resultater>. Accessed 28 Mar 2023
- Norwegian Biodiversity Information Centre (2019) Natur i Norge. T32. Semi-naturlig eng. Artsdatabanken, Trondheim, Norway. https://artsdatabanken.no/Pages/171950/Semi-naturlig_eng. Accessed 28 Mar 2023
- Pastore MA, Hobbie SE, Reich PB (2021) Sensitivity of grassland carbon pools to plant diversity, elevated CO₂, and soil nitrogen addition over 19 years. *PNAS* 118(17):e2016965118. <https://doi.org/10.1073/pnas.2016965118>
- Pe'er G, Dicks LV, Visconti P, Arlettaz R, Baldi A, Benton TG, Collins S, Dieterich M, Gregory RD, Hartig F, Henle K, Hobson PR, Kleijn D, Neumann RK, Robijns T, Schmidt J, Schwartz A, Sutherland WJ, Turbe A et al (2014) EU agricultural reform fails on biodiversity. *Science* 344:1090–1092
- Poepplau C, Don A, Vesterdal L, Leifeld J, van Wesemael B, Schumacher J, Gensior A (2011) Temporal dynamics of soil organic carbon after land-use change in the temperate zone – carbon response functions as a model approach. *Glob Chang Biol* 17:2415–2427. <https://doi.org/10.1111/j.1365-2486.2011.02408.x>
- Poirier V, Roumet C, Munson AD (2018) The root of the matter: Linking root traits and soil organic matter stabilization processes. *Soil Biol Biochem* 120:246–259
- Pykälä J (2007) Maintaining plant species richness by cattle grazing: mesic semi-natural grasslands as focal habitats. Publications in Botany from the University of Helsinki, no. 36, University of Helsinki, Faculty of Biosciences, Helsinki, Finland. <http://urn.fi/URN:ISBN:978-952-10-4205-8>
- Rasse DP, Rumpel C, Dignac MF (2005) Is soil carbon mostly root carbon? Mechanisms for a specific stabilisation. *Plant Soil* 269:341–356
- Ratajczak Z, Nippert JB, Ocheltree TW (2014) Abrupt transition of mesic grassland to shrubland: evidence for thresholds, alternative attractors, and regime shifts. *Ecology* 95(9):2633–2645
- Röös E, Bajzelj B, Smith P, Patel M, Little D, Garnett T (2017) Greedy or needy? Land use and climate impacts of food in 2050 under different livestock futures. *Global Env Change* 47:1–12
- Rumpel C, Kögel-Knabner I (2011) Deep soil organic matter—a key but poorly understood component of terrestrial C cycle. *Plant Soil* 338:143–158. <https://doi.org/10.1007/s11104-010-0391-5>
- Scasta JD, Duchardt C, Engle DM, Miller JR, Debinski DM, Harr RN (2016) Constraints to restoring fire and grazing ecological processes to optimize grassland vegetation structural diversity. *Ecol Eng* 95:865–875. <https://doi.org/10.1016/j.ecoleng.2016.06.096>
- Schnoor TK, Lekberg Y, Rosendahl S, Olsson PA (2011) Mechanical soil disturbance as a determinant of arbuscular mycorrhizal fungal communities in semi-natural grassland. *Mycorrhiza* 21:211–220
- SLU Swedish Species Information Centre (2020) SLU Artdatabanken. Rödlistade arter i Sverige 2020. SLU, Uppsala <https://www.artdatabanken.se/publikationer/bestall-publikationer/bestall-rodlista-2020/>. Accessed 28 Mar 2023
- Smith P et al (2019) How to measure, report and verify soil carbon change to realize the potential of soil carbon sequestration for atmospheric greenhouse gas removal. *Glob Chang Biol* 26:219–241. <https://doi.org/10.1111/gcb.14815>
- Sosa-Hernández MA, Leifheit EF, Ingrassia R, Rillig MC (2019) Subsoil arbuscular mycorrhizal fungi for sustainability and climate-smart agriculture: a solution right under our feet? *Front Microbiol* 10:744
- Stanley PL, Rowntree JE, Beede DK, MS DL, Hamm MW (2018) Impacts of soil carbon sequestration on life cycle greenhouse gas emissions in Midwestern USA beef finishing systems. *Agric Syst* 162:249–258
- Steinbeiss S, Beßler H, Engels C, Temperton VM, Buchmann N, Roscher C, Kreuziger Y, Baade J, Habekost M, Gleixner G (2008) Plant diversity positively affects short-term soil carbon storage in experimental grasslands. *Glob Chang Biol* 14:2937–2949
- Still CJ, Berry JA, Collatz GJ, RS DF (2003) Global distribution of C₃ and C₄ vegetation: Carbon cycle implications. *Global Biogeochem Cycles* 17(1):1006. <https://doi.org/10.1029/2001GB001807>
- Strand LT, Fjellstad W, Jackson-Blake L, De Wit HA (2020) Afforestation of a pasture in Norway did not result in higher soil carbon, 50 years after planting. *Landsc Urban Plan* 207. <https://doi.org/10.1016/j.landurbplan.2020.104007>
- Sørensen MV, Strimbeck R, Nystuen KO et al (2018) Draining the pool? Carbon storage and fluxes in three alpine plant communities. *Ecosystems* 21:316–330. <https://doi.org/10.1007/s10021-017-0158-4>

- Teague R, Kreuter U (2020) Managing grazing to restore soil health, ecosystem function, and ecosystem services. *Frontiers in Sust Food Systems* 4:157. <https://doi.org/10.3389/fsufs.2020.534187>
- Tisserant E, Malbreil M, Kuo A, Kohler A, Symeonidi A, Balestrini R, Charron P, Duensing N, Frei dit Frey N, Gianinazzi-Pearson V, Gilbert LB, Handa Y, Herr JR, Hijri M, Koul R, Kawaguchi M, Krajinski F, Lammers PJ, Masclaux FG et al (2013) Genome of an arbuscular mycorrhizal fungus provides insight into the oldest plant symbiosis. *PNAS* 110:20117–20122
- United Nations (2022) International Year of Rangelands and Pastoralists (IYRP) 2026. IYPR resolution approved by the General Assembly. 15 March 2022. https://iyrp.info/sites/iyrp.org/files/IYRP%20adopted_Res_76-253_0.pdf. Accessed 28 Mar 2023
- Veldman JW, Aleman JC, Alvarado ST, Anderson TM, Archibald S, Bond WJ, Boutton TW, Buchmann N, Buisson E, Canadell JG, de Sá DM, Diaz-Toribio MH, Durigan G, Ewel JJ, Fernandes GW, Fidelis A, Fleischman F, Good SP, Griffith DM et al (2019) Comment on “The global tree restoration potential”. *Science* 366 (6463):eaay7976. <https://doi.org/10.1126/science.aay7976>
- Ward SE et al (2016) Legacy effects of grassland management on soil carbon to depth. *Glob Chang Biol* 22:2929–2938
- Wilson JB, Peet RK, Dengler J, Pärtel M (2012) Plant species richness: the world records. *J Veg Sci* 23:796–802
- World Resources Institute (2019) Creating a sustainable food future. World Resources Institute, Washington DC, USA. <https://research.wri.org/wrr-food>. Accessed 28 Mar 2023
- Yang X, Wang B, An S (2021) Root derived C rather than root biomass contributes to the soil organic carbon sequestration in grassland soils with different fencing years. *Plant Soil* 469:161–172. <https://doi.org/10.1007/s11104-021-05144-z>
- Ziter C, AS MD (2013) Nutrients and defoliation increase soil carbon inputs in grassland. *Ecology* 94:106–116

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.