

Bees increase seed set of wild plants while the proportion of arable land has a variable effect on pollination in European agricultural landscapes

Lina Herbertsson^{1,a,*}, Johan Ekroos^{1,b}, Matthias Albrecht², Ignasi Bartomeus³, Péter Batáry⁴, Riccardo Bommarco⁵, Paul Caplat^{1,6}, Tim Diekötter⁷, Jenny M. Eikestam¹, Martin H. Entling⁸, Sunniva Farbu¹, Nina Farwig⁹, Juan P. Gonzalez-Varo^{3,c}, Annika L. Hass^{10,23}, Andrea Holzschuh¹¹, Sebastian Hopfenmüller^{11,d}, Anna Jakobsson¹², Birgit Jauker¹³, Anikó Kovács-Hostyánszki¹⁴, Wera Kleve⁵, William E. Kunin¹⁵, Sandra A.M. Lindström^{5,16,17}, Sarah Mullen¹⁸, Erik Öckinger⁵, Theodora Petanidou¹⁹, Simon G. Potts²⁰, Eileen F. Power¹⁸, Maj Rundlöf¹⁶, Kathrin Seibel¹, Virve Söber²¹, Annika Söderman¹, Ingolf Steffan-Dewenter¹¹, Jane C. Stout¹⁸, Tiit Teder^{21,22}, Teja Tscharntke²³ & Henrik G. Smith^{1,16}

¹Centre for Environmental and Climate Science, Lund University, Lund, Sweden

²Agroscope, Agroecology and Environment, Zürich, Switzerland

³Estación Biológica de Doñana EBD-CSIC, Department of Integrative Ecology, Sevilla, Spain

⁴Centre for Ecological Research, Institute of Ecology and Botany, Lendület Landscape and Conservation Ecology, Vácrtót, Hungary

⁵Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden

⁶School of Biological Sciences, Queen's University, Belfast, UK

⁷Institute for Natural Resource Conservation, Department of Landscape Ecology, Kiel University, Kiel, Germany

⁸IES Landau, Institute for Environmental Sciences, University of Koblenz-Landau, Landau, Germany

⁹Department of Biology, Conservation Ecology, Philipps-Universität Marburg, Marburg, Germany

¹⁰Functional Agrobiodiversity, University of Göttingen, Göttingen, Germany

¹¹Department of Animal Ecology and Tropical Biology, Biocenter, University of Würzburg, Würzburg, Germany

¹²Högskolan Väst, Trollhättan, Sweden

¹³Department of Animal Ecology, Justus Liebig University Giessen, Giessen, Germany

¹⁴Lendület Ecosystem Services Research Group, Institute of Ecology and Botany, Centre for Ecological Research, Vácrtót, Hungary

¹⁵School of Biology, University of Leeds, Leeds, UK

¹⁶Department of Biology, Lund University, Lund, Sweden

¹⁷The Swedish Rural Economy and Agricultural Society, Kristianstad, Sweden

¹⁸School of Natural Sciences, Trinity College Dublin, Dublin, Republic of Ireland

¹⁹Department of Geography, University of the Aegean, Mytilene, Greece

²⁰Centre for Agri-Environmental Research, School of Agriculture, Policy and Development, University of Reading, Reading, UK

²¹Department of Zoology, Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia

²²Department of Ecology, Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Prague, Czech Republic

²³Agroecology, University of Göttingen, Göttingen, Germany

Current affiliations

^aDepartment of Biology, Lund University, Lund, Sweden

^bPlant Production Sciences, Department of Agricultural Sciences, University of Helsinki, Helsinki, Finland

^cDepartamento de Biología, IVAGRO, Universidad de Cádiz, Puerto Real, Spain

^dEvolutionary Ecology and Conservation Genomics, Ulm University, Ulm, Germany

*Corresponding author: lina.herbertsson@biol.lu.se

© 2021 Lina Herbertsson, Johan Ekroos, Matthias Albrecht, Ignasi Bartomeus, Péter Batáry, Riccardo Bommarco, Paul Caplat, Tim Diekötter, Jenny M. Eikestam, Martin H. Entling, Sunniva Farbu, Nina Farwig, Juan P. Gonzalez-Varo, Annika L. Hass, Andrea Holzschuh, Sebastian Hopfenmüller, Anna Jakobsson, Birgit Jauker, Anikó Kovács-Hostyánszki, Wera Kleve, William E. Kunin, Sandra A.M. Lindström, Sarah Mullen, Erik Öckinger, Theodora Petanidou, Simon G. Potts, Eileen F. Power, Maj Rundlöf, Kathrin Seibe, Virve Söber, Annika Söderman, Ingolf Steffan-Dewenter, Jane C. Stout, Tiit Teder, Teja Tscharntke, Henrik G. Smith.

This article is published and distributed in Open Access under the terms of the [Creative Commons Attribution License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution, and reproduction in any medium, provided the original work (author and source) is properly cited.

Plant Ecology and Evolution is published by Meise Botanic Garden and Royal Botanical Society of Belgium

ISSN: 2032-3913 (print) – 2032-3921 (online)

Background and aims – Agricultural intensification and loss of farmland heterogeneity have contributed to population declines of wild bees and other pollinators, which may have caused subsequent declines in insect-pollinated wild plants.

Material and methods – Using data from 37 studies on 22 pollinator-dependent wild plant species across Europe, we investigated whether flower visitation and seed set of insect-pollinated plants decline with an increasing proportion of arable land within 1 km.

Key results – Seed set increased with increasing flower visitation by bees, most of which were wild bees, but not with increasing flower visitation by other insects. Increasing proportion of arable land had a strongly variable effect on seed set and flower visitation by bees across studies.

Conclusion – Factors such as landscape configuration, local habitat quality, and temporally changing resource availability (e.g. due to mass-flowering crops or honey bee hives) could have modified the effect of arable land on pollination. While our results highlight that the persistence of wild bees is crucial to maintain plant diversity, we also show that pollen limitation due to declining bee populations in homogenized agricultural landscapes is not a universal driver causing parallel losses of bees and insect-pollinated plants.

Keywords – Habitat loss; landscape complexity; landscape simplification; pollination; pollinating insects; semi-natural.

INTRODUCTION

Agricultural intensification threatens the richness and abundance of vascular plants in European agricultural landscapes, likely because of a combination of drivers such as nitrogen deposition, weed control, and the loss and degradation of semi-natural habitats (Robinson & Sutherland 2002; Bobbink et al. 2010; Bilz et al. 2011).

Although plant communities are strongly influenced by the availability of water, light, and minerals, which determine the degree of stress and interspecific competition (e.g. Theodose & Bowman 1997; Qi et al. 2018), the regeneration of plants in a community is also affected by seed production (Turnbull et al. 2000), which in turn depends on pollen transfer (Burd 1994). Consequently, loss of suitable pollinators can distort plant regeneration and reshape plant communities (Fontaine et al. 2005). Thus, because about 90% of the flowering plant species are pollinated by insects (Ollerton et al. 2011), observed declines of pollinators, such as butterflies (van Swaay et al. 2006), bees (Scheper et al. 2014), and hoverflies (Hallmann et al. 2021), could exacerbate the threat to plant populations (Clough et al. 2014; Papanikolaou et al. 2017).

A major driver of the pollinator decline is agricultural intensification, which is associated with loss of habitat and increased exposure to agrochemicals (IPBES 2016). In agricultural landscapes where semi-natural habitats have been converted to arable land (hereafter: homogeneous landscapes), pollinators can suffer from a lack of forage plants (Wallisdeevries et al. 2012; Scheper et al. 2014). Depending on their specific needs, groups of pollinators, such as bees, butterflies, and hoverflies, may also suffer from lack of nesting habitats (Persson et al. 2015), larval host plants (Kuussaari et al. 2007), and larval microhabitats (Power et al. 2016). In addition to high-quality habitats (e.g. semi-natural habitats and low-intensive permanent grasslands), pollinators often use complementary resources from field borders and arable fields (Smith et al. 2014). Therefore, they

do not only suffer from loss of high-quality habitat, but also from the general loss of flower resources at a landscape scale (Persson & Smith 2013; Mallinger et al. 2016).

Pollinators respond differently to variation in landscape homogeneity (Jauker et al. 2009; Ekroos et al. 2010; Ekroos et al. 2013). For example, population declines and scarcity in homogeneous landscapes are biased towards species with high habitat specificity, narrow diet requirements, or low mobility (Biesmeijer et al. 2006; Ekroos et al. 2010; Bommarco et al. 2012), and some generalist species can even benefit from certain aspects of modern agricultural landscapes (Westphal et al. 2003). Therefore, it is possible that the lack of certain pollinators in homogeneous landscapes can be mitigated by the presence of other pollinators, and thus have little impact on wild plant pollination and regeneration.

Parallel declines of bees and insect-pollinated plants may indicate loss of pollination in contemporary landscapes (Biesmeijer et al. 2006) and correlations between species loss in wild bees and insect-pollinated plants have been observed along gradients of land-use intensity (Clough et al. 2014) and agricultural intensification (Gabriel & Tschardtke 2007). This could simply reflect the fact that bees need flowers, or that the two groups respond negatively to an external factor, such as intensive agriculture (Biesmeijer et al. 2006). Yet, the fact that insect-pollinated plants show steeper declines over time and landscape-scale land-use gradients than wind- or self-pollinated plants indicates that loss of suitable pollinators plays a role (Biesmeijer et al. 2006; Clough et al. 2014).

It is important to consider that more homogenous landscapes not only support fewer pollinators, but that there are also fewer flowers to visit (Persson & Smith 2013). Therefore, it is possible that the net number of visits per flower or per plant is only weakly or not at all affected by increasing landscape homogenisation and that the loss of insect-pollinated plants is driven entirely by factors unrelated to pollination, such as reduced mowing and grazing of permanent grasslands (Tyler et al. 2018), or

management changes (Kleijn & Snoeijs 1997; Honová et al. 2007). For example, herbicides and fertilisers can benefit the relative abundance of grasses at the expense of forbs (Kleijn & Snoeijs 1997; Honová et al. 2007). This calls for an assessment of whether flower visitation and seed set of insect-pollinated wild plants decline with increasing landscape homogeneity.

In this study, we aimed to address this knowledge gap. Ideally, this requires time-series data from landscapes characterized by different trajectories of change over time, which unfortunately is not available. We therefore capitalized on studies using a time-for-space substitution approach, in which the proportion of arable land varied between different study landscapes. We compiled data from such studies from eight European countries, on 22 highly pollinator-dependent native plant taxa. Using these data, we assessed the relationships between (i) flower visitation and seed set, (ii) the proportion of arable land and flower visitation, and (iii) the proportion of arable land and seed set. We expected (i) seed set to increase with flower visitation, and both (ii) flower visitation and (iii) seed set to decline with increasing proportion of arable land. By selecting a meta-analytical tool to address this aim, we were able to assess the general directions of the relationships, but also if these varied more than expected among studies.

MATERIAL AND METHODS

Data

Following an initial screening of the published literature, we contacted European researchers investigating wild plant pollination. We did this in 2015. We requested published and unpublished data from studies where seed set of a native insect-pollinated plant species had been measured at a minimum of five sites, which were separated by at least 1 km. Data should be from Europe, and most of the sites should be situated in landscapes (1 km radius) where at least 50% of the land was covered by agriculture, including arable land, permanent crops, pastures, etc. In addition, while there was no requirement that the original aim of the studies was to assess the effect of landscape homogeneity, we required variation in landscape homogeneity among the sites within each study. We identified suitable data for 22 plant taxa (for details, see supplementary file 1A), from both potted plants in experimental field studies (15 studies, 10 taxa) and naturally occurring plant populations (22 studies, 15 taxa). The data were collected in eight European countries (supplementary file 1A) between the years 2000 and 2015. When the identified studies included data on more than one plant species, we treated the data for each species as a separate study. All studies from the same country were later grouped (see statistical methods).

Flower visitation of the focal plants (i.e. focal observations of the study plants) was recorded in 14 of the studies, but we excluded one because bees and other flower visitors were not distinguished in that study. In another study, only bees had visited the flowers during the observations. In nine studies, flower visitation was quantified as the number of observed visitors divided by the number of flowers at

the site, or sub-site for studies including replication within sites. In the four remaining studies, the number of observed flower visitors was recorded per group of plants, using the same number of plants at all sites within a study. To keep the data as homogeneous and comparable among studies as possible, we did not include other types of estimates on pollinator densities such as data deriving from e.g. transect walks and pan traps, which we considered less relevant from a pollination perspective. To account for differences in sampling effort (sampling time and the number of observed flowers or plants) among the studies, we z-transformed flower visitation within each study (see statistical methods).

We extracted land-cover information for landscape sectors with 1 km radius around each site, or sub-site for studies with replications within sites. This radius corresponds to a common foraging distance of wild bees (Walther-Hellwig & Frankl 2000; Greenleaf et al. 2007) and has been shown to be relevant when assessing landscape effects on flower visitation by bees (e.g. Steffan-Dewenter et al. 2001). We extracted data from the CORINE online database (<https://www.eea.europa.eu/data-and-maps/data/corine-biotopes>) with a minimum mapping unit of 25 ha for polygon data from 2006. We expected that the proportion of arable land had remained rather stable during the time span in this study (2000–2015), because for example in the UK, only around 1% of the total area was reclassified between 2006 and 2012, and the changes were biased towards forested land (Cole et al. 2018).

Statistical methods

To analyse the relationships between (i) flower visitation and seed set, (ii) proportion of arable land and flower visitation, and (iii) proportion of arable land and seed set, we used hierarchical meta-analyses (R package metafor; Viechtbauer 2010) in R v.4.1.0 (R Core Team 2021). This is a meta-analytical tool that allows the use of factors to group data that are not independent, such as studies from the same country.

We estimated seed set as the proportional seed set per fruit, or as the number of seeds per fruit when the total number of ovules had not been measured (supplementary file 1B). To account for the different scales at which these estimates were reported (supplementary file 1B), we z-transformed (mean = 0, SD = 1) seed set within each study. For the same reason (supplementary file 1B), we z-transformed flower visitation by bees and other insects (after log-transforming the latter two), respectively, within each study. For studies where pollinator data included zeros, we added the minimum non-zero value (1 or smaller) for that particular study, before the log-transformation.

To prepare for the meta-analyses, we calculated effect sizes and variances for each study using either linear models when data were collected from independent sites, or linear mixed-effects models when we needed to specify random factors to account for geographical dependence among sub-sites from the same study site. For each of the studies (supplementary file 1A) with enough data we tested the following relationships:

- Flower visitation by bees and seed set (13 studies)
- Flower visitation by other insects and seed set (12 studies)

Table 1 – Results from the meta-analyses. Interrelations between the proportion of arable land, abundance of flower-visiting bees, as well as other flower-visiting insects, and seed set. For each meta-analysis, the number of studies (Studies (n)), the overall estimates (Pearson’s r), their associated 95% confidence intervals (CI₉₅) and p values are shown, in addition to Q statistics and the associated significance (Q_p) that indicates if effect sizes are heterogeneous; column I² (%) shows how much of the variability is caused by among study variation, rather than by sampling error. Potted: potted plants; nat. occ.: naturally occurring plants.

Variable	Studies (n)	Q	Q _p	I ² (%)	Pearson’s r	CI ₉₅	p
Flower visitation by bees							
Proportion of arable land	13	26.65	< 0.01	30.00	0.02	-1.12–1.08	0.97
Flower visitation by other insects							
Proportion of arable land	12	9.59	0.57	25.41	-0.66	-1.78–0.45	0.24
Seed set							
Flower visitation by bees	13	9.34	0.67	< 0.01	0.19	0.32–0.34	0.02
Flower visitation by other insects	12	17.05	0.11	23.14	-0.05	-0.26–0.16	0.64
Proportion of arable land (potted)	15	15.20	0.36	< 0.01	-0.002	-0.73–0.73	> 0.99
Proportion of arable land (nat. occ.)	22	48.32	< 0.01	10.20	-0.44	-1.25–0.36	0.28

- Proportion of arable land and flower visitation by bees (13 studies)
- Proportion of arable land and flower visitation by other insects (12 studies, because in one study, only bees visited the flowers)
- Proportion of arable land and seed set:
 - in potted plants (15 studies)
 - in naturally occurring plants (22 studies)

For each of these relationships, we then ran a separate random-effects meta-analysis, using the effect sizes and variances obtained in the previous step. In these meta-analyses, we specified country as a grouping factor to account for non-independence among studies from the same country, since we expected land-use history, climate, and political decisions to have shaped the landscape similarly within, but not necessarily among, countries. Since the shared phylogenetic history of plant species can pose a potential bias in meta-analyses (Vamosi et al. 2006; Chamberlain et al. 2012), we ran extra models with plant family as grouping factor, which had no qualitative impact on the results (supplementary file 2A). Because the plant species were all insect-pollinated, and therefore did not represent a random selection of angiosperms, we did not find it necessary to further control for the phylogeny of the plants. When extracting effect sizes and variances, some of the sub-replicated studies, for which we used linear mixed models, had a random variance close to zero. We therefore aggregated data from these studies and instead used mean values per site (aggregation of plant, insect, and landscape data from all sub-sites within a site). This had no qualitative impact on the results (supplementary file 2B). Because it is possible that potted plants and plants from naturally occurring populations respond differently to pollination-related effects of variation in the proportion of arable land, we ran the meta-analysis on the proportion of arable land and seed set separately for these two groups. For flower visitation by bees and other insects, we were not able to make this distinction due to the lower number of studies (n = 13 and n = 12, respectively).

We used Wald’s test, with restricted maximum likelihood, to evaluate if the effect differed from zero. We used Q statistics to evaluate if the heterogeneity among effect sizes was larger than expected by chance (Higgins & Thompson 2002), which would indicate that an important moderator is missing from the model (Del Re 2015). In addition, we used the inconsistency index I² to evaluate the proportion of total variation resulting from heterogeneity among studies rather than sampling error (Higgins & Thompson 2002). To ease interpretation of fig. 1, we used Fisher’s z-transformation and transformed the z values that had been used in the meta-analyses, to Pearson’s r correlations (Batáry et al. 2018).

RESULTS

Bees constituted 57 ± 29% (mean ± SD) of the total number of flower visitors across the studies where pollinator data had been collected and bees had been distinguished from other insects. The remaining 43% (referred to as other insects) were butterflies, moths, hoverflies, bee flies, and other unidentified flower-visiting insects. Only eight of the studies distinguished between honey bees and other bees. In these studies, wild bees were more common than honey bees, with honey bees constituting 11 ± 11% (mean ± SD) of the individual flower-visiting bees.

Seed set increased with increasing flower visitation by bees, but not with other insects (table 1, fig. 1). We found no overall relationship between the proportion of arable land and flower visitation by bees or by other insects (p > 0.05, table 1, fig. 2). The effect of the proportion of arable land on flower visitation by bees was significantly context-dependent, as shown by heterogeneous effect sizes among studies (Q_p < 0.05, table 1, fig. 2). The inconsistency index I² showed that true heterogeneity among studies explained 30% of the variation (table 1). Flower visitation by other insects was unrelated to the proportion of arable land in a consistent manner across studies (Q_p > 0.05, p > 0.05, table 1, fig. 2).

The proportion of arable land did not have a significant relationship with seed set for either naturally occurring or potted plants (p > 0.05, table 1, fig. 2). However, seed set of

naturally occurring plants was significantly heterogeneously related to the proportion of arable land among studies ($Q_p < 0.05$, table 1, fig. 2) and the inconsistency index I^2 shows that true heterogeneity explained 10.2% of the variation. Notably, we observed considerable variation in the direction of the effect across studies, even within species or genera (fig. 2). This was not the case concerning potted plants ($Q_p > 0.05$, $p > 0.05$, table 1, fig. 2). For seed set of naturally occurring plants, we tested if the heterogeneous effect sizes related to mean latitude (per study) and mean proportion of arable land (per study), but the addition of these moderators did not explain the heterogeneity (supplementary file 2C). We did not test if the addition of the above moderators explained the heterogeneity in the relationship between the proportion of arable land and flower visitation by bees, because of the low number of studies ($n = 13$).

DISCUSSION

As expected, seed set in the study plants increased with flower visitation by bees, but not by other insects, possibly because of a biased choice of bee pollinated plants among researchers. When splitting the different groups of bees, more than 90% of the flower-visiting bees were wild bees (similar to Hung et al. 2018), likely reflecting that honey bees tend to aggregate on mass-flowering resources (Rasmussen et al. 2021). This highlights the importance of maintaining these insects for the pollination of wild plants in agricultural landscapes. Despite that the abundance of wild bees often declines with the proportion of arable land in the surrounding landscape (Da Encarnação Coutinho et al. 2018), neither seed set, nor flower visitation declined with the proportion of arable land across all studies. For naturally occurring

plants as well as flower-visiting bees, the relationship with increasing proportions of arable land instead varied strongly among the studies, as shown by a significantly heterogeneous response (table 1), including strongly positive as well as strongly negative effects of increasing proportion of arable land in individual studies (fig. 2). Similar patterns for bees and seed set suggest that the effect on seed set is driven by flower-visitation and pollen transfer.

For potted plants, we found no relationship at all between increasing proportions of arable land and seed set, neither consistent ($p > 0.05$), nor inconsistent ($Q_p > 0.05$). Based on our results it is, however, unclear whether the observed difference between potted and naturally occurring plants is explained by a true difference between them or reflects a difference in sample size between the groups ($n = 15$ for potted plants, compared to $n = 22$ for naturally occurring plants).

Habitat fragmentation can disrupt flower visitation and pollination (Steffan-Dewenter & Tschamtkke 1999; Aguilar et al. 2006; Cranmer et al. 2011; Aguilar et al. 2019), but we show that an increasing proportion of arable land instead has a strongly variable impact on flower visitation by bees and seed set of wild plants, at least for the subset of plants that is covered in this study. While a complete lack of effect could have been explained by an inappropriate landscape gradient (Winfree et al. 2009) or scale (Westphal et al. 2006), or by the fact that both wild bees and the flowers they visit are rare in homogeneous landscapes (i.e. similar per capita flower visitation as in heterogeneous landscapes, cf. Persson & Smith 2013), a strongly variable effect needs a further explanation. We suggest that it results from variation in the extent to which the proportion of arable land reflects pollinator-relevant effects in the different study landscapes.




Figure 1 – Seed set in the tested plant species increased with flower visitation by bees (pink square), but not with other insects (blue triangle). Estimated mean values (squares and triangles) and 95% confidence intervals (error bars) from the meta-analyses are shown.

Correlation between seed set and proportion of arable land




Figure 2 – Forest plot showing the relationship with proportion of arable land for flower visitation by bees (open pink squares), flower visitation by other insects (open blue triangles), and seed set (grey circles) in naturally occurring (open circles) and potted (filled circles) plants. The estimated correlation coefficients (squares, circles, and triangles) and 95% confidence intervals (error bars) for each of the included studies are shown. The summary effects (mean \pm 95% CI) of increasing proportion of arable land are shown as diamonds.

Land-use change can affect pollinators in very different ways depending on how it alters floral resource availability (Winfree et al. 2011). While an increasing proportion of arable land often has a negative effect on wild bees when it occurs at the expense of flower-rich semi-natural habitat (Kennedy et al. 2013; Clough et al. 2014), it can instead have a positive effect when it substitutes flower-poor habitats, such as forests (Winfree et al. 2011). In addition, resource availability is not always low in landscapes dominated by arable land but depends on crop management (Tuck et al. 2014) and the configuration of arable land (Martin et al. 2019). When fields are small, or subject to organic farming, landscapes with high proportions of arable land can contain sufficient floral resources to maintain high bee densities (Rundlöf et al. 2008; Carrié et al. 2018; Hass et al. 2018; Martin et al. 2019). Therefore, the way in which increasing proportions of arable land affect seed set and flower visitation by bees depends on the type of arable land as well as the land-cover type it replaces.

Furthermore, while flower densities are usually higher in heterogeneous landscapes than in homogeneous landscapes, the characteristics of co-flowering plants themselves can affect each other in contrasting ways (Carvalho et al. 2014). They can either facilitate pollination by simultaneously attracting pollinators to a spot or compete for flower visitors (Hegland et al. 2009) and suffer from intraspecific pollen transfer (Morales & Traveset 2008). Hence, the effect of increasing proportion of arable land on flower visitation and seed set may depend on flower availability in the local habitat (cf. Herbertsson et al. 2018) and differ between plant species depending on their relative attractiveness to bees (Mesgaran et al. 2017). The focus on individual plant species in this study, compared to averaging over a whole plant community as in Clough et al. (2014), may thus have resulted in a larger heterogeneity of landscape effects.

Common flowering crops, such as oilseed rape and red clover, can increase bee densities (Westphal et al. 2003; Rundlöf et al. 2014), alter the bumble bee community composition (Diekötter et al. 2010), and redistribute bees across the landscape (Kovács-Hostyánszki et al. 2013; Holzschuh et al. 2016; Marja et al. 2018). Consequently, their effect on flower visitation depends on the location and phenology of the wild plants relative to the crop and on the extent to which they share pollinators with the crop (Cussans et al. 2010; Kovács-Hostyánszki et al. 2013; Herbertsson et al. 2017; Magrach et al. 2018). Flowering crops are often supplemented with honey bees, so that the abundance of honey bees on natural vegetation in agricultural landscapes increases with cultivation of bee-pollinated crops (Holzschuh et al. 2016; Magrach et al. 2017), affecting pollination networks (Magrach et al. 2017) and flower visitation of wild plants. In this study, we had no information on the extent to which flowering crops and bee hives occurred in the surrounding landscapes, and we could therefore not test if any of these variables caused the heterogeneity.

CONCLUSION

To conclude, we show that the relation between increasing proportion of arable land and pollination of wild plants is

highly context-dependent. We found no consistent evidence that an increasing proportion of arable land reduces per capita flower visitation and seed set, which instead varied across the studies. While this suggests that other factors, such as habitat degradation, are stronger drivers of the loss of insect-pollinated plants in landscapes dominated by arable land, we also show that lack of flower-visiting bees can harm seed set in such plants, which could affect their long-term persistence. In order to develop adequate conservation solutions for wild insect-pollinated plants, it is therefore important to identify why the increasing proportion of arable land has a variable, and sometimes contrasting, effect on flower visitation and seed set.

DATA AVAILABILITY STATEMENT

All data used in the analyses are presented in supplementary file 3.

SUPPLEMENTARY FILES

Supplementary file 1 – Contains information about the data sets used in the meta-analyses.

<https://doi.org/10.5091/plecevo.2021.1884.2525>

Supplementary file 2 – Contains extended results.

<https://doi.org/10.5091/plecevo.2021.1884.2527>

Supplementary file 3 – Contains all data used in the meta-analyses.

<https://doi.org/10.5091/plecevo.2021.1884.2529>

ACKNOWLEDGEMENTS

We thank Carole Chateil, Volker Gaebele, Emmanuelle Porcher, and Marie Winsa for data contribution; Yann Clough and Peter Olsson for help on analyses and extraction of landscape data. The original idea was conceived as part of the project ‘STEP - Status and trends of European pollinators’ funded by the European Union in the 7th Framework Programme (grant 244090), which funded RB, ISD, HGS, SP, MR, LH, JME, and KS. JE and PC were funded by MULTAGRI/FORMAS. MA was funded by the EU FP5 project ‘Evaluating current European agri-environment schemes to quantify and improve nature conservation efforts in agricultural landscapes’ (EASY; QLRT-2001-01495) and the Swiss Federal Office for Science and Technology (01-0524-2). AJ and EÖ were funded by FORMAS. TTe and VS were supported by institutional research funding IUT (IUT20-33) of the Estonian Ministry of Education and Research, and TTe also received funding (grant no. 42900/1312/3166) from the Internal Grant Agency of the Faculty of Environmental Sciences, Czech University of Life Sciences Prague. IB was funded by MC-CIG BeeFun project: PCIG14-GA-2013-631653 and thanks the Doñana NP staff members for granting access to the Park and Curro Molina for conducting the field work. JS, EFP, and SM were funded by DAFM and IRC. LH was funded by a grant from Formas to HGS and by a mobility grant from Formas (2018-01466). ALH was funded by the FarmLand project supported by the German Ministry of Research and Education (FKZ: 01LC1104A). AKH was supported by the NKFIH project

(FK123813) and was a Bolyai Fellow of MTA. PB was supported by the NKFIH (KKP 133839).

AUTHOR CONTRIBUTIONS

HGS, MR, and RB formulated the original idea; LH and JE developed the idea with contribution from HGS; LH compiled and analyzed the data with input from HGS, JE, PC, and PB. AH, ALH, AKH, AJ, AS, BJ, EÖ, HGS, IB, ISD, JE, JCS, JME, JPGV, LH, MA, MHE, MR, NF, PB, RB, SF, SH, SL, SP, TD, TTe, TTj, VS, VG, WK, and WEK provided data. LH interpreted the data and wrote the manuscript with contributions from all authors.

REFERENCES

- Aguilar R., Ashworth L., Galetto L. & Aizen M.A. 2006. Plant reproductive susceptibility to habitat fragmentation: review and synthesis through a meta-analysis. *Ecology Letters* 9: 968–980. <https://doi.org/10.1111/j.1461-0248.2006.00927.x>
- Aguilar R., Cristóbal-Pérez E.J., Balvino-Olvera F.J., et al. 2019. Habitat fragmentation reduces plant progeny quality: a global synthesis. *Ecology Letters* 22: 1163–1173. <https://doi.org/10.1111/ele.13272>
- Batáry P., Kurucz K., Suarez-Rubio M. & Chamberlain D.E. 2018. Non-linearities in bird responses across urbanization gradients: a meta-analysis. *Global Change Biology* 24: 1046–1054. <https://doi.org/10.1111/gcb.13964>
- Biesmeijer J.C., Roberts S.P.M., Reemer M., et al. 2006. Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science* 313: 351–354. <https://doi.org/10.1126/science.1127863>
- Bilz M., Kell S.P., Maxted N. & Lansdown R.V. 2011. European Red List of Vascular Plants. Publications Office of the European Union, Luxembourg. <https://doi.org/10.2779/8515>
- Bobbink R., Hicks K., Galloway J., et al. 2010. Global assessment of nitrogen deposition effects on terrestrial plant diversity: a synthesis. *Ecological Applications* 20: 30–59. <https://doi.org/10.1890/08-1140.1>
- Bommarco R., Lundin O., Smith H.G. & Rundlöf M. 2012. Drastic historic shifts in bumble-bee community composition in Sweden. *Proceedings of the Royal Society B* 279: 309–315. <https://doi.org/10.1098/rspb.2011.0647>
- Burd M. 1994. Bateman's principle and plant reproduction: the role of pollen limitation in fruit and seed set. *The Botanical Review* 60: 83–139. <https://doi.org/10.1007/BF02856594>
- Carrié R., Ekroos J. & Smith H.G. 2018. Organic farming supports spatiotemporal stability in species richness of bumblebees and butterflies. *Biological Conservation* 227: 48–55. <https://doi.org/10.1016/j.biocon.2018.08.022>
- Carvalho L.G., Biesmeijer J.C., Benadi G., et al. 2014. The potential for indirect effects between co-flowering plants via shared pollinators depends on resource abundance, accessibility, and relatedness. *Ecology Letters* 17: 1389–1399. <https://doi.org/10.1111/ele.12342>
- Chamberlain S.A., Hovick S.M. & Dibble C.J. 2012. Does phylogeny matter? Assessing the impact of phylogenetic information in ecological meta-analysis. *Ecology Letters* 15: 627–636. <https://doi.org/10.1111/j.1461-0248.2012.01776.x>
- Clough Y., Ekroos J., Báldi A., et al. 2014. Density of insect-pollinated grassland plants decreases with increasing surrounding land-use intensity. *Ecology Letters* 17: 1168–1177. <https://doi.org/10.1111/ele.12325>
- Cole B., Smith G. & Balzter H. 2018. Acceleration and fragmentation of CORINE land cover changes in the United Kingdom from 2006–2012 detected by Copernicus IMAGE2012 satellite data. *International Journal of Applied Earth Observation and Geoinformation* 73: 107–122. <https://doi.org/10.1016/j.jag.2018.06.003>
- Cranmer L., McCollin D. & Ollerton J. 2011. Landscape structure influences pollinator movements and directly affects plant reproductive success. *Oikos* 121: 562–568. <https://doi.org/10.1111/j.1600-0706.2011.19704.x>
- Cussans J., Goulson D., Sanderson R., Goffe L., Darvill B. & Osborne J.L. 2010. Two bee-pollinated plant species show higher seed production when grown in gardens compared to arable farmland. *PLoS ONE* 5: e11753. <https://doi.org/10.1371/journal.pone.0011753>
- Da Encarnação Coutinho J.G., Garibaldi L.A. & Viana B.F. 2018. The influence of local and landscape scale on single response traits in bees: a meta-analysis. *Agriculture, Ecosystems and Environment* 256: 61–73. <https://doi.org/10.1016/j.agee.2017.12.025>
- Del Re A.C. 2015. A practical tutorial on conducting meta-analysis in R. *The Quantitative Methods for Psychology* 11: 37–50. <https://doi.org/10.20982/tqmp.11.1.p037>
- Diekötter T., Kadoya T., Peter F., Wolters V. & Jauker F. 2010. Oilseed rape crops distort plant–pollinator interactions. *Journal of Applied Ecology* 47: 209–214. <https://doi.org/10.1111/j.1365-2664.2009.01759.x>
- Ekroos J., Heliölä J. & Kuussaari M. 2010. Homogenization of lepidopteran communities in intensively cultivated agricultural landscapes. *Journal of Applied Ecology* 47: 459–467. <https://doi.org/10.1111/j.1365-2664.2009.01767.x>
- Ekroos J., Rundlöf M. & Smith H.G. 2013. Trait-dependent responses of flower-visiting insects to distance to semi-natural grasslands and landscape heterogeneity. *Landscape Ecology* 28: 1283–1292. <https://doi.org/10.1007/s10980-013-9864-2>
- Fontaine C., Dajoz I., Meriguet J. & Loreau M. 2005. Functional diversity of plant–pollinator interaction webs enhances the persistence of plant communities. *PLoS Biology* 4: e1. <https://doi.org/10.1371/journal.pbio.0040001>
- Gabriel D. & Tschardt T. 2007. Insect pollinated plants benefit from organic farming. *Agriculture, Ecosystems & Environment* 118: 43–48. <https://doi.org/10.1016/j.agee.2006.04.005>
- Greenleaf S.S., Williams N.M., Winfree R. & Kremen C. 2007. Bee foraging ranges and their relationship to body size. *Oecologia* 153: 589–596. <https://doi.org/10.1007/s00442-007-0752-9>
- Hass A.L., Kormann U.G., Tschardt T., et al. 2018. Landscape configurational heterogeneity by small-scale agriculture, not crop diversity, maintains pollinators and plant reproduction in western Europe. *Proceedings of the Royal Society B* 285: 20172242. <https://doi.org/10.1098/rspb.2017.2242>
- Hallmann A., Ssymank A., Sorg M., de Kroon H. & Jongejans E. 2021. Insect biomass decline scaled to species diversity: general patterns derived from a hoverfly community. *Proceedings of the National Academy of Sciences of the United States of America* 118: e2002554117. <https://doi.org/10.1073/pnas.2002554117>
- Hegland S.J., Grytnes J. & Totland Ø. 2009. The relative importance of positive and negative interactions for pollinator attraction in a plant community. *Ecological Research* 24: 929–936. <https://doi.org/10.1007/s11284-008-0572-3>
- Herbertsson L., Rundlöf M. & Smith H.G. 2017. The relation between oilseed rape and pollination of later flowering plants

- varies across plant species and landscape contexts. *Basic and Applied Ecology* 24: 77–85. <https://doi.org/10.1016/j.baae.2017.08.001>
- Herbertsson L., Jönsson A.M., Andersson G.K.S., et al. 2018. The impact of sown flower strips on plant reproductive success in Southern Sweden varies with landscape context. *Agriculture, Ecosystems & Environment* 259: 127–134. <https://doi.org/10.1016/j.agee.2018.03.006>
- Higgins J.P.T. & Thompson S.G. 2002. Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine* 21: 1539–1558. <https://doi.org/10.1002/sim.1186>
- Holzschuh A., Dainese M., González-Varo J.P., et al. 2016. Mass-flowering crops dilute pollinator abundance in agricultural landscapes across Europe. *Ecology Letters* 19: 1228–1236. <https://doi.org/10.1111/ele.12657>
- Honová D., Hecman M., Klaudivsová M., Pavlů V., Kocourková D. & Hakl J. 2007. Species composition of an alluvial meadow after 40 years of applying nitrogen, phosphorus and potassium fertilizer. *Preslia* 79: 245–258.
- Hung K.L.J., Kingston J.L., Albrecht M., Holway D.A & Kohn J.R. 2018. The worldwide importance of honey bees as pollinators in natural habitats. *Proceedings of the Royal Society B* 285: 20172140. <https://doi.org/10.1098/rspb.2017.2140>
- Jauker F., Diekötter T., Schwartzbach F. & Wolters V. 2009. Pollinator dispersal in an agricultural matrix: opposing responses of wild bees and hoverflies to landscape structure and distance from main habitat. *Landscape Ecology* 24: 547–555. <https://doi.org/10.1007/s10980-009-9331-2>
- Kennedy C.M., Lonsdorf E., Neel M.C., et al. 2013. A global quantitative synthesis of local and landscape effects on wild bee pollinators in agroecosystems. *Ecology Letters* 16: 584–599. <https://doi.org/10.1111/ele.12082>
- Kleijn D. & Snoeijs G.I.J. 1997. Field boundary vegetation and the effects of agrochemical drift: botanical change caused by low levels of herbicide and fertilizer. *Journal of Applied Ecology* 34: 1413–1425. <https://doi.org/10.2307/2405258>
- Kovács-Hostyánszki A., Haenke S., Batáry P., et al. 2013. Contrasting effects of mass-flowering crops on bee pollination of hedge plants at different spatial and temporal scales. *Ecological Applications* 23: 1938–1946. <https://doi.org/10.1890/12-2012.1>
- Kuussaari M., Heliölä J., Pöyry J. & Saarinen K. 2007. Contrasting trends of butterfly species preferring semi-natural grasslands, field margins and forest edges in northern Europe. *Journal of Insect Conservation* 11: 351–366. <https://doi.org/10.1007/s10841-006-9052-7>
- Magrach A., González-Varo J.P., Boiffier M., Vilà M. & Bartomeus I. 2017. Honeybee spillover reshuffles pollinator diets and affects plant reproductive success. *Nature Ecology & Evolution* 1: 1299–1307. <https://doi.org/10.1038/s41559-017-0249-9>
- Magrach A., Holzschuh A., Bartomeus I., et al. 2018. Plant-pollinator networks in semi-natural grasslands are resistant to the loss of pollinators during blooming of mass-flowering crops. *Ecography* 41: 62–74. <https://doi.org/10.1111/ecog.02847>
- Mallinger R.E., Gibbs J. & Gratton C. 2016. Diverse landscapes have a higher abundance and species richness of spring wild bees by providing complementary floral resources over bees' foraging periods. *Landscape Ecology* 31: 1523–1535. <https://doi.org/10.1007/s10980-015-0332-z>
- Marja R., Viik E., Mänd M., Phillips J., Klein A-M. & Batáry P. 2018. Crop rotation and agri-environment schemes determine bumblebee communities via flower resources. *Journal of Applied Ecology* 55: 1714–1724. <https://doi.org/10.1111/1365-2664.13119>
- Martin E.A., Dainese M., Clough Y., et al. 2019. The interplay of landscape composition and configuration: new pathways to manage functional biodiversity and agroecosystem services across Europe. *Ecology Letters* 22: 1083–1094. <https://doi.org/10.1111/ele.13265>
- Mesgaran M.B., Bouhours J., Lewis M.A. & Cousens R.D. 2017. How to be a good neighbour: facilitation and competition between two co-flowering species. *Journal of Theoretical Biology* 422: 72–83. <https://doi.org/10.1016/j.jtbi.2017.04.011>
- Morales C.L. & Traveset A. 2008. Interspecific pollen transfer: magnitude, prevalence and consequences for plant fitness. *Critical Reviews in Plant Sciences* 27: 221–238. <https://doi.org/10.1080/07352680802205631>
- Ollerton J., Winfree R. & Tarrant S. 2011. How many flowering plants are pollinated by animals? *Oikos* 120: 321–326. <https://doi.org/10.1111/j.1600-0706.2010.18644.x>
- Papanikolaou A.D., Kühn I., Frenzel M., et al. 2017. Wild bee and floral diversity co-vary in response to the direct and indirect impacts of land use. *Ecosphere* 8: e02008. <https://doi.org/10.1002/ecs2.2008>
- Persson A.S. & Smith H.G. 2013. Seasonal persistence of bumblebee populations is affected by landscape context. *Agriculture, Ecosystems & Environment* 165: 201–209. <https://doi.org/10.1016/j.agee.2012.12.008>
- Persson A.S., Rundlöf M., Clough Y. & Smith H.G. 2015. Bumble bees show trait-dependent vulnerability to landscape simplification. *Biodiversity and Conservation* 24: 3469–3489. <https://doi.org/10.1007/s10531-015-1008-3>
- Power E.F., Jackson Z. & Stout J.C. 2016. Organic farming and landscape factors affect abundance and richness of hoverflies (Diptera, Syrphidae) in grasslands. *Insect Conservation and Diversity* 9: 244–253. <https://doi.org/10.1111/icad.12163>
- Qi M., Sun T., Xue S., Yang W., Shao D. & Martínez-López J. 2018. Competitive ability, stress tolerance and plant interactions along stress gradients. *Ecology* 99: 848–857. <https://doi.org/10.1002/ecy.2147>
- Rasmussen C., Dupont Y.L., Bang Madsen H., et al. 2021. Evaluating competition for forage plants between honey bees and wild bees in Denmark. *PLoS ONE* 16: e0250056. <https://doi.org/10.1371/journal.pone.0250056>
- R Core Team 2021. R: a language and environment for statistical computing. Version 4.1.0. R Foundation for Statistical Computing, Vienna. Available from <https://www.r-project.org/> [accessed 8 Oct. 2021].
- Robinson R.A. & Sutherland W.J. 2002. Post-war changes in arable farming and biodiversity in Great Britain. *Journal of Applied Ecology* 39: 157–176. <https://doi.org/10.1046/j.1365-2664.2002.00695.x>
- Rundlöf M., Nilsson H. & Smith H.G. 2008. Interacting effects of farming practice and landscape context on bumblebees. *Biological Conservation* 141: 417–426. <https://doi.org/10.1016/j.biocon.2007.10.011>
- Rundlöf M., Persson A.S., Smith H.G. & Bommarco R. 2014. Late-season mass-flowering red clover increases bumble bee queen and male densities. *Biological Conservation* 172: 138–145. <https://doi.org/10.1016/j.biocon.2014.02.027>
- Scheper J., Reemer M., van Kats R., et al. 2014. Museum specimens reveal loss of pollen host plants as key factor driving wild bee decline in The Netherlands. *Proceedings of the National*

- Academy of Sciences of the United States of America* 111: 17552–17557. <https://doi.org/10.1073/pnas.1412973111>
- Smith H.G., Birkhofer K., Clough Y., Ekroos J., Olsson O. & Rundlöf M. 2014. Beyond dispersal: the role of animal movement in modern agricultural landscapes. In: Hansson L.-A. & Åkesson S. (eds) *Animal movement across scales*. Oxford University Press, Oxford. <https://doi.org/10.1093/acprof:oso/9780199677184.003.0004>
- Steffan-Dewenter I., Münzenberg U. & Tschardt T. 2001. Pollination, seed set and seed predation on a landscape scale. *Proceedings of the Royal Society B* 268: 1685–1690. <https://doi.org/10.1098/rspb.2001.1737>
- Steffan-Dewenter I. & Tschardt T. 1999. Effects of habitat isolation on pollinator communities and seed set. *Oecologia* 121: 432–440. <https://doi.org/10.1007/s004420050949>
- Theodose T.A. & Bowman W.D. 1997. The influence of interspecific competition on the distribution of an alpine graminoid: evidence for the importance of plant competition in an extreme environment. *Oikos* 79: 101–114. <https://doi.org/10.2307/3546095>
- Tuck S.L., Winqvist C., Mota F., Ahnström J., Turnbull L.A. & Bengtsson J. 2014. Land-use intensity and the effects of organic farming on biodiversity: a hierarchical meta-analysis. *Journal of Applied Ecology* 51: 746–755. <https://doi.org/10.1111/1365-2664.12219>
- Turnbull L.A., Crawley M.J. & Rees M. 2000. Are plant populations seed-limited? A review of seed sowing experiments. *Oikos* 88: 225–238. <https://doi.org/10.1034/j.1600-0706.2000.880201.x>
- Tyler T., Herbertsson L., Olsson P.A., et al. 2018. Climate warming and land-use changes drive broad-scale floristic changes in Southern Sweden. *Global Change Biology* 24: 2607–2621. <https://doi.org/10.1111/gcb.14031>
- Vamosi J.C., Knight T.M., Steets J.A., Mazer S.J., Burd M. & Ashman T.-L. 2006. Pollination decays in biodiversity hotspots. *Proceedings of the National Academy of Sciences of the United States of America* 103: 956–961. <https://doi.org/10.1073/pnas.0507165103>
- van Swaay C., Warren M. & Loïs G. 2006. Biotope use and trends of European butterflies. *Journal of Insect Conservation* 10: 189–209. <https://doi.org/10.1007/s10841-006-6293-4>
- Viechtbauer W. 2010. Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software* 36: 1–48. <https://doi.org/10.18637/jss.v036.i03>
- WallisdeVries M., Swaay C. & Plate C. 2012. Changes in nectar supply: a possible cause of widespread butterfly decline. *Current Zoology* 58: 384–391. <https://doi.org/10.1093/czoolo/58.3.384>
- Walther-Hellwig K. & Frankl R. 2000. Foraging habitats and foraging distances of bumblebees, *Bombus* spp. (Hym., Apidae), in an agricultural landscape. *Journal of Applied Entomology* 124: 299–306. <https://doi.org/10.1046/j.1439-0418.2000.00484.x>
- Westphal C., Steffan-Dewenter I. & Tschardt T. 2003. Mass flowering crops enhance pollinator densities at a landscape scale. *Ecology Letters* 6: 961–965. <https://doi.org/10.1046/j.1461-0248.2003.00523.x>
- Westphal C., Steffan-Dewenter I. & Tschardt T. 2006. Bumblebees experience landscapes at different spatial scales: possible implications for coexistence. *Oecologia* 149: 289–300. <https://doi.org/10.1007/s00442-006-0448-6>
- Winfrey R., Aguilar R., Vázquez D.P., LeBuhn G. & Aizen M.A. 2009. A meta-analysis of bees' responses to anthropogenic disturbance. *Ecology* 90: 2068–2076. <https://doi.org/10.1890/08-1245.1>
- Winfrey R., Bartomeus I. & Cariveau D.P. 2011. Native pollinators in anthropogenic habitats. *Annual Review of Ecology, Evolution, and Systematics* 42: 1–22. <https://doi.org/10.1146/annurev-ecolsys-102710-145042>

Communicating editor: Renate Wesselingh.

Submission date: 14 Apr. 2021

Acceptance date: 13 Sep. 2021

Publication date: 23 Nov. 2021