



Recently fixed carbon fuels microbial activity several meters below the soil surface

Andrea Scheibe¹, Carlos A. Sierra^{2,3}, and Marie Spohn^{1,4}

¹Bayreuth Center of Ecology and Environmental Research (BayCEER), University of Bayreuth, Bayreuth, Germany

²Department of Biogeochemical Processes, Max Planck Institute for Biogeochemistry, Jena, Germany

³Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden

⁴Department of Soil and Environment, Swedish University of Agricultural Sciences, Uppsala, Sweden

Correspondence: Marie Spohn (marie.spohn@slu.se)

Received: 23 September 2022 – Discussion started: 26 September 2022

Revised: 29 January 2023 – Accepted: 3 February 2023 – Published: 21 February 2023

Abstract. The deep soil, > 1 m, harbors a substantial share of the global microbial biomass. Currently, it is not known whether microbial activity several meters below the surface is fueled by recently fixed carbon or by old carbon that persisted in soil for several hundred years. Understanding the carbon source of microbial activity in deep soil is important to identify the drivers of biotic processes in the critical zone. Therefore, we explored carbon cycling in soils in three climate zones (arid, mediterranean, and humid) of the Coastal Cordillera of Chile down to a depth of 6 m, using carbon isotopes. Specifically, we determined the $^{13}\text{C} : ^{12}\text{C}$ ratio ($\delta^{13}\text{C}$) of soil and roots and the $^{14}\text{C} : ^{12}\text{C}$ ratio ($\Delta^{14}\text{C}$) of soil organic carbon and $\text{CO}_2\text{-C}$ respired by microorganisms. We found that the $\Delta^{14}\text{C}$ of the respired $\text{CO}_2\text{-C}$ was significantly higher than that of the soil organic carbon in all soils. Further, we found that the $\delta^{13}\text{C}$ of the soil organic carbon changed only in the upper decimeters (by less than 6‰). Our results show that microbial activity several meters below the soil surface is mostly fueled by recently fixed carbon that is on average much younger than the total soil organic carbon present in the respective soil depth increments, in all three climate zones. Further, our results indicate that most decomposition that leads to enrichment of ^{13}C occurs in the upper decimeters of the soils, which is possibly due to stabilization of organic carbon in the deep soil. In conclusion, our study demonstrates that microbial processes in the deep soil several meters below the surface are closely tied to input of recently fixed carbon.

1 Introduction

The deep soil (> 1 m) harbors not only a large part of the global soil microbial biomass (Pedersen, 1997; Krumholz, 2000; Akob and Küsel, 2011), but also large amounts of organic carbon (C) (Rumpel and Kögel-Knabner, 2011; Marin-Spiotta et al., 2014; Jackson et al., 2017; Moreland et al., 2021; Marín-Spiotta and Hobbey, 2022). In the uppermost meter of soils, microorganisms have been shown to rely primarily on young organic C, tying microbial activity belowground closely to photosynthetic activity aboveground (Trumbore, 2000; van Hees et al., 2005; Högberg and Read, 2006; Högberg et al., 2008). Whether this relationship holds for deep soils (> 1 m) has hardly been investigated yet, which limits the understanding of biotic processes, such as biogenic weathering, in the deep critical zone.

The age of C metabolized by microorganisms can be estimated by measuring the ^{14}C signature of the C emitted by the soil microbial biomass in the form of CO_2 in incubations (Trumbore, 2000). To our knowledge, and according to the International Soil Radiocarbon Database (Lawrence et al., 2020), the ^{14}C signature of CO_2 respired by microorganisms in soil below 1.0 m has been measured only in permafrost soils (Dutta et al., 2006; Lee et al., 2012; Gentsch et al., 2018), showing that microbial respiration during winter relies strongly on C that has persisted in these soils for centuries (Dutta et al., 2006; Lee et al., 2012; Gentsch et al., 2018; Pedron et al., 2022). For non-permafrost soils, there are no data on the $\Delta^{14}\text{C}$ of CO_2 respired by microorganisms for depth increments below 90–100 cm (Nagy et al., 2018; Lawrence et al., 2020). The $\Delta^{14}\text{C}$ of total soil CO_2 at greater




Figure 1. Location of the three study sites in the Coastal Cordillera of Chile together with a photo of the vegetation and soil at each of the sites. The map is taken from Werner et al. (2018) and was created based on Luebert and Plissock (2017).

depths has only been measured in the field (Trumbore et al., 1995; Fierer et al., 2005), but the total soil CO₂ can be largely composed of CO₂ respired by roots (Hanson et al., 2000) and therefore masks the information about soil C cycling by microbial processes.

To test the hypothesis that microbial activity in deep, non-permafrost soil is driven by young organic C, we explored C cycling in soils of the Coastal Cordillera of Chile using C isotopes. The bedrock in this region is extraordinarily deeply weathered (Vázquez et al., 2016; Hayes et al., 2020; Krone et al., 2021), possibly because of the CO₂ and organic acids produced by soil microorganisms (Berner, 1997; Berg and Banwart, 2000; Hoffland et al., 2004; Finlay et al., 2020; Uroz et al., 2022). We studied soils in the arid, mediterranean, and humid climate zone that correspond to the vegetation zones arid shrubland, sclerophyllous forest, and humid temperate forest, respectively, in the Coastal Cordillera of Chile (Fig. 1). For this purpose, three to four soil profiles per site were dug down to a depth of 6.0 m (humid and mediterranean site) or 2.0 m (arid site). Soil and root samples were collected at regular intervals. To determine the age of C metabolized by microorganisms, we determined the $\Delta^{14}\text{C}$ of the C that was emitted by soil microorganisms in the form of CO₂ in incubations. Further, we determined the $\Delta^{14}\text{C}$ of the soil total organic carbon (TOC) to estimate its age, and we also measured its concentration. To quantify the extent of microbial decomposition of organic matter in soil, we determined the $\delta^{13}\text{C}$ of TOC and roots. Finally, in order to estimate microbial biomass and activity, we extracted and quantified DNA from soil and determined the soil-microbial respiration rate.

2 Material and methods

2.1 Study sites

All three study sites are located along a precipitation gradient in the Coastal Cordillera of Chile between 30 and 38° S (Fig. 1; Table 1). Mean annual precipitation is about 87, 436, and 1084 mm at the arid, mediterranean, and humid site, respectively. The arid site is located in the private reserve Santa Gracia (Table 1). The vegetation is classified as arid shrubland. The vegetation is sparse (10% cover) and dominated by *Proustia cuneifolia*, *Cordia decandra*, and *Adesmia* spp. (Spohn and Holzheu, 2021). The mediterranean site is located ~4.2 km south of the national park La Campana, in a water protection area. The soils are covered by a low sclerophyllous forest. The humid site is located on the eastern border of the national park Nahuelbuta. The soils are covered by a mixed humid, temperate forest with coniferous and deciduous tree species. Further information about climate, geology, pedogenesis, and vegetation is provided in Bernhard et al. (2018), Oeser et al. (2018), and Brucker and Spohn (2019).

2.2 Soils and sampling

The soils at all three sites have formed from plutonic bedrock of similar granitoid lithology (Oeser et al., 2018). They were not affected by glaciation during the last glacial maximum (~19 000–23 000 years ago; Hulton et al., 2002) and have not received volcanic ashes (Oeser et al., 2018). The soils of the arid and mediterranean site are classified as Cambisols, and the soils at the humid site are classified as Cambisols and Umbrisols (IUSS Working Group WRB, 2015).

The soil sampling was conducted at the arid site in March 2019 and at the mediterranean and humid site in March 2020. For the soil sampling, three soil profiles down to a depth of 200 cm were excavated on each site in an area of 1 km². One of them was further extended to a depth of 600 cm at the mediterranean and humid site. Furthermore, at the humid site, an additionally fourth profile was established down to 400 cm soil depth. At the arid site, the soil profiles were chosen to be less deep because the soils at this site are less deeply developed than at the other two sites (Bernhard et al., 2018). The soil sampling was conducted the day after excavation from the bottom to the top of the profiles in the following soil increments: from 600 to 200 cm in 50 cm increments and from 200 to 20 cm in 20 cm increments. For the uppermost 20 cm, the following soil increments were sampled: 0–5, 5–10, and 10–20 cm. A subsample of the organic layer on top of each profile was collected at the mediterranean and humid site (there is no organic layer at the arid site). In addition, root samples were taken to complement the roots collected with the soil.

The percentage of fine soil (< 2 mm) is smaller at the arid site than at the two other sites throughout all soil depth incre-

Table 1. Properties of the three sites in the Coastal Cordillera in Chile, including mean annual temperature (MAT), mean annual precipitation (MAP), and the locations of the soil profiles.

| Climate zone | Site name | MAT [°C] | MAP [mm yr ⁻¹] | Biome | Profile | Longitude | Latitude |
|---------------|--------------|-------------|-------------------------------|-------------------------------------|---------|------------|------------|
| Arid | Santa Gracia | 16.1 | 87 | Arid shrubland | 1 | -71.159439 | -29.759769 |
| | | | | | 2 | -71.160226 | -29.759037 |
| | | | | | 3 | -71.161234 | -29.759465 |
| Mediterranean | La Campana | 14.9 | 436 | Mediterranean sclerophyllous forest | 1 | -71.043710 | -33.028375 |
| | | | | | 2 | -71.041269 | -33.028585 |
| | | | | | 3 | -71.047170 | -33.028718 |
| Humid | Nahuelbuta | 14.1 | 1084 | Humid, temperate forest | 1 | -72.95065 | -37.79371 |
| | | | | | 2 | -72.95125 | -37.79017 |
| | | | | | 3 | -72.94868 | -37.79533 |
| | | | | | 4 | -72.95206 | -37.79517 |

ments (Table S1 in Supplement). The percentage of fine soil (< 2 mm) strongly decreases with increasing soil depth at the arid site and less strongly at the humid site (Table S1). At the arid site, the soil texture of the fine soil fraction (< 2 mm), averaged over all profiles, is loamy sand in the uppermost 80 cm, silty sand from 80 to 100 cm soil depth, and pure sand below 100 cm soil depth. At the mediterranean site, the soil texture in the uppermost 160 cm is medium to slightly loamy sand and below 160 cm soil depth pure sand. The soil texture at the humid site is loamy. All soils are free of carbonates.

2.3 Sample preparation and soil physical analyses

All field-moist soil samples were sieved (\emptyset 2 mm) and roots and other debris removed by hand. The weights of both fine soil fraction ($sf_{<2\text{mm}}$) and gravel or stone fraction ($sf_{>2\text{mm}}$) were determined for each soil increment. After subtracting the soil water content (SWC), the proportion of the fine soil fraction ($sf_{<2\text{mm}}$) per mass of the total sample was calculated. Subsamples of the sieved soil (< 2 mm) were either immediately dried at 60 °C (for chemical and isotope analyses), or stored at 5 °C (for incubation experiments), or frozen at -20 °C (for DNA analyses). The subsample of the organic layer was dried at 40 °C (for chemical and isotope analyses). For each soil increment, the SWC of the field-moist soil was determined gravimetrically. The soil water-holding capacity (WHC) was determined for eight soil increments down to 600 cm (0–5, 5–10, 10–20, 40–60, 100–120, 180–200, 350–400, and 550–600 cm), in order to facilitate the adjustment of comparable SWCs in the incubation experiment. The WHCs for the other soil increments were obtained by extrapolation between measured soil increments. For the determination of the WHC, field-moist soil samples were oversaturated with water for 24 h and then left on a sand bath for 24 h. Afterwards, the samples were weighed before and after drying at 105 °C for 24 h.

For the mediterranean and humid site, fine roots (\emptyset < 2 mm) per individual soil increment were collected during the sieving process. At the arid site, fine roots were not present in all soil increments, and therefore roots of different increments were combined to a composite sample to acquire enough biomass for the isotope analyses. All root samples were washed and subsequently dried at 60 °C.

2.4 Chemical and stable isotope analyses

Subsamples of the organic layer, dried fine soil (< 2 mm) and roots were milled and analyzed for their carbon (C) concentration as well as stable carbon isotope ratio ($\delta^{13}\text{C}$) using an element analyzer (NA 1108, CE Instruments, Milan, Italy) coupled to an isotope-ratio mass spectrometer (delta S, Finnigan MAT, Bremen, Germany) via a ConFlo III interface (Finnigan MAT, Bremen, Germany). Subsequently, the soil total organic C (TOC) concentrations were re-calculated per total soil mass ($sf_{<2\text{mm}} + sf_{>2\text{mm}}$).

2.5 Soil respiration rates

To determine the soil respiration rate for each soil depth increment, a subsample of the fine soil (< 2 mm) was incubated in a 350 mL incubation jar with septum. The amount of soil (dry-weight equivalent) used for the incubation differed between soil depth increments: 20 g (mediterranean and humid) or 30 g (arid) for 0–20 cm, 40 g for 20–100 cm, 60 g for 100–200 cm, 80 g for 200–400 cm, and 100 g for 400–600 cm. After adjusting the SWC to 60% WHC, the incubation jars were closed air-tight and the samples pre-incubated at 20 °C in the dark for 3 (arid) or 4 (mediterranean and humid) days. Subsequent to the pre-incubation period, the incubation jars were aerated for several minutes, and closed again, and the CO₂ concentration within the incubation jars was immediately analyzed (time point T0). Afterwards, the samples were incubated at 20 °C in the dark for 6 weeks. During the whole incubation period, the CO₂ concentration in the incubation

jars was measured weekly (and additionally for mediterranean and humid site at day 3). For this purpose, a gas sample of 50 or 100 μL was collected from the headspace of each incubation jar using a syringe and immediately measured at a gas chromatograph (SRI 8610C, SRI Instruments, USA). At the time point of gas injection, air temperature, air pressure, and inner pressure of the incubation jar were noted for each sample at each measurement time point. The measured $\text{CO}_2\text{-C}$ concentration in parts per million was converted into $\text{CO}_2\text{-C}$ by taking into account the sum of air pressure and inner pressure of the incubation jar (p), the molar mass of C ($M_C = 12.01 \text{ g mol}^{-1}$), the air temperature (T), the universal gas constant ($R = 8.314 \text{ [(kg m}^2\text{) (s}^2\text{ mol K)}^{-1}]$), the conversion of [g] into [mg], and the ideal gas law using the following Eq. (1).

$$\text{CO}_2\text{-C [mg m}^{-3}] = \frac{p \text{ [kg (m s}^{-2}\text{)}^{-1}] \cdot M_C \text{ [g mol}^{-1}] \cdot \text{CO}_2 \text{ [ppm]}}{R \text{ [(kg m}^2\text{) (s}^2\text{ mol K)}^{-1}] \cdot T \text{ [K]}} \cdot 1000 \quad (1)$$

The soil respiration rate was calculated across the linear increase in the CO_2 concentration over the 6-week incubation period using a linear regression. Afterwards, the soil respiration rate per mass of the total sample [$\text{mg CO}_2\text{-C m}^{-3} \text{ d}^{-1}$] was calculated for the volume of headspace (Vol_{HS}) and the amount of dry soil (soil_{dw}) within each incubation jar, considering the proportion of the fine soil fraction ($\text{sf}_{<2\text{mm}}$) in the total soil ($\text{sf}_{>2\text{mm}} + \text{sf}_{<2\text{mm}}$) and including the conversion of [mg] into [μg] using Eq. (2).

$$\text{CO}_2\text{-C [\mu g (g d)}^{-1}] = \frac{\text{CO}_2\text{-C [mg (m}^3 \text{ d)}^{-1}] \cdot \text{Vol}_{\text{HS}} \text{ [m}^3\text{]} \cdot \left(\frac{\text{sf}_{<2\text{mm}} \text{ [g]}}{\text{sf}_{>2\text{mm}} \text{ [g]} + \text{sf}_{<2\text{mm}} \text{ [g]}} \right)}{\text{soil}_{\text{dw}} \text{ [g]}} \cdot 1000 \quad (2)$$

The Vol_{HS} was calculated as the difference between the total volume of the incubation jar (350 mL) and the volume of soil at 60 % WHC.

2.6 Extraction and quantification of DNA

Total DNA was extracted from 400 mg of field-moist soil (< 2 mm) using a kit (FastDNA Spin Kit for Soil, MP Biomedicals, Solon, USA) following the instructions of the producer. The volume of the extracted DNA eluate was determined, and afterwards the DNA was quantified using a PicoGreen assay (Quant-iT PicoGreen dsDNA Assay Kit, Invitrogen, Life Technologies Corporation, Eugene, OR, USA) measured at a fluorescence microplate reader (FLx800, BioTek, Winooski, VT, USA). By considering the proportion of the fine soil fraction ($\text{sf}_{<2\text{mm}}$) in the total soil ($\text{sf}_{<2\text{mm}} + \text{sf}_{>2\text{mm}}$) and the dry-weight equivalent of the soil, the amount of DNA per mass of the total sample was calculated.

2.7 Radiocarbon (^{14}C) ratio of respired $\text{CO}_2\text{-C}$ and soil TOC

To determine the radiocarbon (^{14}C) ratio of respired $\text{CO}_2\text{-C}$, fine soil (< 2 mm) subsamples were incubated in 1050 mL incubation jars with septum. For the arid study site, between 100–340 g soil (dry-weight equivalent) was incubated for the different soil increments of the three profiles. For the mediterranean and humid site, the amount of soil (dry weight equivalent) used for the incubation differed between soil depth increments: 180 g (humid) or 250 g (mediterranean) for 0–10 cm, 240 g (humid) or 280 g (mediterranean) for 10–200 cm, 260 g (humid), or 330 g (mediterranean) for 200–600 cm. Immediately after adjusting the soil water content to 60 % WHC, the incubation jars were closed air-tight and flushed with synthetic air (hydrocarbon free, Rießner-Gase GmbH, Lichtenfels, Germany) for ~ 6 min with a flow rate of $\sim 700 \text{ cm}^3 \text{ min}^{-1}$ to remove the atmospheric CO_2 within the incubation jars. Afterwards the samples were incubated at 20 °C in the dark, and the CO_2 concentration was monitored. Once a minimum amount of $\sim 1 \text{ mg CO}_2\text{-C}$ within the headspace of the incubation jar was reached, the gas in the headspace was transferred into a 400 mL vacuum container (miniature air sampling canister, Restek, Bellefonte, USA) for storage until further processing. For samples with a minimum amount of only $\sim 0.5 \text{ mg CO}_2\text{-C}$ within the headspace of the incubation jar, the CO_2 was directly extracted from the incubation jar.

The ^{14}C analyses of the respired $\text{CO}_2\text{-C}$ and the soil TOC were conducted using the accelerator mass spectrometry (AMS) facility in Jena, Germany (Steinhof et al., 2004). From the storage container or directly from the incubation jar, the respired CO_2 was transferred into a glass tube cooled by liquid nitrogen and containing an iron catalyst. The following reduction of CO_2 to graphite was carried out in the presence of hydrogen (H_2) at 600 °C. The resulting graphite-coated iron was pressed into targets and measured for ^{14}C using the AMS MICADAS (Ionplus, Dietikon, Switzerland). To determine the ^{14}C ratio of soil TOC samples, milled soil subsamples were combusted and graphitized following Steinhof et al. (2017). The radiocarbon ratio is reported as $\Delta^{14}\text{C}$ in per mil [‰], which is the fraction with respect to the standard isotope ratio (oxalic acid standard SRM-4990C; Steinhof et al., 2017) including the normalization for $\delta^{13}\text{C}$ (fractionation correction) and the correction for the decay between 1950 and the measurement time (2020/21; Stuiver and Polach, 1977). The analytical uncertainty is 2 ‰ .

We refrain from presenting radiocarbon ages in years because in open systems, such as soils, radiocarbon is continuously exchanged with the atmosphere and traditional radiocarbon dating cannot be done (Trumbore et al., 2016). A mean age estimate using the available radiocarbon data would require fitting a compartmental model to the data, but there were not enough data, for instance about C input that

covers the relevant timescale (last 1000 years), to constrain such a model.

2.8 Statistical analyses

Data were analyzed for significant differences between sites and soil depths and their interaction effects, using a two-way ANOVA followed by Tukey's post hoc test ("Tukey adjusted pairwise comparison" using the R function *emmeans*). For this purpose, the residuals of the two-way ANOVA model were tested for normality using the Shapiro–Wilk test and for variance homogeneity using Levene's test. In cases in which the assumptions of normality and variance homogeneity were violated the data were rank-transformed. In addition, we tested whether TOC and microbial-derived CO₂–C from the same soil depth increment differ significantly in their ¹⁴C signature in specific soil depth increments since differences might potentially only occur in specific depth ranges. Differences between radiocarbon data (¹⁴C) of the soil total organic carbon (TOC) and respired CO₂–C for individual soil depth increments per site were analyzed using Welch's *t* test (in the case of normality) or the Wilcoxon rank sum test (in the case of violation of normality). For soil depth increments with a low number of observations (*n*), several depth increments were combined (as indicated in the results). For all analyses, *p* < 0.05 was considered the threshold for statistical significance. All statistical analyses were conducted in R (version 4.2.2, R Core Team, 2022).

3 Results

The $\Delta^{14}\text{C}$ of the TOC was significantly lower in soils of the arid site than in soils of the mediterranean and humid site (Fig. 2a and Table S2). The $\Delta^{14}\text{C}$ of the soil TOC decreased significantly with increasing soil depth at all three sites (yet the interaction effect between site and soil depth was not statistically significant; Table S2). In the uppermost 160 cm, the mean $\Delta^{14}\text{C}$ of the TOC decreased on average from -103‰ to -761‰ at the arid site, from 70‰ to -267‰ at the mediterranean site, and from -30‰ to -469‰ at the humid site (Fig. 2a). The $\Delta^{14}\text{C}$ of the CO₂ respired by soil microorganisms also decreased with depth, but less than the $\Delta^{14}\text{C}$ of the soil TOC (Fig. 2b). In the uppermost 1 m of the soils, the $\Delta^{14}\text{C}$ of the respired CO₂–C tended to be positive, while it was negative at depth > 1 m. The respired CO₂–C at a depth of 200–600 cm had a higher $\Delta^{14}\text{C}$, and thus was younger, at the humid than at the mediterranean site (Fig. 2b). The $\Delta^{14}\text{C}$ of the CO₂ respired by soil microorganisms was significantly higher than the $\Delta^{14}\text{C}$ of the TOC in the same depth increment in the soils of all three sites (Fig. 2), indicating that the respired C is younger than the soil TOC. These differences were statistically significant in most depth increments of the humid and arid site and in one depth increment of the soil at the mediterranean site (Fig. 2b and c). The slope of the linear

model that describes the $\Delta^{14}\text{C}$ of the respired C as a function of the $\Delta^{14}\text{C}$ of the TOC was 0.15 for the Mediterranean site, 0.23 for the arid site, and 0.26 for the humid site (Fig. 2c).

The $\delta^{13}\text{C}$ of the TOC differed significantly between sites, and also the interaction effect of site and soil depth was statistically significant (Fig. 3a and Table S2). The $\delta^{13}\text{C}$ of the TOC increased at the humid site from a mean of $-28.7 \pm 0.2\text{‰}$ in the organic layer to $-23.2 \pm 0.2\text{‰}$ in the depth section 160–180 cm (Fig. 3a). At the mediterranean site, it increased from $-27.2 \pm 0.6\text{‰}$ in the organic layer to $-23.1 \pm 0.3\text{‰}$ at a depth of 40–60 cm and did not change consistently with depth in the subsoil. At the arid site, the $\delta^{13}\text{C}$ of the TOC hardly changed in the whole soil profile (Fig. 3a). The $\delta^{13}\text{C}$ of the roots did not change significantly with depth (Fig. 3b and Table S2). The $\delta^{13}\text{C}$ of the roots increased in the order humid < mediterranean < arid in all depth increments (Fig. 3b), similar to the $\delta^{13}\text{C}$ of the TOC (Fig. 3a).

The concentration of DNA in the soils increased in the order arid < mediterranean < humid across all depth increments (Fig. 4). In all soils, the DNA concentration decreased strongly with depth (Fig. 4 and Table S2). Similarly, the TOC concentration increased in the order arid < mediterranean < humid across all depth increments (Fig. 5). The TOC concentration decreased by 93 % at the arid and by 96 % at the mediterranean and humid site, respectively, in the uppermost 200 cm from top to bottom (Fig. 5). The microbial respiration rate in the soils of the humid and mediterranean site was higher than at the arid site in all depth sections (Fig. S1 in the Supplement and Table S2). The microbial respiration rates decreased by more than 99 % at all three sites in the uppermost 200 cm from top to bottom (Fig. S1).

4 Discussion

4.1 Microorganisms live on young carbon, even several meters below the soil surface

We found that microbial respiration in the soils of all three climate zones is fueled by modern C in the uppermost 1 m of the soils and by relatively young C, as compared to the TOC, below 1 m depth (Fig. 2a, b, c). Most C that is respired by microorganisms in the upper part of the soil is likely directly derived from roots. This is consistent with the fact that annual dead root biomass input to soil makes up about 10 % and 34 % of the fine root biomass in forests and shrublands, respectively (Gill and Jackson, 2000), while root exudates amount to 20 %–40 % of the photosynthetically fixed C (Badri and Vivanco, 2009). This is further supported by studies showing that rhizodeposition is an important driver of microbial respiration (Phillips et al., 2013). Although roots are absent below 160 cm (arid site) and 350 cm (humid site) (Fig. 3b), the C respired by microorganisms in the deep soil




Figure 2. Radiocarbon ratio ($\Delta^{14}\text{C}$; mean \pm standard error) of (a) soil total organic carbon (TOC) and (b) respired CO_2 -C as well as (c) their relationship in soils at three sites (arid, mediterranean, humid) located along a precipitation gradient in the Coastal Cordillera of Chile (0–200 cm: $n = 3$, except for humid site with $n = 4$; > 200 cm: $n = 1$, except for humid site 200–400 cm with $n = 2$). Different letters in (a) and (b) indicate significant differences ($p \leq 0.05$) among the sites tested by Tukey's post hoc test after two-way ANOVA. Asterisks indicate significant differences between the radiocarbon ratio of the total soil organic carbon (TOC) and that of the respired CO_2 -C for each site, tested across all soil depth increments (see legend) and also separately for the different sites and depth increments (* $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$). The colors of the asterisks indicate the site the asterisks refer to. Lines in (a) and (b) indicate which soil depth increments were combined (separately per site) for statistical analysis.




Figure 3. ^{13}C ratio ($\delta^{13}\text{C}$; mean \pm standard error) of (a) soil total organic carbon (TOC) and (b) roots at three sites (arid, mediterranean, humid) located along a precipitation gradient in the Coastal Cordillera of Chile (0–200 cm: $n = 3$, except for humid site with $n = 4$; 200–600 cm: $n = 1$, except for humid site 200–400 cm with $n = 2$). Different letters in (a) and (b) indicate significant differences ($p \leq 0.05$) among the sites tested by Tukey’s post hoc test after two-way ANOVA. Lines in (a) and (b) indicate which soil depth increments were combined (separately per site) for statistical analysis.

is still relatively young (Fig. 2b, c), suggesting that microbial activity in the deep soil is mostly fueled by young dissolved organic carbon (DOC) that is rapidly transported downwards in the soil. Indeed, studies have shown that DOC fluxes from the top to the subsoil can be as high as $200 \text{ kg ha}^{-1} \text{ yr}^{-1}$ (Michalzik et al., 2001). Notably, the respired $\text{CO}_2\text{-C}$ at a depth of 200–600 cm tends to be younger at the humid than at the mediterranean site (Fig. 2b and c). As the humid site experiences higher precipitation and has a larger C content in the topsoil than the other two sites, it is likely that the transport of young C from the topsoil to the deep soil is the largest at this site. This possibility is supported by a meta-analysis that found a positive correlation between precipitation and DOC fluxes from the topsoil to the subsoil (Michalzik et al., 2001).

Our findings suggest that microorganisms throughout the soil, even several meters below the soil surface, metabolize organic C that is relatively young and substantially younger than the soil TOC. We interpret these findings to indicate preferential metabolism of organic C that is directly derived from roots or has been transported rapidly downwards from the topsoil to deeper soil horizons as DOC (Sanderman et al., 2008; Rumpel and Kögel-Knabner, 2011; Kaiser and Kalbitz, 2012; Phillips et al., 2013; Marín-Spiotta and Hobbey,




Figure 4. Total DNA content (mean \pm standard error) of soil depth increments down to (a) 100 cm, (b) 200 cm, and (c) 600 cm depth at three sites (arid, mediterranean, humid) located along a precipitation gradient in the Coastal Cordillera of Chile (0–200 cm: $n = 3$, except for humid site with $n = 4$; > 200 cm: $n = 1$, except for humid site 200–400 cm with $n = 2$). Please note the differences in the scale of the x axes with a logarithmic scale for (a) and (b) and a linear scale for (c). Different letters indicate significant differences ($p \leq 0.05$) between the three sites tested by Tukey’s post hoc test after two-way ANOVA.




Figure 5. Total organic carbon (TOC) content (mean \pm standard error) of soil depth increments down to (a) 100 cm, (b) 200 cm, and (c) 600 cm depth at three sites (arid, mediterranean, humid) located along a precipitation gradient in the Coastal Cordillera of Chile (0–200 cm: $n = 3$, except for humid site with $n = 4$; > 200 cm: $n = 1$, except for humid site 200–400 cm with $n = 2$). Different letters in both (a) and (b) indicate significant differences ($p \leq 0.05$) among the sites tested by Tukey's post hoc test after two-way ANOVA. Lines in (a) and (c) indicate which soil depth increments were combined (separately per site) for statistical analysis.

2022). The young organic matter input to soil is very likely not yet stabilized against microbial decomposition by sorption to mineral surfaces. In addition, they are likely more carbohydrate-rich and thus energy-rich compared to organic matter that has already been processed by the microbial community (Ni et al., 2020). Yet, it has to be considered that the soil was sieved before its incubation. Sieving destroys soil aggregates and potentially renders organic C available to microorganisms that was previously protected in microaggregates. Thus, it is likely that under in situ conditions, the CO₂-C that is respired by microorganisms is even younger (and thus has an even higher $\Delta^{14}\text{C}$) than in our incubation experiment.

Our findings are in accord with studies about topsoil, such as Trumbore (2000), who reported that whereas TOC in boreal, temperate, and tropical forest topsoils was between 200 and 1200 years old, the CO₂-C respired by heterotrophic bacteria in these soil types was only 30, 8, and 3 years old, respectively. Our study is the first to report the $\Delta^{14}\text{C}$ of CO₂ respired by soil microorganisms below 1.0 m in non-permafrost soils and shows that microbial processes in deep soil are closely connected to recently fixed CO₂ from the atmosphere. This is in contrast to permafrost soils, in which microbial respiration during large parts of the year relies strongly on C that has persisted in these soils for centuries (Dutta et al., 2006; Lee et al., 2012; Gentsch et al., 2018; Pedron et al., 2022). Owing to the replication of this study in three climate zones, our result that microbial activity in the deep soil of non-permafrost soils is fueled mostly by young organic matter may be generalizable across a large part of the terrestrial biosphere.

4.2 Strong decomposition of soil organic matter only occurs in the upper decimeters of the soils

While soil organic matter moves downwards slowly, it interacts with the soil matrix and gets partly decomposed by microorganisms (Sanderman et al., 2008; Kaiser and Kalbitz, 2012). Microbial enzymes preferably decompose organic compounds with a high proportion of the lighter ¹²C, which leads to enrichment of the heavier ¹³C isotope in soils (Ehleringer et al., 2000; Wynn et al., 2006; Balesdent et al., 2018). In the Coastal Cordillera soils, we found that intensive decomposition of organic matter that leads to ¹³C enrichment was restricted to the upper decimeters of the soils (Fig. 3a). Decomposition is likely restricted to the topsoil because organic matter becomes increasingly stabilized against microbial decomposition over time due to occlusion and sorption to mineral surfaces, which renders the organic compounds spatially inaccessible (Kögel-Knabner et al., 2008; Schmidt et al., 2011; Dungait et al., 2012). In addition, intensive decomposition of organic matter might also be restricted to the uppermost decimeters of the soils because the energy content of the soil organic matter declines with increasing soil depth (Ni et al., 2020). Further, it could be

that recently fixed carbon that enters the soil in the upper decimeters also leads to priming and thus intensive decomposition in the topsoil (Bernard et al., 2022). In addition, it is possible that the change in the carbon stable isotope ratio is partly related to the dilution of atmospheric ^{13}C - CO_2 by ^{13}C -depleted CO_2 derived from burning of fossil fuels (i.e., “Suess effect”; Keeling et al., 1979).

At the arid site, there is no enrichment of ^{13}C in the soil TOC with increasing soil depth (Fig. 3a), indicating that little organic matter processing by microorganisms occurs in this soil, probably because of the lack of moisture which hampers decomposition (Moyano et al., 2013; Seuss et al., 2022). At the mediterranean and the humid sites, the $\delta^{13}\text{C}$ of the soil TOC increases with increasing depth down to 80 and 180 cm, respectively (Fig. 3a), suggesting that microbial processing of organic matter occurred at greater depths at the humid site than at the mediterranean site, which is likely because of the higher soil moisture at the humid site than at the mediterranean site. In addition, the higher decomposition of organic matter at greater depth at the humid site compared to the other sites might also be related to the trees at this site that seem to cause a larger root density in deeper soil horizons than the vegetation at the other sites and thus likely stimulate microbial activity in the subsoil.

4.3 The age of total soil organic carbon increases with soil depth and aridity

While the soil TOC content decreases with soil depth and aridity of the sites (Fig. 5), the age of the TOC increases (Fig. 2a), suggesting that deep soils in an arid climate can retain C for long periods of time. Our findings agree with a meta-analysis (Mathieu et al., 2015) reporting that soils in an arid climate have a lower ^{14}C abundance in the topsoil and, particularly, in the deeper layers than soils in a humid climate. Consistent with our results, the mean differences in $\Delta^{14}\text{C}$ among dry and humid climate in that study amounted to 40‰ in the topsoils and 100‰ in the subsoils, indicating that the soil TOC in arid areas is older than in humid areas, particularly in the subsoil (Mathieu et al., 2015). This is partly because the microbial community in the subsoil feeds primarily on young DOC and not on old pools of the TOC (see above). Another reason for why the soil TOC at the arid site is particularly old is likely the low microbial biomass (Fig. 4) and activity (Fig. S1). This is supported by in situ measurements of the soil CO_2 concentration at the arid site at different soil depths that detected low CO_2 concentrations throughout the year, suggesting that microbial activity and organic matter decomposition is low at this site (Spohn and Holzheu, 2021).

4.4 Conclusions

Based on the unique measurements of microbially respired ^{14}C - CO_2 down to a soil depth of 6 m in different climate zones, this study reveals that microbial activity several meters below the soil surface is fueled by recently fixed C and that strong microbial decomposition of the soil TOC only occurs in the upper decimeters of the soils. Thus, in contrast to deep permafrost soils, in which microorganisms respire old C during large parts of the year, microorganisms in deep, non-permafrost soils feed on recent C input. Our results suggest that deeply developed soils can restrain C from the atmosphere for climate-relevant periods of time because the microbial community in the subsoil mostly feeds on young carbon derived from roots and DOC. Taken together, our results show that different layers of the critical zone are tightly connected and that processes in the deep soil depend on carbon that recently entered the ecosystem through fixation of CO_2 from the atmosphere.

Data availability. Data are available here: <https://doi.org/10.5281/zenodo.7389812> (Scheibe et al., 2022).

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/bg-20-827-2023-supplement>.

Author contributions. MS and CAS conceptualized the study, MS and AS conducted the fieldwork, AS conducted the lab work, AS did the data analysis with input from MS and CAS, MS wrote the manuscript, AS and CAS contributed to the manuscript, and MS acquired funding for the project.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Special issue statement. This article is part of the special issue “Earth surface shaping by biota (Esurf/BG/SOIL/ESD/ESSD inter-journal SI)”. It is not associated with a conference.

Acknowledgements. We are grateful to the Chilean National Park Service (CONAF) for providing access to the sample locations and on-site support for our research. We thank the whole Deep Earth-Shape team for collaboration during the field sampling campaigns. We thank Renate Krauß and Harald Isreal Suaznabar Olguin for their technical support in the laboratory. We acknowledge the Laboratory of Isotope Biogeochemistry at the University of Bayreuth

for stable isotope measurements. We acknowledge the support from the ^{14}C Analysis Facility at the Max Planck Institute for Biogeochemistry in Jena, and we thank Axel Steinhof and Manuel Rost for their technical support. The study was funded by the German Science Foundation (DFG) as part of the priority research program SPP-1803 “EarthShape: Earth Surface Shaping by Biota” (grant no. DFG SP1389/5-2).

Financial support. This research has been supported by the Deutsche Forschungsgemeinschaft (grant no. SP1389/5-2).

This open-access publication was funded by the University of Bayreuth.

Review statement. This paper was edited by Sara Vicca and Serita Frey and reviewed by Karis McFarlane and Hannah Holland-Moritz.

References

- Akob, D. M. and Küsel, K.: Where microorganisms meet rocks in the Earth’s Critical Zone, *Biogeosciences*, 8, 3531–3543, <https://doi.org/10.5194/bg-8-3531-2011>, 2011.
- Badri, D. V. and Vivanco, J. M.: Regulation and function of root exudates, *Plant Cell Environ.*, 32, 666–681, <https://doi.org/10.1111/j.1365-3040.2009.01926.x>, 2009.
- Balesdent, J., Basile-Doelsch, I., Chadoeuf, J., Cornu, S., Derrien, D., Fekiacova, Z., and Hatté, C.: Atmosphere–soil carbon transfer as a function of soil depth, *Nature*, 559, 599–602, <https://doi.org/10.1038/s41586-018-0328-3>, 2018.
- Berg, A. and Banwart, S. A.: Carbon dioxide mediated dissolution of Ca-feldspar: implications for silicate weathering, *Chem. Geol.*, 163, 25–42, [https://doi.org/10.1016/S0009-2541\(99\)00132-1](https://doi.org/10.1016/S0009-2541(99)00132-1), 2000.
- Berner, R. A.: The rise of plants and their effect on weathering and atmospheric CO_2 , *Science*, 276, 544–546, <https://doi.org/10.1126/science.276.5312.544>, 1997.
- Bernard, L., Basile-Doelsch, I., Derrien, D., Fanin, N., Fontaine, S., Guenet, B., Karimi, B., Marsden, C., and Maron, P.-A.: Advancing the mechanistic understanding of the priming effect on soil organic matter mineralization, *Funct. Ecol.*, 36, 1355–1377, <https://doi.org/10.1111/1365-2435.14038>, 2022.
- Bernhard, N., Moskwa, L.-M., Schmidt, K., Oeser, R. A., Aburto, F., Bader, M. Y., Baumann, K., von Blanckenburg, F., Boy, J., van den Brink, L., Brucker, E., Büdel, B., Canessa, R., Dippold, M. A., Ehlers, T. A., Fuentes, J. P., Godoy, R., Jung, P., Karsten, U., Köster, M., Kuzyakov, Y., Leinweber, P., Neidhardt, H., Matus, F., Mueller, C. W., Oelmann, Y., Osés, R., Osses, P., Paulino, L., Samolov, E., Schaller, M., Schmid, M., Spielvogel, S., Spohn, M., Stock, S., Stroncik, N., Tielbörger, K., Übernickel, K., Scholten, T., Seguel, O., Wagner, D., and Kühn, P.: Pedogenic and microbial interrelations to regional climate and local topography: New insights from a climate gradient (arid to humid) along the Coastal Cordillera of Chile, *Catena*, 170, 335–355, <https://doi.org/10.1016/j.catena.2018.06.018>, 2018.
- Brucker, E. and Spohn, M.: Formation of soil phosphorus fractions along a climate and vegetation gradient in the Coastal Cordillera of Chile, *Catena*, 180, 203–211, <https://doi.org/10.1016/j.catena.2019.04.022>, 2019.
- Dungait, J. A., Hopkins, D. W., Gregory, A. S., and Whitmore, A. P.: Soil organic matter turnover is governed by accessibility not recalcitrance, *Glob. Change Biol.*, 18, 1781–1796, <https://doi.org/10.1111/j.1365-2486.2012.02665.x>, 2012.
- Dutta, K., Schuur, E. A. G., Neff, J. C., and Zimov, S. A.: Potential carbon release from permafrost soils of Northeastern Siberia, *Glob. Change Biol.*, 12, 2336–2351, <https://doi.org/10.1111/j.1365-2486.2006.01259.x>, 2006.
- Ehleringer, J. R., Buchmann, N., and Flanagan, L. B.: Carbon isotope ratios in belowground carbon cycle processes, *Ecol. Appl.*, 10, 412–422, [https://doi.org/10.1890/1051-0761\(2000\)010\[0412:CIRIBC\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0412:CIRIBC]2.0.CO;2), 2000.
- Fierer, N., Chadwick, O. A., and Trumbore, S. E.: Production of CO_2 in soil profiles of a California annual grassland, *Ecosystems*, 8, 412–429, <https://doi.org/10.1007/s10021-003-0151-y>, 2005.
- Finlay, R. D., Mahmood, S., Rosenstock, N., Bolou-Bi, E. B., Köhler, S. J., Fahad, Z., Rosling, A., Wallander, H., Belyazid, S., Bishop, K., and Lian, B.: Reviews and syntheses: Biological weathering and its consequences at different spatial levels – from nanoscale to global scale, *Biogeosciences*, 17, 1507–1533, <https://doi.org/10.5194/bg-17-1507-2020>, 2020.
- Gentsch, N., Wild, B., Mikutta, R., Čapek, P., Diáková, K., Schrupf, M., Turner, S., Minnich, C., Schaarschmidt, F., Shibistova, O., Schneckner, J., Ulrich, T., Gittel, A., Šantrůčková, H., Bárta, J., Lashchinskiy, N., Fuß, R., Richter, A. and Guggenberger, G.: Temperature response of permafrost soil carbon is attenuated by mineral protection, *Glob. Change Biol.*, 24, 3401–3415, <https://doi.org/10.1111/gcb.14316>, 2018.
- Gill, R. A. and Jackson, R. B.: Global patterns of root turnover for terrestrial ecosystems, *New Phytol.*, 147, 13–31, <https://doi.org/10.1046/j.1469-8137.2000.00681.x>, 2000.
- Hanson, P. J., Edwards, N. T., Garten, C. T., and Andrews, J. A.: Separating root and soil microbial contributions to soil respiration: a review of methods and observations, *Biogeochemistry*, 48, 115–146, <https://doi.org/10.1023/A:1006244819642>, 2000.
- Hayes, N. R., Buss, H. L., Moore, O. W., Krám, P., and Pancost, R. D.: Controls on granitic weathering fronts in contrasting climates, *Chem. Geol.*, 535, 119450, <https://doi.org/10.1016/j.chemgeo.2019.119450>, 2020.
- Hoffland, E., Kuyper, T. W., Wallander, H., Plassard, C., Gorbushina, A. A., Haselwandter, K., Holmström, S., Landeweert, R., Lundström, U. S., Rosling, A., Sen, R., Smits, M. M., van Hees, P. A. W., and van Breemen, N.: The role of fungi in weathering, *Front. Ecol. Environ.*, 2, 258–264, [https://doi.org/10.1890/1540-9295\(2004\)002\[0258:TROFIW\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2004)002[0258:TROFIW]2.0.CO;2), 2004.
- Högberg, P. and Read, D. J.: Towards a more plant physiological perspective on soil ecology, *Trends Ecol. Evol.*, 21, 548–554, <https://doi.org/10.1016/j.tree.2006.06.004>, 2006.
- Högberg, P., Högberg, M. N., Göttlicher, S. G., Betson, N. R., Keel, S. G., Metcalfe, D. B., Campbell, C., Schindlbacher, A., Hurry, V., Lundmark, T., Linder, S., and Näsholm, T.: High temporal resolution tracing of photosynthate carbon from the tree

- canopy to forest soil microorganisms, *New Phytol.*, 177, 220–228, <https://doi.org/10.1111/j.1469-8137.2007.02238.x>, 2008.
- Hulton, N. R. J., Purves, R. S., McCulloch, R. D., Sugden, D. E., and Bentley, M. J.: The last Glacial Maximum and deglaciation in southern South America, *Quaternary Sci. Rev.*, 21, 233–241, [https://doi.org/10.1016/S0277-3791\(01\)00103-2](https://doi.org/10.1016/S0277-3791(01)00103-2), 2002.
- IUSS Working Group WRB: World Reference Base for Soil Resources 2014, update 2015, International soil classification system for naming soils and creating legends for soil maps, World Soil Resources Reports No. 106, FAO, Rome, 2015.
- Jackson, R. B., Lajtha, K., Crow, S. E., Hugelius, G., Kramer, M. G., and Piñeiro, G.: The ecology of soil carbon: pools, vulnerabilities, and biotic and abiotic controls, *Annu. Rev. Ecol. Evol. S.*, 48, 419–445, <https://doi.org/10.1146/annurev-ecolsys-112414-054234>, 2017.
- Kaiser, K. and Kalbitz, K.: Cycling downwards—dissolved organic matter in soils, *Soil Biol. Biochem.*, 52, 29–32, <https://doi.org/10.1016/j.soilbio.2012.04.002>, 2012.
- Keeling, C. D., Mook, W. G., and Tans, P. P.: Recent trends in the $^{13}\text{C}/^{12}\text{C}$ ratio of atmospheric carbon dioxide, *Nature*, 277, 121–123, <https://doi.org/10.1038/277121a0>, 1979.
- Kögel-Knabner, I., Guggenberger, G., Kleber, M., Kandeler, E., Kalbitz, K., Scheu, S., Eusterhues, K., and Leinweber, P.: Organo-mineral associations in temperate soils: Integrating biology, mineralogy, and organic matter chemistry, *J. Plant Nutr. Soil Sc.*, 171, 61–82, <https://doi.org/10.1002/jpln.200700048>, 2008.
- Krone, L. V., Hampl, F. J., Schwerdtelm, C., Bryce, C., Ganzert, L., Kitte, A., Übernickel, K., Dielforder, A., Aldaz, S., Oses-Pedraza, R., Perez, J. P. H., Sanchez-Alfaro, P., Wagner, D., Weckmann, U., and von Blanckenburg, F.: Deep weathering in the semi-arid Coastal Cordillera, Chile, *Sci. Rep.*, 11, 1–15, <https://doi.org/10.1038/s41598-021-90267-7>, 2021.
- Krumholz, L. R.: Microbial communities in the deep subsurface, *Hydrogeol. J.*, 8, 4–10, <https://doi.org/10.1007/s1004400050003>, 2000.
- Lawrence, C. R., Beem-Miller, J., Hoyt, A. M., Monroe, G., Sierra, C. A., Stoner, S., Heckman, K., Blankinship, J. C., Crow, S. E., McNicol, G., Trumbore, S., Levine, P. A., Vinduřková, O., Todd-Brown, K., Rasmussen, C., Hicks Pries, C. E., Schädel, C., McFarlane, K., Doetterl, S., Hatté, C., He, Y., Treat, C., Harden, J. W., Torn, M. S., Estop-Aragonés, C., Asefaw Berhe, A., Keiluweit, M., Della Rosa Kuhnen, Á., Marin-Spiotta, E., Plante, A. F., Thompson, A., Shi, Z., Schimel, J. P., Vaughn, L. J. S., von Fromm, S. F., and Wagai, R.: An open-source database for the synthesis of soil radiocarbon data: International Soil Radiocarbon Database (ISRaD) version 1.0, *Earth Syst. Sci. Data*, 12, 61–76, <https://doi.org/10.5194/essd-12-61-2020>, 2020.
- Lee, H., Schuur, E. A., Inglett, K. S., Lavoie, M., and Chanton, J. P.: The rate of permafrost carbon release under aerobic and anaerobic conditions and its potential effects on climate, *Glob. Change Biol.*, 18, 515–527, <https://doi.org/10.1111/j.1365-2486.2011.02519.x>, 2012.
- Luebert, F. and Plissock, P.: Sinopsis bioclimática y vegetacional de Chile, 2nd Edn., Editorial Universitaria, Santiago, Chile, 384 pp., 2017.
- Marín-Spiotta, E. and Hobbey, E. U.: Deep soil carbon, in: Multi-Scale Biogeochemical Processes in Soil Ecosystems: Critical Reactions and Resilience to Climate Changes, edited by: Yang, Y., Keiluweit, M., Senesi, N., and Xing B., John Wiley & Sons, Inc., 193–206, <https://doi.org/10.1002/9781119480419.ch9>, 2022.
- Marin-Spiotta, E., Chaopricha, N. T., Plante, A. F., Diefendorf, A. F., Mueller, C. W., Grandy, A. S., and Mason, J. A.: Long-term stabilization of deep soil carbon by fire and burial during early Holocene climate change, *Nat. Geosci.*, 7, 428–432, <https://doi.org/10.1038/NGEO2169>, 2014.
- Mathieu, J. A., Hatté, C., Balesdent, J., and Parent, É.: Deep soil carbon dynamics are driven more by soil type than by climate – a worldwide meta-analysis of radiocarbon profiles, *Glob. Change Biol.*, 21, 4278–4292, <https://doi.org/10.1111/gcb.13012>, 2015.
- Michalzik, B., Kalbitz, K., Park, J.-H., Solinger, S., and Matzner, E.: Fluxes and concentrations of dissolved organic carbon and nitrogen – a synthesis for temperate forests, *Biogeochemistry*, 52, 173–205, <https://doi.org/10.1023/A:1006441620810>, 2001.
- Moreland, K., Tian, Z., Berhe, A. A., McFarlane, K. J., Hartsoogh, P., Hart, S. C., Bales, R., and O’Geen, A. T.: Deep in the Sierra Nevada critical zone: saprock represents a large terrestrial organic carbon stock, *Environ. Res. Lett.*, 16, 124059, <https://doi.org/10.1088/1748-9326/ac3bfe>, 2021.
- Moyano, F. E., Manzoni, S., and Chenu, C.: Responses of soil heterotrophic respiration to moisture availability: An exploration of processes and models, *Soil Biol. Biochem.*, 59, 72–85, <https://doi.org/10.1016/j.soilbio.2013.01.002>, 2013.
- Nagy, R. C., Porder, S., Brando, P., Davidson, E. A., Figueira, A. M. E. S., Neill, C., Riskin, S., and Trumbore, S.: Soil carbon dynamics in soybean cropland and forests in Mato Grosso, Brazil, *J. Geophys. Res.-Biogeo.*, 123, 18–31, <https://doi.org/10.1002/2017JG004269>, 2018.
- Ni, X., Liao, S., Tan, S., Peng, Y., Wang, D., Yue, K., Wu, F., and Yang, Y.: The vertical distribution and control of microbial necromass carbon in forest soils, *Global Ecol. Biogeogr.*, 29, 1829–1839, <https://doi.org/10.1111/geb.13159>, 2020.
- Oeser, R. A., Stroncik, N., Moskwa, L.-M., Bernhard, N., Schaller, M., Canessa, R., van den Brink, L., Köster, M., Brucker, E., Stock, S., Fuentes, J. P., Godoy, R., Matus, F. J., Pedraza, R. O., McIntyre, P. O., Paulino, L., Seguel, O., Bader, M. Y., Boy, J., Dippold, M. A., Ehlers, T. A., Kühn, P., Kuzyakov, Y., Leinweber, P., Scholten, T., Spielvogel, S., Spohn, M., Übernickel, K., Tielbörger, K., Wagner, D., and von Blanckenburg, F.: Chemistry and microbiology of the Critical Zone along a steep climate and vegetation gradient in the Chilean Coastal Cordillera, *Catena*, 170, 183–203, <https://doi.org/10.1016/j.catena.2018.06.002>, 2018.
- Pedersen, K.: Microbial life in deep granitic rock, *FEMS Microbiol. Rev.*, 20, 399–414, <https://doi.org/10.1111/j.1574-6976.1997.tb00325.x>, 1997.
- Pedron, S. A., Welker, J. M., Euskirchen, E. S., Klein, E. S., Walker, J. C., Xu, X., and Czimczik, C. I.: Closing the winter gap – Year-round measurements of soil CO₂ emission sources in Arctic tundra, *Geophys. Res. Lett.*, 49, e2021GL097347, <https://doi.org/10.1029/2021GL097347>, 2022.
- Phillips, C. L., McFarlane, K. J., Risk, D., and Desai, A. R.: Biological and physical influences on soil ^{14}C seasonal dynamics in a temperate hardwood forest, *Biogeosciences*, 10, 7999–8012, <https://doi.org/10.5194/bg-10-7999-2013>, 2013.
- R Core Team: A Language and Environment for Statistical Computing R Foundation for Statistical Computing, Austria, Vienna,

- <https://www.R-project.org/> (last access: 14 February 2023), 2022.
- Rumpel, C. and Kögel-Knabner, I.: Deep soil organic matter – a key but poorly understood component of terrestrial C cycle, *Plant Soil*, 338, 143–158, <https://doi.org/10.1007/s11104-010-0391-5>, 2011.
- Sanderman, J., Baldock, J. A., and Amundson, R.: Dissolved organic carbon chemistry and dynamics in contrasting forest and grassland soils, *Biogeochemistry*, 89, 181–198, <https://doi.org/10.1007/s10533-008-9211-x>, 2008.
- Scheibe, A., Sierra, C. A., and Spohn, M.: Recently fixed carbon fuels microbial activity several meters below the soil surface, Zenodo [data set], <https://doi.org/10.5281/zenodo.7389812>, 2022.
- Schmidt, M. W., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. A., Kleber, M., Kögel-Knabner, I., Lehmann, J., Manning, D. A. C., Nannipieri, P., Rasse, D. P., Weiner, S., and Trumbore, S. E.: Persistence of soil organic matter as an ecosystem property, *Nature*, 478, 49–56, <https://doi.org/10.1038/nature10386>, 2011.
- Seuss, I., Scheibe, A., and Spohn, M.: N₂ fixation is less sensitive to changes in soil water content than carbon and net nitrogen mineralization, *Geoderma*, 424, 115973, <https://doi.org/10.1016/j.geoderma.2022.115973>, 2022.
- Spohn, M. and Holzheu, S.: Temperature controls diel oscillation of the CO₂ concentration in a desert soil, *Biogeochemistry*, 156, 279–292, <https://doi.org/10.1007/s10533-021-00845-0>, 2021.
- Steinhof, A., Adamec, G., Gleixner, G., van Klinken, G. J., and Wagner, T.: The new ¹⁴C analysis laboratory in Jena, Germany, *Radiocarbon*, 46, 51–58, <https://doi.org/10.1017/S0033822200039345>, 2004.
- Steinhof, A., Altenburg, M., and Machts, H.: Sample preparation at the Jena ¹⁴C laboratory, *Radiocarbon*, 59, 815–830, <https://doi.org/10.1017/RDC.2017.50>, 2017.
- Stuiver, M. and Polach, H. A.: Radiocarbon – Discussion reporting of ¹⁴C data, *Radiocarbon*, 19, 355–363, <https://doi.org/10.1017/S0033822200003672>, 1977.
- Trumbore, S.: Age of soil organic matter and soil respiration: radiocarbon constraints on belowground C dynamics, *Ecol. Appl.*, 10, 399–411, [https://doi.org/10.1890/1051-0761\(2000\)010\[0399:AOSOMA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0399:AOSOMA]2.0.CO;2), 2000.
- Trumbore, S. E., Davidson, E. A., Barbosa de Camargo, P., Nepstad, D. C., and Martinelli, L. A.: Belowground cycling of carbon in forests and pastures of eastern Amazonia, *Global Biogeochem. Cy.*, 9, 515–528, <https://doi.org/10.1029/95GB02148>, 1995.
- Trumbore, S. E., Sierra, C. A., and Hicks Pries, C. E.: Radiocarbon nomenclature, theory, models, and interpretation: Measuring age, determining cycling rates, and tracing source pools, in: *Radiocarbon and Climate Change*, edited by: Schuur, E. A. G., Druffel, E., and Trumbore, S. E., Springer International Publishing, 45–82, https://doi.org/10.1007/978-3-319-25643-6_3, 2016.
- Uroz, S., Picard, L., and Turpault, M. P.: Recent progress in understanding the ecology and molecular genetics of soil mineral weathering bacteria, *Trends Microbiol.*, 30, 882–897, <https://doi.org/10.1016/j.tim.2022.01.019>, 2022.
- van Hees, P. A., Jones, D. L., Finlay, R., Godbold, D. L., and Lundström, U. S.: The carbon we do not see – the impact of low molecular weight compounds on carbon dynamics and respiration in forest soils: a review, *Soil Biol. Biochem.*, 37, 1–13, <https://doi.org/10.1016/j.soilbio.2004.06.010>, 2005.
- Vázquez, M., Ramírez, S., Morata, D., Reich, M., Braun, J. J., and Carretier, S.: Regolith production and chemical weathering of granitic rocks in central Chile, *Chem. Geol.*, 446, 87–98, <https://doi.org/10.1016/j.chemgeo.2016.09.023>, 2016.
- Werner, C., Schmid, M., Ehlers, T. A., Fuentes-Espoz, J. P., Steinkamp, J., Forrest, M., Liakka, J., Maldonado, A., and Hickler, T.: Effect of changing vegetation and precipitation on denudation – Part 1: Predicted vegetation composition and cover over the last 21 thousand years along the Coastal Cordillera of Chile, *Earth Surf. Dynam.*, 6, 829–858, <https://doi.org/10.5194/esurf-6-829-2018>, 2018.
- Wynn, J. G., Harden, J. W., and Fries, T. L.: Stable carbon isotope depth profiles and soil organic carbon dynamics in the lower Mississippi Basin, *Geoderma*, 131, 89–109, <https://doi.org/10.1016/j.geoderma.2005.03.005>, 2006.