

Sveriges lantbruksuniversitet Swedish University of Agricultural Sciences

SLU Risk Assessment of Plant Pests

SLU.ua.2022.2.6-1487 April 19, 2022

# Feedback on a list of plant pests with candidates for risk assessments – Batch 3

## Background

Three plant pests were recently found in EFSAs media and literature horizon scanning and evaluated using EFSAs PeMoScoring tool (EFSA 2021). The pests were scored as either positive (above a certain threshold value) or negative (below the threshold).

The three species were:

- *Neofusicoccum mediterraneum* (positive)
- Citrus concave gum-associated virus (negative)
- Chinese wheat mosaic virus (negative)

SLU Risk Assessment of Plant Pests was requested by the Swedish Board of Agriculture to provide feedback in terms of (i) whether any of these pests are present in Sweden and (ii) whether there are some special reasons to exclude or prioritize any of the pests for further pest categorizations (i.e., in addition to those provided by an EFSA PeMoScoring evaluation of these species (EFSA 2022, unpublished)).

## Methods

The scientific names, including synonyms (see below), were matched against the Swedish Taxonomic Database that provides information about which species that are established in Sweden (Artfakta 2022), and against the databases iNaturalist (2022) and GBIF (2022). Synonyms were obtained from EPPO Global Database (EPPO 2022). Location records for the fungal species was also searched in USDA Fungal databases (Farr & Rossman 2021). A search was conducted for all the scientific names and filtering for "Sweden" in Web of Science (2021). Further, such searches were also conducted using the search engine Google, but restricting the searches to Swedish webpages. Finally, searches were conducted with Google Scholar combining the scientific names with the word "Sweden". Information about the pests was also requested from a Swedish expert on plant viruses (see Acknowledgement).

### **Results and discussion**

The search procedure described in the Methods section above did not reveal any information indicating that any of the three species has been found in Sweden. Some noteworthy information about the pests are presented below.

#### Neofusicoccum mediterraneum

*Neofusicoccum mediterraneum* is a fungal pathogen described as a new species in 2007 (Crous et al. 2007) and no synonyms were found (Farr and Rossman 2022). The fungus was first discovered in Greece and Italy (Crous et al. 2007). The species has since been reported from numerous different hosts and from many locations in EU, i.e. in Italy and Spain<sup>1</sup>. The fungus is also reported from Algeria, Turkey, Tunisia, India, Iran, Sri Lanka, South Africa and USA (California) (Farr and Rossman 2022; Yildiz et al. 2022; Abouzkhar et al. 2022). Given that the fungus has been present in the Mediterranean area for more than 15 years and that it has been reported from several locations, hosts and production systems suggest that it may be widespread in this region.

Some of its hosts that may be especially relevant for Swedish conditions are *Fraxinus* sp., *Juniperus* sp. and *Rubus* sp. (Inderbitzin et al. 2010; Farr and Rossman 2022).

#### Citrus concave gum-associated virus

The preferred name of the Citrus concave gum-associated virus according to EPPO (2022) is Citrus coguvirus [EPPO code: CCGAV0] and the abbreviation CCGaV is also listed as another scientific name. It may be noted that coguviruses have recently been discovered and that little is known about them, e.g., how they spread and to which degree they cause disease (A. Kvarnheden, pers. comm. 2022). For the Citrus coguvirus there is a possibility of entry in Sweden with infected propagation material from e.g., areas in Europe where it occurs in apple trees (A. Kvarnheden, pers. comm. 2022).

#### Chinese wheat mosaic virus

The Chinese wheat mosaic virus [EPPO code: WCHMV0] has the following synonyms; CWMV, WChMV, Wheat China mosaic furovirus and Wheat China mosaic virus (EPPO 2022). The virus was added to EPPO Alert List in 1999 but deleted in 2001 since there was no indication that it was a threatening disease (EPPO 1999; EPPO 2001). It is transmitted by a

<sup>&</sup>lt;sup>1</sup> Examples of reports from Italy and Spain. Italy: Puglia - *Olea europaea* (Crous et al. 2007; Lazzizera et al. 2008), Sicily, orchard - *Juglans regia* (Gusella et al. 2020), Sicily, urban area - *Ficus microcarpa* (Fiorenza et al. 2022), Sicily, orchards (4 farms) - *Pistacia vera* (Gusella et al. 2021), Sardinia, plantations (5 sites) - *Eucalyptus camaldulensis* (Deidda et al. 2016). Spain: Andalucia (2 locations) - *O. europaea* and *P. vera* (Moral et al. 2010), Valencia, commercial packinghouses - *Diospyros kaki* (Palou et al. 2013), Castilla y León - *Vitis vinifera* (Martin et al. 2011), Sevilla - *Vitis* sp. (Varela et al. 2011), southern Spain, orchards - *J. regia* (López-Moral et al. 2020), Málaga, orchards - *Persea americana* (Arjona-Girona et al. 2019), Valencia, orchards - *Eriobotrya japonica* (González-Domínguez et al. 2016), Mallorca Island, orchards (2 locations) - *Prunus dulcis* (Olmo et al. 2016).

vector, i.e., *Polymyxa graminis*, which is present in Sweden (Lindsten & Tapio 1986; Eckersten et al. 2015).

It may be noted that viruses that are spread with *Polymyxa* are difficult (or impossible) to eradicate if they become introduced, e.g. due to that resting spores of *Polymyxa* can remain in the soil with infectious virus for several years (A. Kvarnheden, pers. comm. 2022). It may also be noted that there are other furoviruses, that cause mosaic disease in wheat and that are vectored by *P. graminis*, which are closely related to CWMV and which occurs in Europe and there appears to be a debate whether these should be regarded as conspecific (Hairi and Meyer 2007; Tamada and Kondo 2013). None of these are however currently found in Sweden (A. Kvarnheden, pers. comm. 2022).

## Conclusion

No evidence was found that any of the three pests have been found in Sweden.

Some noteworthy information was found for the pests but no additional reasons to exclude or prioritize these pests for further pest categorizations beyond those provided by the EFSA PeMoScoring evaluations (EFSA 2022, unpublished).

## Acknowledgement

We would like to thank Professor Anders Kvarnheden at the Department of Plant Biology, SLU for providing information about the viruses.

## Authors

This report was prepared by SLU Risk Assessment of Plant Pests at the Swedish University of Agricultural Sciences:

Niklas Björklund, Dept. of Ecology, Swedish University of Agricultural Sciences, P.O. Box 7044, SE-750 07 Uppsala, Sweden. Visiting address: Ullsväg 16, E-mail: Niklas.Bjorklund@slu.se

Johanna Boberg, Dept. of Forest Mycology and Plant Pathology, Swedish University of Agricultural Sciences, PO Box 7026, SE-750 07 Uppsala, Sweden. Visiting address: Almas allé 5, E-mail: Johanna.Boberg@slu.se

## References

Arjona-Girona, I., Ruano-Rosa, D., & López-Herrera, C. J. (2019). Identification, pathogenicity and distribution of the causal agents of dieback in avocado orchards in Spain. Spanish journal of agricultural research, 17(1), e1003-e1003. <u>https://doi.org/10.5424/sjar/2019171-13561</u>

Crous, P.W., Groenewald, J.Z., Wingfield, M.J., and Phillips, A.J.L. (2007) *Neofusicoccum mediterraneum* Crous, M.J. Wingf. & A.J.L. Phillips, sp. nov. Fungal Planet: 19.

Deidda, A., Buffa, F., Linaldeddu, B. T., Pinna, C., Scanu, B., Deiana, V., ... & Floris, I. (2016). Emerging pests and diseases threaten *Eucalyptus camaldulensis* plantations in Sardinia, Italy. iForest-Biogeosciences and Forestry, 9(6), 883. <u>https://doi.org/10.3832/ifor1805-009</u>

Eckersten, H., Djurle, A., Albihn, A., Båge, R., Andersson, L., De Toro, A., ... & Magnusson, U. (2015). Framtida risker och hot mot svensk spannmåls-respektive mjölkproduktion. Utgivare: SLU, Framtidens lantbruk - djur växter och markanvändning, ISBN: 978-91-576-9334-1, LINK [Accessed 12 April 2022]

EPPO (1999) EPPO Alert List. EPPO Reporting Service no. 08-1999. Num. article: 1999/134 https://gd.eppo.int/reporting/article-3454

EPPO (2001) Mini data sheet on Wheat China mosaic furovirus. https://gd.eppo.int/download/doc/1043\_minids\_WCHMV0.pdf

EPPO (2022) EPPO Global Database (available online). https://gd.eppo.int

Farr, D.F., & Rossman, A.Y. (2022) Fungal Databases, U.S. National Fungus Collections, ARS, USDA. Retrieved April 6, 2022, from <u>https://nt.ars-grin.gov/fungaldatabases/</u>

Fiorenza, A., Aiello, D., Costanzo, M. B., Gusella, G., & Polizzi, G. (2022). A new disease for Europe of *Ficus microcarpa* caused by Botryosphaeriaceae Species. Plants, 11(6), 727. https://www.mdpi.com/2223-7747/11/6/727

González-Domínguez, E., Alves, A., León, M., & Armengol, J. (2017). Characterization of Botryosphaeriaceae species associated with diseased loquat (*Eriobotrya japonica*) in Spain. Plant Pathology, 66(1), 77-89. <u>https://doi.org/10.1111/ppa.12556</u>

Gusella, G., Giambra, S., Conigliaro, G., Burruano, S., & Polizzi, G. (2021). Botryosphaeriaceae species causing canker and dieback of English walnut (Juglans regia) in Italy. Forest Pathology, 51(1), e12661. <u>https://doi.org/10.1111/efp.12661</u>

Gusella, G., Lawrence, D. P., Aiello, D., Luo, Y., Polizzi, G., & Michailides, T. (2021). Etiology of Botryosphaeria Panicle and Shoot Blight of Pistachio (*Pistacia vera*) caused by Botryosphaeriaceae in Italy. Plant Disease, 106:4, 1192-1202. <u>https://doi.org/10.1094/PDIS-08-21-1672-RE</u>

Hariri, D., & Meyer, M. (2007). A new furovirus infecting barley in France closely related to the Japanese soil-borne wheat mosaic virus. *European journal of plant pathology*, *118*(1), 1-10. https://doi.org/10.1007/s10658-006-9095-3 Inderbitzin, P., Bostock, R.M., Trouillas, F.P. & Michailides, T.J. (2010) A six locus phylogeny reveals high species diversity in Botryosphaeriaceae from California almond, Mycologia, 102:6, 1350-1368, <u>https://doi.org/10.3852/10-006</u>

Lazzizera, C., Frisullo, S., Alves, A., & Phillips, A. J. L. (2008). Morphology, phylogeny and pathogenicity of Botryosphaeria and *Neofusicoccum* species associated with drupe rot of olives in southern Italy. Plant Pathology, 57(5), 948-956. <u>https://doi.org/10.1111/j.1365-3059.2008.01842.x</u>

Lindsten, K & Tapio, E. (1986) Jordburna växtviroser – en förbisedd grupp av sjukdomar? Växtskyddsnotiser, nr 4-5, p. 83-84 <u>https://www.slu.se/globalassets/ew/org/inst/ekol/vaxtskyddsnotiser/vaxtskyddsnotiser-arkiv/1986\_50\_4-5.pdf</u> [Accessed 12 April 2022]

López-Moral, A., Lovera, M., Raya, M. D. C., Cortés-Cosano, N., Arquero, O., Trapero, A., & Agustí-Brisach, C. (2020). Etiology of branch dieback and shoot blight of English walnut caused by Botryosphaeriaceae and Diaporthe Species in Southern Spain. Plant Disease, 104(2), 533-550. <u>https://doi.org/10.1094/PDIS-03-19-0545-RE</u>

Martin, M. T., Martin, L., & Cuesta, M. J. (2011). First report of *Neofusicoccum mediterraneum* and N. australe causing decay in Vitis vinifera in Castilla y León, Spain. Plant Disease, 95(7), 876-876. <u>https://doi.org/10.1094/PDIS-03-11-0172</u>

Moral, J., Muñoz-Díez, C., González, N., Trapero, A., & Michailides, T. J. (2010). Characterization and pathogenicity of Botryosphaeriaceae species collected from olive and other hosts in Spain and California. Phytopathology, 100(12), 1340-1351.

Olmo, D., Armengol, J., León, M., & Gramaje, D. (2016). Characterization and pathogenicity of Botryosphaeriaceae species isolated from almond trees on the island of Mallorca (Spain). Plant Disease, 100(12), 2483-2491. <u>https://doi.org/10.1094/PDIS-05-16-0676-RE</u>

Palou, L., Montesinos-Herrero, C., Besada, C., & Taberner, V. (2013). Postharvest fruit rot of persimmon (*Diospyros kaki*) in Spain caused by *Lasiodiplodia theobromae* and *Neofusicoccum* spp. Journal of Phytopathology, 161(9), 625-631. <u>https://doi.org/10.1111/jph.12111</u>

Tamada, T., & Kondo, H. (2013). Biological and genetic diversity of plasmodiophoridtransmitted viruses and their vectors. Journal of general plant pathology, 79(5), 307-320. https://doi.org/10.1007/s10327-013-0457-3

Varela, C. P., Fernández, V. R., Casal, O. A., & Vázquez, J. M. (2011). First report of cankers and dieback caused by *Neofusicoccum mediterraneum* and *Diplodia corticola* on grapevine in Spain. Plant Disease, 95(10), 1315-1315. <u>https://doi.org/10.1094/PDIS-05-11-0429</u>