



# Assessing freshwater fish biodiversity of Kumbe River, Papua (Indonesia) through environmental DNA metabarcoding

Arif Wibowo<sup>A,\*</sup> , Kurniawan Kurniawan<sup>A</sup>, Dwi Atminarso<sup>A,B</sup>, Tri Heru Prihadi<sup>C</sup>, Lee J. Baumgartner<sup>B</sup>, Meaghan L. Rourke<sup>B,D</sup>, Satoshi Nagai<sup>E</sup>, Nicolas Hubert<sup>F</sup> and Anti Vasemagi<sup>G,H</sup>

For full list of author affiliations and declarations see end of paper

**\*Correspondence to:**

Arif Wibowo  
Research Center for Conservation of Marine and Inland Water Resources, National Research and Innovation Agency, Jalan Raya Jakarta-Bogor Km. 48 Cibinong, Bogor, West Java 16911, Indonesia  
Email: [wibowo@daad-alumni.de](mailto:wibowo@daad-alumni.de)

**Handling Editor:**

Alan Lymbery

**Received:** 25 December 2021

**Accepted:** 29 July 2022

**Published:** 25 August 2022

**Cite this:**

Wibowo A *et al.* (2023)  
*Pacific Conservation Biology*, **29**(4), 340–350.  
doi: [10.1071/PC21078](https://doi.org/10.1071/PC21078)

© 2023 The Author(s) (or their employer(s)). Published by CSIRO Publishing.  
This is an open access article distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND).

OPEN ACCESS

## ABSTRACT

**Context.** The ability to accurately assess biodiversity is a critical first step towards effective conservation and management. However, assessment of biodiversity using conventional monitoring programs is often constrained by high cost and a lack of taxonomic expertise. Environmental DNA (eDNA) metabarcoding may be a useful tool to efficiently catalogue biodiversity in areas that cannot be easily assessed using other methods. **Aims.** Here, we evaluated the potential of eDNA metabarcoding for assessing fish biodiversity and distribution in the Kumbe River, Papua Province, Indonesia. **Methods.** We selected four sampling locations and collected seven eDNA samples from each location. We used eDNA metabarcoding of the Cytochrome-*b* gene to characterise the fish community. **Key results.** A total of 23 species were detected, three of which comprised 92% of sequence reads detected: *Melanotaenia goldiei* (32%), *Craterocephalus randi* (31%), and the invasive tilapia *Oreochromis niloticus* (29%). Only five species that were previously detected using conventional methods were detected by metabarcoding: *M. goldiei*, *Craterocephalus stercusmuscarum*, *O. niloticus*, *Neoarius graeffei*, and *Arius arius*. We detected 18 species (70% native) that have never been recorded from the Kumbe River. **Conclusions.** This work has demonstrated that fish biodiversity is substantially underestimated in the Kumbe River. Environmental DNA metabarcoding is a promising rapid, non-invasive and cost-effective method for assessing fish biodiversity in Papua. **Implications.** The findings support future investment in eDNA metabarcoding to characterise the fish biodiversity in Papua. This will assist in allocating the limited resources for conservation and management to areas most at risk from anthropogenic impacts.

**Keywords:** biodiversity, eDNA, freshwater fish, Indonesia, metabarcoding, Papua, river, tropical.

## Introduction

Defined by Walter Rosen in 1986 during the first American forum on biological diversity, biodiversity represents the variety and variability among living organisms and ecosystems where they occur (Pereira *et al.* 2013; Bartkowski *et al.* 2015). The challenge for biodiversity conservation is to understand when and how major perturbations such as climate change cause significant impact (Yachi and Loreau 1999; Mace *et al.* 2012). Thus, it is the responsibility of individual countries to preserve biodiversity for its environmental, productive, consumptive, social, ethical and aesthetic value (Pauchard 2017).

Indonesia is one of the most biodiverse countries, and its freshwater fish biodiversity is no exception. A total of 1200 native fish species have been reported across the archipelago (Kottelat and Whitten 1996; Hubert *et al.* 2015). The inventory of Indonesian freshwater species is ongoing, and recent studies have suggested that fish diversity may be underestimated (Kadariusman *et al.* 2012; Hubert *et al.* 2019; Sholihah *et al.* 2020, 2021a, 2021b). Thus, the incomplete understanding of fish biodiversity presents challenges for effective conservation and management. While traditional methods can be implemented to document biodiversity, these can be cost-prohibitive and time consuming. In

addition, it also requires detailed expertise in fish taxonomy, which becomes limiting as the number of taxonomists declines. Optimising monitoring strategies will be critical to determine change in ecosystems; and monitoring over time is a crucial step toward sustainable use of resources (Xiao *et al.* 2016; Trebitz *et al.* 2017). This challenge is particularly pronounced in Indonesia, where aquatic biodiversity-related research output is not balanced with biodiversity complexity (Wibowo *et al.* 2018; Gustiano *et al.* 2021; Kurniawan *et al.* 2021) and understanding of the dynamics behind the origin and maintenance of freshwater biodiversity is still fragmentary (de Bruyn *et al.* 2013, 2014; Sholihah *et al.* 2021a, 2021b).

Threats to Indonesian aquatic biodiversity have escalated in recent decades through agricultural development (Cleary and DeVantier 2011; Imai *et al.* 2018; Austin *et al.* 2019), introduction of exotic species (Herder *et al.* 2012; Dahruddin *et al.* 2017), and pollution (Garg *et al.* 2018). As such, locally decreasing ichthyodiversity is of concern, and taxonomic knowledge gaps are challenging (Dahruddin *et al.* 2017; Hubert *et al.* 2019). Papua, the western most Indonesian province of the Island of New Guinea, contains 16% of known Indonesian freshwater fish diversity and 20% of endemic species, including all fish species within the genus *Melanotaenia* (Rainbowfish) (Hubert *et al.* 2015). New species or expanded distributions of known species are still being reported (Nugraha *et al.* 2015; Wibowo *et al.* 2017; Ditya *et al.* 2018). However, much of the fish biodiversity of Papua is still undiscovered (Kadariusman *et al.* 2012; Koh *et al.* 2013; Hubert *et al.* 2015).

An alternative approach to identify fish biodiversity is to collect DNA shed by the target species into the water. All aquatic organisms release DNA into the environment via mucus, urine, faeces, or dead tissue (Goldberg *et al.* 2016; Carraro *et al.* 2018), where it is then known as environmental DNA (eDNA). Water samples containing eDNA from the entire aquatic community (e.g. bacteria, algae, fish) can be filtered to extract eDNA and the target species can be identified using the relevant primer sets (Turner *et al.* 2015; Pont *et al.* 2018). Environmental DNA sampling can be used to detect a single species or multiple species (known as metabarcoding) (Taberlet *et al.* 2012; Rees *et al.* 2014; Pont *et al.* 2018). Although species-specific eDNA sampling may be more sensitive than metabarcoding – especially for rare species (Bylemans *et al.* 2019) – eDNA metabarcoding is a rapid, cost-effective, and non-invasive technique (Janosik and Johnston 2015; Seymour 2019; Sigsgaard *et al.* 2020). Environmental DNA has been successfully implemented in the surveillance of rare, invasive, or migratory fish species in various aquatic systems (Nathan *et al.* 2015; Sigsgaard *et al.* 2015; Simpfendorfer *et al.* 2016; Rice *et al.* 2018; Itakura *et al.* 2019; McElroy *et al.* 2020). However, eDNA metabarcoding has rarely been applied in Indonesia, with only three studies in freshwater environments. For instance, eDNA was successfully implemented in detection of

invasive alligator gar *Atractosteus spatula* (Ulayya *et al.* 2020) and crayfish *Cherax quadricarinatus* (Djalil *et al.* 2018) in lakes of West Java. Assessing fish biodiversity using eDNA was carried out in the Maninjau lake of Sumatra, where 26 species were detected in the lake, including five native and 21 exotic fish species (Roesma *et al.* 2021).

The objective of this study was to evaluate the utility of eDNA metabarcoding to rapidly characterise the fish community of the Kumbe River in West Papua and assess its potential in a tropical context as a suitable alternative method to traditional sampling to characterise fish diversity.


## Materials and methods

### Sample collection and preservation

Sampling was undertaken at four locations in the Kumbe River, which is located in the southern part of Papua Island (Fig. 1). The Kumbe River is a meandering tropical river with a total length of 242 km and width ranging from 97 to 700 m (Anonymous 2010). The average annual rainfall of this area is 2160 mm and average annual temperature of 23°C (Peel *et al.* 2007). We collected seven 15 mL water samples (a total of 28 samples) from each of four sampling locations (Fig. 1). Immediately after collection, we added 1.5 mL of 3 M sodium acetate and 33 mL of absolute ethanol, mixed with the water sample and then stored it at room temperature in the field. Once in the laboratory, water samples were placed at –20°C for DNA preservation until DNA extraction.

### DNA extraction and PCR amplification

Water samples were centrifuged for 35 min at 5500g at 24°C to precipitate DNA and suspended material, and supernatant was discarded (Valiere and Taberlet 2000). Precipitates were resuspended in 100 µL ultrapure water and DNA was extracted using the QIAamp Tissue Extraction Kit (Qiagen). We extracted DNA from each precipitate individually and DNA extraction was performed in a DNA laminar flow hood to avoid contamination. A negative control containing all reagents but no eDNA was included in each batch of extractions to monitor contamination. The concentration of extracted DNA ranged from 1 to 20 ng/µL. Polymerase chain reaction (PCR) amplification was performed using the primers 5'-TGCCAACGGAGCATCATTC-3' and 5'-ATAAAGGTAGGAGCCGTAGT-3', which amplify a 79 base pairs (bp) segment of the mitochondrial Cytochrome b (Cyt-b) (Ficetola *et al.* 2008). Each PCR reaction contained 2.5 µL of 1 × KAPA HiFi Hotstart ReadyMix (Kapa Biosystems), 0.5 µL of each primer (1 mM) and 5 µL of DNA extract; 4.0 µL PCR grade water was added up to a reaction volume of 12.5 µL. A negative control was included with each PCR batch to monitor the



**Fig. 1.** (a) Map of the study location in the Kumbe River, Papua Island, Indonesia and photos of each sampling site, (b) Alfasera: S'07°24'23.9"; E'140°37'04.6", (c) Inggun: S'07°59'05.3"; E'140°27'53.3", (d) Yakui: S'08°02'26.3"; E'140°31'47.5", and (e) Sakor: S'08°06'51.6"; E'140°29'98.9".

contamination of the reagents used. The PCR conditions were as follows: one cycle at 95°C for 5 min; 40 cycles at 95°C for 30 s, 50°C for 30 s, 72°C for 30 s; and one cycle at 72°C for 5 min. PCR products were visualised under UV light using a 2% agarose gel stained with Midori Green Advanced (Nippon Genetics Europe). PCR products were purified using the NucleoMag 96 PCR Kit (Macherey-Nagel).

### Ion Torrent PGM preparation and reads

We pooled individually tagged DNA samples into two different sequencing libraries. Libraries were set up as described by Meyer and Kircher (2010) but modified as follows for Ion Torrent Personal Genome Machine (PGM) sequencing. The adapter mix was prepared using an adapter marked with a multiplex rating (MID). Adapters are designed to include individual MID's and to match Ion Torrent-specific priming sites. The total molar concentration of the adapter mixture was 20 M for each adapter. We used a 100 bp long PCR product as the positive control DNA template. We carried out blunt-end reaction with half of the volume, i.e. we added 10 µL of blunt-end master mix to 25 µL of the sample.

The reaction was purified using solid phase reversible immobilisation (SPRI) beads with a template: bead ratio of

1:1.8. In the post-ligation purification steps, the template: bead ratio was 1:1. In order to verify adapter ligation success, the SPRI-purified MID-tagged positive control library and a subset of MID-tagged library aliquots were separated side by side with non-MID-tagged templates on 2% agarose gel stained with Midori Green Advanced electrophoresis (45 min, 95 V) and visualised using UV light. We measured DNA concentration of the samples using a Qubit Fluorometer (Life Technologies) and mixed amplicon libraries subsequently in equimolar ratios. The stock library was re-amplified in two different reactions as follows: 5 µL pooled library stock was added to a master mix comprising of 5 U of Herculase II polymerase (Agilent technologies, catalogue number 600677), 1 × Herculase II reaction buffer, 25 mM each deoxynucleoside triphosphate, 10 µM each primer, and added PCR grade water up to 50 µL. We designed the re-amplification primers based on the sequence of the IonTorrent-specific priming sites and thus generated millions of MID-tagged copies including binding sites necessary for subsequent sequencing with IonTorrent technology. The thermocycling profile included a 30 s denaturation at 98°C followed by 15 cycles consisting of a 20 s denaturation at 98°C, a 30 s annealing at 64°C, and a 30 s elongation at 72°C. Final elongation was conducted at 72°C for 5 min. To clean the re-amplified

library, size-selection was conducted by separating the entire library using 2% Size-Select Agarose E- Gel and E-Gel Electrophoresis System (Life Technologies) according to the manufacturer's instructions. The DNA concentration of amplified library pool was measured using a Qubit Fluorometer. The library pool stock was then diluted to a final concentration of 26 pmol. For template preparation, an 18  $\mu$ L aliquot of the library dilution (approximately  $2.8 \times 10^8$  molecules) was transferred into the sequencing reaction set-up. Emulsion PCR and Ion Torrent PGM sequencing were carried out on two 314 chips (Life Sciences, catalogue number 4462923) according to the manufacturer's protocol (Publication Part Number: 4471974 Rev. C).

The resulting reads were binned and renamed by MID, i.e. the original individual samples using the software Geneious Pro 6.1. Then the original-specific primers (Ficetola *et al.* 2008) were trimmed off, reads were trimmed for poor quality parts using a 0.05 error probability limit and reads shorter than 100 bp were excluded using Geneious Pro 6.1. Thus, only reads comprising both the Ion Torrent adapter with a MID and the original primer were passed on for further analysis. Subsequent analyses were performed employing supercomputers at the IT Centre for fisheries (Yokohama, Japan). Sequences were assigned to species using the BLASTN 2.2.25+ algorithm against the reference database (GenBank). We examined Basic Local Alignment Search Tool (BLAST) results showing a sequence similarity of at least 94% and we also considered species distribution when confirming species identity (Wibowo *et al.* 2017).

### The analysis of BLAST result

The BLAST output was imported into the program MEGAN (Huson *et al.* 2007) to further summarise the results. We employed default parameters, except for following settings: Min support = 1, Min score = 100, Min complexity = 0. We utilised a built-in comparison tool in MEGAN for database comparison. We discarded all hits below 94% identity. We also discarded hits with e-value over  $1e-20$ . We further analysed the hits one by one to see if there were reads matching more than one species in GenBank. In case of multiple hits, we (a) identified the hit with the longest alignment length, (b) identified the hit with a match to an adult specimen over a juvenile specimen (given the greater confidence in morphological identification of an adult specimen), and (c) reviewed the database entry for errors and omitted incorrect entries.

## Results

A total of 629 976 raw reads were detected by sequencing of amplicons from 28 water samples and filtered to 297 438 reads. Thus, altogether 332 538 reads (52.79%) were discarded from further analysis after quality control. The

numbers of filtered sequences obtained from Alfasea, Yakui, Sakor, and Inggun were 71 627, 89 014, 81 030 and 55 767 reads, respectively. BLAST queries resulted in sequence similarities between reads and reference sequences in Genbank ranging between 94 and 98% (Table 1). A total of 23 freshwater fish species were detected (Table 1).

Species diversity was greatest at Inggun (21 species) and Yakui (21 species), followed by Sakor (19 species) and Alfasea (16 species) (Fig. 2a–d). The number of reads per species varied considerably among the seven sampling replicates within sites, with reads proportion ranging from 0.07 to 0.27 for *Oreochromis niloticus*, at Alfasea for instance (Fig. 2a). Some rarer species were not represented in all sampling replicates within sites including *Trichopodus microlepis*, *Arius arius* and *Smilosicyopus bitaeniatus*, which were not represented in replicates one and five at Yakui, and *Melanotaenia goldiei*, which was not represented in replicate six at Inggun.

Fourteen of the 23 species were detected at all sites (*M. goldiei*, *O. niloticus*, *Craterocephalus randi*, *Craterocephalus stercusmuscarum*, *Craterocephalus nouhuysi*, *Craterocephalus gloveri*, *S. bitaeniatus*, *Neogobius melanostomus*, *Oryzias haugiensis*, *Atherinella schultzi*, *Acanthopagrus berda*, *Diplospinus multistriatus*, *Anguilla australis* and *Labeobarbus natalensis*), five species were detected at three sites (*Melanotaenia albimarginata*, *M. sp.*, *Neoarius graeffei*, *Neoarius berneyi* and *Oryzias latipes*), two species were detected at in two sites (*Oxyeleotris lineolata* and *Pseudanthias dispar*) and two species were detected at one site (*A. arius* and *T. microlepis*) (Table 1, Fig. 3). A total of 16 species were native (*M. goldiei*, *M. albimarginata*, *M. sp.*, *C. randi*, *C. stercusmuscarum*, *C. nouhuysi*, *C. gloveri*, *N. melanostomus*, *N. berneyi*, *N. graeffei*, *A. berda*, *A. australis*, *D. multistriatus*, *O. lineolata*, *P. dispar* and *A. arius*) and seven species were exotic (*O. niloticus*, *S. bitaeniatus*, *O. haugiensis*, *O. latipes*, *A. schultzi*, *L. natalensis* and *T. microlepis*) (Table 1). If reads were ranked by abundance (Table 1), three species dominated all sites, consisting of *M. goldiei* (32.03% of the all reads), *C. randi* (31.19% of all reads), and *O. niloticus* (28.77% of reads). The 23 detected species belonged to 13 families and 16 genera. Atherinidae (35% of all sites; four species) was the most abundant family in terms of sequencing reads, followed by Melanotaeniidae (32% of all sites; three species) and Cichlidae (29% of all sites; two species). Sequences from these three families dominated all studied locations (Fig. 4a). These families mostly represented the genera *Craterocephalus*, *Melanotaenia* and *Oreochromis* (Fig. 4b).

A total of 23 species have previously been detected in the Kumbe River using conventional methods and DNA barcoding (Table 1). We detected five of these species (*N. graeffei*, *M. goldie*, *C. stercusmuscarum*, *O. niloticus* and *A. arius*) using metabarcoding, and an additional 18 species (69.56% native species) not previously recorded from the Kumbe River.

**Table 1.** The list of fish species detected at Kumbé River collected based on eDNA and previous conventional sampling.

Family	Species	Identification method	Similarity	Reads	Abundance (%)	Reference	Status
Adrianichthyidae	<i>Oryzias haugiangensis</i>	eDNA/BLAST	96%	1330	0.45	Present study	Exotic
Adrianichthyidae	<i>Oryzias latipes</i>	eDNA/BLAST	96%	7	0.00	Present study	Exotic
Ambassidae	<i>Parambassis gulliver</i>	Morphology identification	–	–	–	Ditya et al. (2018)	Native
Anabantidae	<i>Anabas testudineus</i>	Morphology identification	–	–	–	Ditya et al. (2018)	Native
Anguillidae	<i>Anguilla australis</i>	eDNA/BLAST	95%	33	0.01	Present study	Native
Apogonidae	<i>Glossamia aprion</i>	DNA barcoding	–	–	–	Wibowo et al. (2017)	Native
Ariidae	<i>Arius arius</i>	Morphology identification	–	–	–	Ditya et al. (2018)	Native
Ariidae	<i>Arius arius</i>	eDNA/BLAST	94%	1	0.00	Present study	Native
Ariidae	<i>Neoarius aff. graeffei</i>	eDNA/BLAST	95%	5	0.00	Present study	Native
Ariidae	<i>Neoarius berneyi</i>	eDNA/BLAST	98%	28	0.01	Present study	Native
Ariidae	<i>Neoarius graeffei</i>	Morphology identification and DNA barcoding	–	–	–	Wibowo et al. (2017); Ditya et al. (2018)	Native
Atherinidae	<i>Craterocephalus gloveri</i>	eDNA/BLAST	96%	37	0.01	Present study	Native
Atherinidae	<i>Craterocephalus nouhuysi</i>	eDNA/BLAST	98%	5131	1.73	Present study	Native
Atherinidae	<i>Craterocephalus randi</i>	eDNA/BLAST	97%	92 774	31.19	Present study	Native
Atherinidae	<i>Craterocephalus stercusmuscarum</i>	DNA barcoding	–	–	–	Wibowo et al. (2017)	Native
Atherinidae	<i>Craterocephalus stercusmuscarum</i>	eDNA/BLAST	96%	7998	2.69	Present study	Native
Atherinopsidae	<i>Atherinella schultzi</i>	eDNA/BLAST	95%	1061	0.36	Present study	Exotic
Belontiidae	<i>Strongylura krefftii</i>	DNA barcoding	–	–	–	Wibowo et al. (2017)	Native
Channidae	<i>Channa striata</i>	Morphology identification	–	–	–	Ditya et al. (2018)	Exotic
Cichlidae	<i>Acanthopagrus berda</i>	eDNA/BLAST	96%	214	0.07	Present study	Native
Cichlidae	<i>Oreochromis mossambicus</i>	Morphology identification	–	–	–	Ditya et al. (2018)	Exotic
Cichlidae	<i>Oreochromis niloticus</i>	Morphology identification	–	–	–	Ditya et al. (2018)	Exotic
Cichlidae	<i>Oreochromis niloticus</i>	eDNA/BLAST	97%	85 582	28.77	Present study	Exotic
Clariidae	<i>Clarias batracus</i>	Morphology identification	–	–	–	Ditya et al. (2018)	Exotic
Clupeidae	<i>Clupeoides venulosus</i>	Morphology identification	–	–	–	Ditya et al. (2018)	Native
Clupeidae	<i>Namatolosa flyensis</i>	Morphology identification and DNA barcoding	–	–	–	Wibowo et al. (2017); Ditya et al. (2018)	Native
Cyprinidae	<i>Labeobarbus natalensis</i>	eDNA/BLAST	96%	16	0.01	Present study	Exotic
Eleotridae	<i>Oxyeleotris lineolata</i>	eDNA/BLAST	95%	2	0.00	Present study	Native
Gempylidae	<i>Diplospinus multistriatus</i>	eDNA/BLAST	95%	16	0.01	Present study	Native
Gobiidae	<i>Neogobius melanostomus</i>	eDNA/BLAST	95%	3378	1.14	Present study	Native
Gobiidae	<i>Smilosicyopus bitaeniatus</i>	eDNA/BLAST	96%	4473	1.50	Present study	Exotic
Melanotaenidae	<i>Iriatherina weneri</i>	DNA barcoding	–	–	–	Wibowo et al. (2017)	Native
Melanotaenidae	<i>Melanotaenia goldie</i>	DNA barcoding	–	–	–	Wibowo et al. (2017)	Native
Melanotaenidae	<i>Melanotaenia splendida inornata</i>	DNA barcoding	–	–	–	Wibowo et al. (2017)	Native
Melanotaeniidae	<i>Melanotaenia albimarginata</i>	eDNA/BLAST	95%	55	0.02	Present study	Native
Melanotaeniidae	<i>Melanotaenia goldiei</i>	eDNA/BLAST	96%	95 279	32.03	Present study	Native
Melanotaeniidae	<i>Melanotaenia sp</i>	eDNA/BLAST	95%	4	0.00	Present study	Native
Osphronemidae	<i>Trichopodus microlepis</i>	eDNA/BLAST	98%	12	0.00	Present study	Exotic
Osteoglossidae	<i>Scleropages jardinii</i>	DNA barcoding	–	–	–	Wibowo et al. (2017)	Native
Plotosidae	<i>Neosilurus ater</i>	Morphology identification and DNA barcoding	–	–	–	Wibowo et al. (2017); Ditya et al. (2018)	Native

(Continued on next page)

Table 1. (Continued).

Family	Species	Identification method	Similarity	Reads	Abundance (%)	Reference	Status
Plotosidae	<i>Porochilus meraukensis</i>	Morphology identification	–	–	–	Ditya et al. (2018)	Native
Serranidae	<i>Pseudanthias dispar</i>	eDNA/BLAST	98%	2	0.00	Present study	Native
Terapontidae	<i>Hephaestus raymondi</i>	Morphology identification	–	–	–	Ditya et al. (2018)	Native
Terapontidae	<i>Pinggala lorentzi</i>	Morphology identification and DNA barcoding	–	–	–	Wibowo et al. (2017); Ditya et al. (2018)	Native
Toxotidae	<i>Toxotes oligolepis</i>	Morphology identification and DNA barcoding	–	–	–	Wibowo et al. (2017); Ditya et al. (2018)	Native

For species detected using eDNA, percent of similarity to the closest species in Genbank, number of reads and relative frequencies are provided.





Fig. 2. Relative abundance of species-specific reads across PCR replicates at (a) Alfasera, (b) Yakui, (c) Sakor, and (d) Inggun.

## Discussion

Environmental DNA was successfully deployed to characterise freshwater fishes species richness and species occurrences at the Kumbe River. This represents one of the first eDNA metabarcoding assessments of fish biodiversity

in Indonesia. Interestingly, *N. graeffei* or blue catfish was the dominant fish species (90%) in the Kumbe River based on monthly fishing over 2 years (Ditya et al. 2018). However, eDNA metabarcoding revealed only five *N. graeffei* reads from the whole dataset, while 91.9% of reads were mapped to only three species (*M. goldiei*, *C. randi* and



**Fig. 3.** Venn diagram of identified number of shared fish species among four sampling locations.


*O. niloticus*). Thus, the lack of consensus between earlier studies employing traditional fishing methods (Wibowo et al. 2017; Ditya et al. 2018) and current eDNA analysis requires further investigation.

Our results showed that only five of the species we detected had been previously recorded (Wibowo et al. 2017; Ditya et al. 2018), and 18 represented new records, highlighting the limited knowledge of fish biodiversity in the Kumbe River. The detection of 18 new species is an important addition and highlights the critical role metabarcoding could have in rapidly describing fish biodiversity in tropical ecosystems. However, given that very low number of mapped reads (<50) for most of the species, the occurrence of these fishes

in the River Kumbe needs to be validated by traditional approaches. Furthermore, it is possible that fish biodiversity in the Kumbe River is greater than reported here given many sequences were produced in this study that could not be identified due to gaps in the reference database. This demonstrates that it is critical that the reference databases be expanded to include more taxonomic groups to improve our understanding of tropical fish biodiversity (Schenekar et al. 2020) by taking every opportunity to collect fin clips from species where identification has been verified.

The inability of metabarcoding to detect 17 species previously recorded in the Kumbe River could be due to a range of factors. From a technical perspective, several reasons may account for this discrepancy: (1) some species may not have been present at the time of eDNA sampling (e.g. migratory species); (2) failure to detect a species using eDNA may be due to low abundance, for example, earlier studies have shown that eDNA metabarcoding may generate false negative results for rare species or lineages (Sato et al. 2017; Burian et al. 2021; Xiong et al. 2022); (3) the incomplete detection of species based on eDNA may be caused by insufficient volume of water filtered, filtration method, PCR inhibitors, primer bias, preferential amplification of abundant species, or an incomplete database of reference sequences (Deiner et al. 2017; Majaneva et al. 2018; Bessey et al. 2020; Schenekar et al. 2020; Rojahn et al. 2021).

We chose to use the *Cyt-b* gene fragment as it is commonly applied for fish metabarcoding studies. Some studies have



**Fig. 4.** Fish community structure (proportion of total reads) at family (a) and genus (b) levels detected using eDNA metabarcoding at four different sites of the Kumbe River.

demonstrated that *Cyt-b* may not perform as well as other regions, such as the 12S rRNA gene (Zhang *et al.* 2020; Shu *et al.* 2021). However, not all studies show consistent patterns and there may be distinct advantages and disadvantages with both markers (Hänfling *et al.* 2016). It is possible that the number of species detected may be maximised using multiple markers, which also enables the verification of taxa detected in each sample (Lecaudey *et al.* 2019). We therefore recommend that future metabarcoding studies in Indonesia evaluate the performance of multiple mitochondrial and nuclear markers to identify the most powerful targets for species identification.

From an ecological perspective, there are several limitations to characterising biodiversity based on eDNA metabarcoding. Environmental DNA production, degradation and transport are the three key factors that determine the quantity of eDNA in the water (Hansen *et al.* 2018). Thus several abiotic factors such as temperature, pH, dissolved oxygen and organic matter can degrade eDNA and reduce species detectability. In this study we did not account for any environmental factors, and thus detectability of some species may have been reduced. If eDNA metabarcoding was to become more widely applied in the tropical waters of Indonesia for conservation purposes – particularly for detection and quantification of specific species – we recommend further research into the role that abiotic and biotic factors play in eDNA production and degradation (Stewart 2019). In addition, false-positives detections may be caused by misidentification of taxa or contamination during field sampling or in the laboratory (Sato *et al.* 2017; Fujii *et al.* 2019). Thus, more comparative studies at species-rich regions are needed to evaluate when eDNA metabarcoding estimates of species richness are on par with, or potentially better than traditional fisheries methods (Ficetola *et al.* 2015; Olds *et al.* 2016; Cilleros *et al.* 2019). Nevertheless, despite the limitations, we have demonstrated that eDNA metabarcoding is a promising method to assess biodiversity in Indonesia. We recommend that eDNA metabarcoding be deployed concurrently with traditional methods for optimal results while methods are being optimised and reference databases are being improved.

Fish biomonitoring using eDNA has been implemented in the detection of invasive species and assessing their distribution (Nathan *et al.* 2015; Dunker *et al.* 2016). The invasive Nile tilapia *O. niloticus* was detected at all sites, which is concerning given it is implicated in extirpations of native fish species (Ogutu-Ohwayo 1990; Ligetvoet *et al.* 1991). The Nile tilapia is an important source of animal protein across its native range, and as such, it has been widely introduced in Papua for aquaculture purposes. Environmental DNA metabarcoding in Indonesian waterways may be a rapid and cost-effective method to track new introductions or expansion of Nile tilapia.

The Kumbe River drains into the Arafura Sea and is expected to contain a diverse fish community comprised of

potamodromous species as well as species that require access to both freshwater and seawater to complete their lifecycle. Given the Kumbe River is a pristine aquatic habitat where land use has had minimal negative impacts on the environment (Wibowo *et al.* 2017; Lasmana *et al.* 2018), eDNA metabarcoding could be used to determine if migratory species are able to access upstream habitat as expected (Duda *et al.* 2021). In addition, eDNA metabarcoding could also be used to monitor spawning patterns of migratory fish species (either potamodromous or diadromous) (Yamanaka and Minamoto 2016; Thalinge *et al.* 2019). For example, if migratory species diversity is similar at upstream and downstream locations, this indicates good habitat connectivity between studied locations (Yamanaka and Minamoto 2016; Carraro *et al.* 2018). We detected *A. australis*, known as catadromous species, at the most upstream site (Alfasera). This likely suggests that the lower reaches of the Kumbe River, where this study was conducted, have sufficient connectivity to allow fish passage upstream. However, as this species is able to climb over migration barriers (Jellyman *et al.* 2017), *A. australis* presence upstream may not indicate that there is clear passage upstream for other species with lower upstream migration capabilities.

## Conclusion

Environmental DNA metabarcoding shows great promise as a novel tool for fish biomonitoring programs in tropical freshwater habitats. It is sensitive, non-invasive and cost-effective as well as enabling a rapid assessment for detection of invasive, rare or endangered species. It provides an alternative approach when traditional sampling is time consuming and requires taxonomic expertise. However, our results highlight that eDNA metabarcoding cannot be routinely applied without further research into sampling design, marker choice, seasonal data and an improved reference databases to reduce false negative detections. Despite this, eDNA metabarcoding will play an important role in the future characterisation of the fish community in tropical river systems.

## References

- Anonymous (2010) Inventarisasi of river in Papua, Indonesia in 2010. Available at <https://lingkunganhidup.papua.go.id/gi/fckimage/file/SLHD/Tabel%20SD-11.pdf>
- Austin KG, Schwantes A, Gu Y, Kasibhatla PS (2019) What causes deforestation in Indonesia? *Environmental Research Letters* 14(2), 024007. doi:10.1088/1748-9326/aaf6db
- Bartkowski B, Lienhoop N, Hansjürgens B (2015) Capturing the complexity of biodiversity: a critical review of economic valuation studies of biological diversity. *Ecological Economics* 113, 1–14. doi:10.1016/j.ecolecon.2015.02.023
- Bessey C, Jarman SN, Berry O, Olsen YS, Bunce M, Simpson T, Power M, McLaughlin J, Edgar GJ, Keesing J (2020) Maximizing fish detection with eDNA metabarcoding. *Environmental DNA* 2(4), 493–504. doi:10.1002/edn3.74



- Burian A, Mauvisseau Q, Bulling M, Domisch S, Qian S, Sweet M (2021) Improving the reliability of eDNA data interpretation. *Molecular Ecology Resources* 21(5), 1422–1433. doi:10.1111/1755-0998.13367
- Bylemans J, Gleeson DM, Duncan RP, Hardy CM, Furlan EM (2019) A performance evaluation of targeted eDNA and eDNA metabarcoding analyses for freshwater fishes. *Environmental DNA* 1(4), 402–414. doi:10.1002/edn3.41
- Carraro L, Hartikainen H, Jokela J, Bertuzzo E, Rinaldo A (2018) Estimating species distribution and abundance in river networks using environmental DNA. *Proceedings of the National Academy of Sciences* 115(46), 11724–11729. doi:10.1073/pnas.1813843115
- Cilleros K, Valentini A, Allard L, Dejean T, Etienne R, Grenouillet G, Iribar A, Taberlet P, Vigouroux R, Brosse S (2019) Unlocking biodiversity and conservation studies in high-diversity environments using environmental DNA (eDNA): a test with Guianese freshwater fishes. *Molecular Ecology Resources* 19(1), 27–46. doi:10.1111/1755-0998.12900
- Cleary DFR, DeVantier L (2011) Indonesia: threats to the country's biodiversity. In 'Encyclopedia of environmental health'. (Ed. JO Nriagu) pp. 187–197. (Elsevier)
- Dahrudin H, Utama A, Busson F, Sauri S, Hanner R, Keith P, Hadiaty R, Hubert N (2017) Revisiting the ichthyodiversity of Java and Bali through DNA barcodes: taxonomic coverage, identification accuracy, cryptic diversity and identification of exotic species. *Molecular Ecology Resources* 17(2), 288–299. doi:10.1111/1755-0998.12528
- de Bruyn M, Rüber L, Nylinder S, Stelbrink B, Lovejoy NR, Lavoué S, Tan HH, Nugroho E, Wowor D, Ng PKL, Siti Azizah MN, Von Rintelen T, Hall R, Carvalho GR (2013) Paleo-drainage basin connectivity predicts evolutionary relationships across three Southeast Asian biodiversity hotspots. *Systematic Biology* 62(3), 398–410. doi:10.1093/sysbio/syt007
- de Bruyn M, Stelbrink B, Morley RJ, Hall R, Carvalho GR, Cannon CH, van den Bergh G, Meijaard E, Metcalfe I, Boitani L, Maiorano L, Shoup R, von Rintelen T (2014) Borneo and Indochina are major evolutionary hotspots for Southeast Asian biodiversity. *Systematic Biology* 63(6), 879–901. doi:10.1093/sysbio/syu047
- Deiner K, Bik HM, Mächler E, Seymour M, Lacoursière-Roussel A, Altermatt F, Creer S, Bista I, Lodge DM, de Vere N, Pfrender ME, Bernatchez L (2017) Environmental DNA metabarcoding: transforming how we survey animal and plant communities. *Molecular Ecology* 26(21), 5872–5895. doi:10.1111/mec.14350
- Ditya YC, Wibowo A, Husnah H (2018) Spatial and temporal distribution of fish in the floodplain of Kumbur River. Papua. *Indonesian Fisheries Research Journal* 24(2), 69–81. doi:10.15578/ifrj.24.1.2018.1-13
- Djalil VN, Farajallah A, Wardiyatno Y (2018) Application of environmental DNA (eDNA) for detection of *Cherax quadricarinatus* (Von Martens 1868) using water sample. *Jurnal Biologi Tropis* 18(2), 134–140. doi:10.29303/jbt.v18i2.878
- Duda JJ, Hoy MS, Chase DM, Pess GR, Brenkman SJ, McHenry MM, Ostberg CO (2021) Environmental DNA is an effective tool to track recolonizing migratory fish following large-scale dam removal. *Environmental DNA* 3(1), 121–141. doi:10.1002/edn3.134
- Dunker KJ, Sepulveda AJ, Massengill RL, Olsen JB, Russ OL, Wenburg JK, Antonovich A (2016) Potential of environmental DNA to evaluate Northern Pike (*Esox lucius*) eradication efforts: an experimental test and case study. *PLoS ONE* 11(9), e0162277. doi:10.1371/journal.pone.0162277
- Ficetola GF, Miaud C, Pompanon F, Taberlet P (2008) Species detection using environmental DNA from water samples. *Biology Letters* 4(4), 423–425. doi:10.1098/rsbl.2008.0118
- Ficetola GF, Pansu J, Bonin A, Coissac E, Giguët-Covex C, De Barba M, Gielly L, Lopes CM, Boyer F, Pompanon F, Raye G, Taberlet P (2015) Replication levels, false presences and the estimation of the presence/absence from eDNA metabarcoding data. *Molecular Ecology Resources* 15(3), 543–556. doi:10.1111/1755-0998.12338
- Fujii K, Doi H, Matsuoka S, Nagano M, Sato H, Yamanaka H (2019) Environmental DNA metabarcoding for fish community analysis in backwater lakes: a comparison of capture methods. *PLoS ONE* 14(1), e0210357. doi:10.1371/journal.pone.0210357
- Garg T, Hamilton SE, Hochard JP, Kresch EP, Talbot J (2018) (Not so) gently down the stream: river pollution and health in Indonesia. *Journal of Environmental Economics and Management* 92, 35–53. doi:10.1016/j.jeem.2018.08.011
- Goldberg CS, Turner CR, Deiner K, Klymus KE, Thomsen PF, Murphy MA, Spear SF, McKee A, Oyler-McCance SJ, Cornman RS, Laramie MB, Mahon AR, Lance RF, Pilliod DS, Strickler KM, Waits LP, Fremier AK, Takahara T, Herder JE, Taberlet P, Gilbert M (2016) Critical considerations for the application of environmental DNA methods to detect aquatic species. *Methods in Ecology and Evolution* 7(11), 1299–1307. doi:10.1111/2041-210X.12595
- Gustiano R, Kurniawan K, Haryono H (2021) Optimizing the utilization of genetic resources of Indonesian native freshwater fish. *Asian Journal of Conservation Biology* 10(2), 189–196. doi:10.53562/ajcb.67022
- Hänfling B, Lawson Handley L, Read DS, Hahn C, Li J, Nichols P, Blackman RC, Oliver A, Winfield IJ (2016) Environmental DNA metabarcoding of lake fish communities reflects long-term data from established survey methods. *Molecular Ecology* 25(13), 3101–3119. doi:10.1111/mec.13660
- Hansen BK, Bekkevold D, Clausen LW, Nielsen EE (2018) The sceptical optimist: challenges and perspectives for the application of environmental DNA in marine fisheries. *Fish and Fisheries* 19(5), 751–768. doi:10.1111/faf.12286
- Herder F, Schliwen UK, Geiger MF, Hadiaty RK, Gray SM, McKinnon JS, Walter RP, Pfaender J (2012) Alien invasion in Wallace's dreamponds: records of the hybridogenic "flowerhorn" cichlid in Lake Matano, with an annotated checklist of fish species introduced to the Malili Lakes system in Sulawesi. *Aquatic Invasions* 7(4), 521–535. doi:10.3391/ai.2012.7.4.009
- Hubert N, Kadarusman, Wibowo A, Busson F, Caruso D, Sulandari S, Nafiqoh N, Pouyau L, Rüber L, Avarre J-C, Herder F, Hanner R, Keith P, Hadiaty RK (2015) DNA barcoding Indonesian freshwater fishes: challenges and prospects. *DNA Barcodes* 3(1), 144–169. doi:10.1515/dna-2015-0018
- Hubert N, Lumbantobing D, Sholihah A, Dahrudin H, Delrieu-Trottin E, Busson F, Sauri S, Hadiaty R, Keith P (2019) Revisiting species boundaries and distribution ranges of *Nemacheilus* spp. (Cypriniformes: Nemacheilidae) and *Rasbora* spp. (Cypriniformes: Cyprinidae) in Java, Bali and Lombok through DNA barcodes: implications for conservation in a biodiversity hotspot. *Conservation Genetics* 20(3), 517–529. doi:10.1007/s10592-019-01152-w
- Huson DH, Auch AF, Qi J, Schuster SC (2007) MEGAN analysis of metagenomic data. *Genome Research* 17(3), 377–386. doi:10.1101/gr.5969107
- Imai N, Furukawa T, Tsujino R, Kitamura S, Yumoto T (2018) Factors affecting forest area change in Southeast Asia during 1980–2010. *PLoS ONE* 13(5), e0197391. doi:10.1371/journal.pone.0197391
- Itakura H, Wakiya R, Yamamoto S, Kaifu K, Sato T, Minamoto T (2019) Environmental DNA analysis reveals the spatial distribution, abundance, and biomass of Japanese eels at the river-basin scale. *Aquatic Conservation: Marine and Freshwater Ecosystems* 29(3), 361–373. doi:10.1002/aqc.3058
- Janosik AM, Johnston CE (2015) Environmental DNA as an effective tool for detection of imperiled fishes. *Environmental Biology of Fishes* 98(8), 1889–1893. doi:10.1007/s10641-015-0405-5
- Jellyman PG, Bauld JT, Crow SK (2017) The effect of ramp slope and surface type on the climbing success of shortfin eel (*Anguilla australis*) elvers. *Marine and Freshwater Research* 68(7), 1317–1324. doi:10.1071/MF16015
- Kadarusman, Hubert N, Hadiaty RK, Sudarto, Paradis E, Pouyau L (2012) Cryptic diversity in Indo-Australian rainbowfishes revealed by DNA barcoding: implications for conservation in a biodiversity hotspot candidate. *PLoS ONE* 7(7), e40627. doi:10.1371/journal.pone.0040627
- Koh LP, Kettle CJ, Sheil D, Lee TM, Giam X, Gibson L, Clements GR (2013) Biodiversity state and trends in Southeast Asia. In 'Encyclopedia of biodiversity'. (Ed. SA Levin) pp. 509–527. (Academic Press)
- Kottelat M, Whitten T (1996) Freshwater biodiversity in Asia with special reference to fish. World Bank Technical paper. (World Bank)
- Kurniawan K, Gustiano R, Kusmini II, Prakoso VA (2021) Genetic resources preservation and utilization of Indonesian native freshwater fish consumption. *Ecology, Environment and Conservation* 27(1), 227–233.
- Lasmana Y, Simanungkalit P, Gifariyono M, Sotyadarpita G, Triadi LB (2018) Potential of tidal lowland for irrigation development in

- Merauke Regency using hydrodynamic modelling 1D2D. *Jurnal Teknik Hidraulik* 9(1), 17–32. doi:10.32679/jth.v9i1.432
- Lecaudey LA, Schletterer M, Kuzovlev VV, Hahn C, Weiss SJ (2019) Fish diversity assessment in the headwaters of the Volga River using environmental DNA metabarcoding. *Aquatic Conservation: Marine and Freshwater Ecosystems* 29(10), 1785–1800. doi:10.1002/aqc.3163
- Ligtvoet W, Witte F, Goldschmidt T, van Oijen MJP, Wanink JH, Goudswaard PC (1991) Species extinction and concomitant ecological changes in Lake Victoria. *Netherlands Journal of Zoology* 42, 214–232. doi:10.1163/156854291X00298
- Mace GM, Norris K, Fitter AH (2012) Biodiversity and ecosystem services: a multilayered relationship. *Trends in Ecology & Evolution* 27(1), 19–26. doi:10.1016/j.tree.2011.08.006
- Majaneva M, Diserud OH, Eagle SHC, Bostrom E, Hajibabaei M, Ekrem T (2018) Environmental DNA filtration techniques affect recovered biodiversity. *Scientific Reports* 8(1), 4682. doi:10.1038/s41598-018-23052-8
- McElroy ME, Dressler TL, Titcomb GC, Wilson EA, Deiner K, Dudley TL, Eliason EJ, Evans NT, Gaines SD, Lafferty KD, Lamberti GA, Li Y, Lodge DM, Love MS, Mahon AR, Pfrender ME, Renshaw MA, Selkoe KA, Jerde CL (2020) Calibrating environmental DNA metabarcoding to conventional surveys for measuring fish species richness. *Frontiers in Ecology and Evolution* 8, 276. doi:10.3389/fevo.2020.00276
- Meyer M, Kircher M (2010) Illumina sequencing library preparation for highly multiplexed target capture and sequencing. *Cold Spring Harbour Protocols* 6, 1–10.
- Nathan LR, Jerde CL, Budny ML, Mahon AR (2015) The use of environmental DNA in invasive species surveillance of the Great Lakes commercial bait trade. *Conservation Biology* 29(2), 430–439. doi:10.1111/cobi.12381
- Nugraha MFI, Pouyaud L, Carman O, Widyastuti U, Junior MZ, Kadarusman, Avarre J-C (2015) Genetic diversity of Boeseman's rainbowfish (*Melanotaenia boesemani*) reared in Indonesian farms compared to endangered natural populations. *Tropical Conservation Science* 8(3), 796–812. doi:10.1177/194008291500800315
- Ogutu-Ohwayo R (1990) The decline of native fishes of Lake Victoria and Kyoga (East Africa) and the impact of the introduced species, other environmental characteristics on amphibian distribution and abundance in mountain lakes of Northern Spain. *Animal Conservation* 9, 171–178.
- Olds BP, Jerde CL, Renshaw MA, Li Y, Evans NT, Turner CR, Deiner K, Mahon AR, Brueseke MA, Shirey PD, Pfrender ME, Lodge DM, Lamberti GA (2016) Estimating species richness using environmental DNA. *Ecology and Evolution* 6(12), 4214–4226. doi:10.1002/ece3.2186
- Pauchard N (2017) Access and benefit sharing under the convention on biological diversity and its protocol: what can some numbers tell us about the effectiveness of the regulatory regime? *Resources* 6(1), 11. doi:10.3390/resources6010011
- Peel MC, Finlayson BL, McMahon TA (2007) Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences* 11(5), 1633–1644. doi:10.5194/hess-11-1633-2007
- Pereira HM, Ferrier S, Walters M, Geller GN, Jongman RHG, Scholes RJ, Bruford MW, Brummitt N, Butchart SHM, Cardoso AC, Coops NC, Dulloo E, Faith DP, Freyhof J, Gregory RD, Heip C, Höft R, Hurtt G, Jetz W, Karp DS, McGeoch MA, Obura D, Onoda Y, Pettorelli N, Reyers B, Sayre R, Scharlemann JPW, Stuart SN, Turak E, Walpole M, Wegmann M (2013) Essential biodiversity variables. *Science* 339(6117), 277–278. doi:10.1126/science.1229931
- Pont D, Rocle M, Valentini A, Civate R, Jean P, Maire A, Roset N, Schabuss M, Zornig H, Dejean T (2018) Environmental DNA reveals quantitative patterns of fish biodiversity in large rivers despite its downstream transportation. *Scientific Reports* 8(1), 10361. doi:10.1038/s41598-018-28424-8
- Rees HC, Maddison BC, Middleditch DJ, Patmore JRM, Gough KC (2014) REVIEW: the detection of aquatic animal species using environmental DNA – a review of eDNA as a survey tool in ecology. *Journal of Applied Ecology* 51(5), 1450–1459. doi:10.1111/1365-2664.12306
- Rice CJ, Larson ER, Taylor CA (2018) Environmental DNA detects a rare large river crayfish but with little relation to local abundance. *Freshwater Biology* 63(5), 443–455. doi:10.1111/fwb.13081
- Roesma DI, Djong HT, Janra MN, Aidil DR (2021) Freshwater vertebrates monitoring in Maninjau Lake, West Sumatra, Indonesia using environmental DNA. *Biodiversitas Journal of Biological Diversity* 22(5), 2794–2802. doi:10.13057/biodiv/d220543
- Rojahn J, Gleeson DM, Furlan E, Haeusler T, Bylemans J (2021) Improving the detection of rare native fish species in environmental DNA metabarcoding surveys. *Aquatic Conservation: Marine and Freshwater Ecosystems* 31(4), 990–997. doi:10.1002/aqc.3514
- Sato H, Sogo Y, Doi H, Yamanaka H (2017) Usefulness and limitations of sample pooling for environmental DNA metabarcoding of freshwater fish communities. *Scientific Reports* 7(1), 14860. doi:10.1038/s41598-017-14978-6
- Schenekar T, Schletterer M, Lecaudey LA, Weiss SJ (2020) Reference databases, primer choice, and assay sensitivity for environmental metabarcoding: lessons learnt from a re-evaluation of an eDNA fish assessment in the Volga headwaters. *River Research and Applications* 36(7), 1004–1013. doi:10.1002/rra.3610
- Seymour M (2019) Rapid progression and future of environmental DNA research. *Communications Biology* 2, 80. doi:10.1038/s42003-019-0330-9
- Sholihah A, Delrieu-Trottin E, Sukmono T, Dahrudin H, Risdawati R, Elvyra R, Wibowo A, Kustiati K, Busson F, Sauri S, Nurhaman U, Dounias E, Zein MSA, Fitriana Y, Utama IV, Muchlisin ZA, Agnès J-F, Hanner R, Wowor D, Steinke D, Keith P, Rüber L, Hubert N (2020) Disentangling the taxonomy of the subfamily Rasborinae (Cypriniformes, Danionidae) in Sundaland using DNA barcodes. *Scientific Reports* 10(1), 2818. doi:10.1038/s41598-020-59544-9
- Sholihah A, Delrieu-Trottin E, Condamine FL, Wowor D, Rüber L, Pouyaud L, Agnès J-F, Hubert N (2021a) Impact of pleistocene eustatic fluctuations on evolutionary dynamics in southeast asian biodiversity hotspots. *Systematic Biology* 70(5), 940–960. doi:10.1093/sysbio/syab006
- Sholihah A, Delrieu-Trottin E, Sukmono T, Dahrudin H, Pouzadoux J, Tilak M-K, Fitriana Y, Agnès J-F, Condamine FL, Wowor D, Rüber L, Hubert N (2021b) Limited dispersal and in situ diversification drive the evolutionary history of Rasborinae fishes in Sundaland. *Journal of Biogeography* 48(9), 2153–2173. doi:10.1111/jbi.14141
- Shu L, Ludwig A, Peng Z (2021) Environmental DNA metabarcoding primers for freshwater fish detection and quantification: in silico and in tanks. *Ecology and Evolution* 11(12), 8281–8294. doi:10.1002/ece3.7658
- Sigsgaard EE, Carl H, Møller PR, Thomsen PF (2015) Monitoring the near-extinct European weather loach in Denmark based on environmental DNA from water samples. *Biological Conservation* 183, 46–52. doi:10.1016/j.biocon.2014.11.023
- Sigsgaard EE, Jensen MR, Winkelmann IE, Møller PR, Hansen MM, Thomsen PF (2020) Population-level inferences from environmental DNA – current status and future perspectives. *Evolutionary Applications* 13(2), 245–262. doi:10.1111/eva.12882
- Simpfendorfer CA, Kyne PM, Noble TH, Goldsburly J, Basitita RK, Lindsay R, Shields A, Perry C, Jerry DR (2016) Environmental DNA detects critically endangered largemouth sawfish in the wild. *Endangered Species Research* 30, 109–116. doi:10.3354/esr00731
- Stewart KA (2019) Understanding the effects of biotic and abiotic factors on sources of aquatic environmental DNA. *Biodiversity and Conservation* 28(5), 983–1001. doi:10.1007/s10531-019-01709-8
- Taberlet P, Coissac E, Pompanon F, Brochmann C, Willerslev E (2012) Towards next-generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology* 21(8), 2045–2050. doi:10.1111/j.1365-294X.2012.05470.x
- Thalinger B, Wolf E, Traugott M, Wanzenböck J (2019) Monitoring spawning migrations of potamodromous fish species via eDNA. *Scientific Reports* 9(1), 15388. doi:10.1038/s41598-019-51398-0
- Trebitz AS, Hoffman JC, Darling JA, Pilgrim EM, Kelly JR, Brown EA, Chadderton WL, Egan SP, Grey EK, Hashsham SA, Klymus KE, Mahon AR, Ram JL, Schultz MT, Stepien CA, Schardt JC (2017) Early detection monitoring for aquatic non-indigenous species: optimizing surveillance, incorporating advanced technologies, and identifying research needs. *Journal of Environmental Management* 202, 299–310. doi:10.1016/j.jenvman.2017.07.045
- Turner CR, Uy KL, Everhart RC (2015) Fish environmental DNA is more concentrated in aquatic sediments than surface water. *Biological Conservation* 183, 93–102. doi:10.1016/j.biocon.2014.11.017

- Ulayya N, Andayani N, Maryanto AE (2020) Development of environmental DNA approaches to detect alligator gar (*Atractosteus spatula*) from water samples. *IOP Conference Series: Earth and Environmental Science* **481**, 012013. doi:10.1088/1755-1315/481/1/012013
- Valiere N, Taberlet P (2000) Urine collected in the field as a source of DNA for species and individual identification. *Molecular Ecology* **9**, 2150–2152.
- Wibowo A, Wahlberg N, Vasemägi A (2017) DNA barcoding of fish larvae reveals uncharacterised biodiversity in tropical peat swamps of New Guinea, Indonesia. *Marine and Freshwater Research* **68**(6), 1079–1087. doi:10.1071/MF16078
- Wibowo A, Shibuno T, Sulit VT (2018) The making of a center of excellence in science and technology on inland fisheries management: the SEAFDEC/IFRDMD. *Fish for the People* **16**(3), 28–33.
- Xiao Y, Ouyang Z, Xu W, Xiao Y, Zheng H, Xian C (2016) Optimizing hotspot areas for ecological planning and management based on biodiversity and ecosystem services. *Chinese Geographical Science* **26**(2), 256–269. doi:10.1007/s11769-016-0803-4
- Xiong F, Shu L, Zeng H, Gan X, He S, Peng Z (2022) Methodology for fish biodiversity monitoring with environmental DNA metabarcoding: the primers, databases and bioinformatic pipelines. *Water Biology and Security* **1**(1), 100007. doi:10.1016/j.watbs.2022.100007
- Yachi S, Loreau M (1999) Biodiversity and ecosystem productivity in a fluctuating environment: the insurance hypothesis. *Proceedings of the National Academy of Sciences* **96**(4), 1463–1468. doi:10.1073/pnas.96.4.1463
- Yamanaka H, Minamoto T (2016) The use of environmental DNA of fishes as an efficient method of determining habitat connectivity. *Ecological Indicators* **62**, 147–153. doi:10.1016/j.ecolind.2015.11.022
- Zhang S, Zhao J, Yao M (2020) A comprehensive and comparative evaluation of primers for metabarcoding eDNA from fish. *Methods in Ecology and Evolution* **11**(12), 1609–1625. doi:10.1111/2041-210X.13485

**Data availability.** The data used in this manuscript are organised by the first author (Arif Wibowo). Access to this data can be negotiated with him.

**Conflicts of interest.** The authors declare no conflicts of interest.

**Declaration of funding.** This study was supported financially by the Research Institute for Inland Fisheries and Extensions (RIIFE), Division of Genetics, Department of Biology, Turku University, the Erasmus Mundus Research Fellowship Programme, Academy of Finland, and the Estonian Ministry of Education and Research (Institutional Research Funding Project IUT8–2).

**Acknowledgements.** We thank Research Institute for Inland Fisheries and Extensions for all support including staff and budget through DIPA 2017–2019. We are also grateful to officers from local fisheries districts in West Papua for sampling collection and fruitful discussion and finally all the reviewers for improving the manuscript.

#### Author affiliations

<sup>A</sup>Research Center for Conservation of Marine and Inland Water Resources, National Research and Innovation Agency, Jalan Raya Jakarta-Bogor Km. 48 Cibinong, Bogor, West Java 16911, Indonesia.

<sup>B</sup>Gulbali Institute for Agriculture, Water and Environment, Charles Sturt University, PO Box 789, Albury, NSW 2640, Australia.

<sup>C</sup>Research Center for Fisheries, National Research and Innovation Agency, Jalan Raya Jakarta-Bogor Km. 48 Cibinong, Bogor, West Java 16911, Indonesia.

<sup>D</sup>New South Wales Department of Primary Industries, Narrandera Fisheries Centre, Narrandera, NSW 2700, Australia.

<sup>E</sup>Japan Fisheries Research and Education Agency, Fisheries Resources Institute, Fisheries Stock Assessment Center, Yokohama, Japan.

<sup>F</sup>Université Montpellier (UMR) 5554 Institut des sciences de l'évolution de Montpellier (ISEM) (IRD, UM, CNRS, EPHE), Université de Montpellier, Place Eugène Bataillon, Montpellier cedex 05 34095, France.

<sup>G</sup>Department of Aquatic Resources, Institute of Freshwater Research, Swedish University of Agricultural Sciences, Drottningholm, Sweden.

<sup>H</sup>Estonian University of Life Sciences, Institute of Veterinary Medicine and Animal Sciences, Tartu, Estonia.