






Seminatural areas act as reservoirs of genetic diversity for crop pollinators and natural enemies across Europe

Joaquín Ortego¹  | Matthias Albrecht² | Andrés Báldi³  | Trine Bilde⁴ | Sean Birk Bek Craig⁴ | José M. Herrera^{5,6} | Amelia S. C. Hood⁷  | David Kleijn⁸ | Corina Maurer^{2,9} | Francisco P. Molina¹ | Erik Öckinger¹⁰ | Simon G. Potts⁷ | Virginia Settepani⁴ | Philip Francis Thomsen⁴ | Marina Trillo¹ | Flóra Vajna³  | Elena Velado-Alonso¹ | Ignasi Bartomeus¹ 

¹Department of Ecology and Evolution, Estación Biológica de Doñana (EBD-CSIC), Seville, Spain

²Agroscope, Agroecology and Environment, Zurich, Switzerland

³Lendület Ecosystem Services Research Group, Institute of Ecology and Botany, HUN-REN Centre for Ecological Research, Vácrátót, Hungary

⁴Department of Biology, Centre for Ecological Genetics, Aarhus University, Aarhus, Denmark

⁵Departamento de Biología, Instituto de Investigación Vitivinícola y Agroalimentaria, Universidad de Cádiz, Puerto Real, Spain

⁶Instituto Mediterráneo para a Agricultura Ambiente e Desenvolvimento, Universidade de Évora, Évora, Portugal

⁷Centre for Agri-Environmental Research (CAER), School of Agriculture, Policy and Development, Reading University, Reading, UK

⁸Plant Ecology and Nature Conservation Group, Wageningen University and Research, Wageningen, The Netherlands

⁹Ecosystems Landscape Evolution, Institute of Terrestrial Ecosystems, Department of Environmental Systems Science, ETH Zürich, Zürich, Switzerland

¹⁰Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden

Correspondence

Joaquín Ortego and Ignasi Bartomeus, Department of Ecology and Evolution, Estación Biológica de Doñana (EBD-CSIC), Avda. Américo Vespucio 26, E-41092 Seville, Spain.
Email: joaquin.ortego@csic.es and nacho.bartomeus@gmail.com

Funding information

Novo Nordisk Challenge Programme, Grant/Award Number: NNF20OC0060118; European Union's Horizon 2020 Research and Innovation Programme, Grant/Award Number: 862480; Programa QUALIFICA, Junta de Andalucía, Consejería de Universidad, Investigación e Innovación, Grant/Award Number: QUAL21020EBD

Abstract

Despite increasing recognition of the importance of the multiple dimensions of biodiversity, including functional or genetic diversity as well as species diversity, most conservation studies on ecosystem service-providing insects focus on simple diversity measures such as species richness and abundance. In contrast, relatively little is known about the genetic diversity and resilience of pollinators or natural enemies of crop pests to population fragmentation and local extinction. The genetic diversity and demographic dynamics of remnant populations of beneficial insects in agricultural areas can be a useful indicator providing additional insights into their conservation status, but this is rarely evaluated. Although gene flow between agricultural and seminatural areas is key to maintaining genetic diversity, its extent and directionality remain largely unexplored. Here, we apply a pan-European sampling protocol to quantify genetic diversity and structure and assess gene flow between agricultural and nearby seminatural landscapes in populations of two key ecosystem service-providing insect species, the lady beetle *Coccinella septempunctata*, an important predator of aphids and

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Authors. *Conservation Science and Practice* published by Wiley Periodicals LLC on behalf of Society for Conservation Biology.

other crop pests, and the bee pollinator *Andrena flavipes*. We show that *A. flavipes* populations are genetically structured at the European level, whereas populations of *C. septempunctata* experience widespread gene flow across the continent and lack any defined genetic structure. In both species, we found that there is high genetic connectivity between populations established in croplands and nearby seminatural areas and, as a consequence, they harbor similar levels of genetic diversity. Interestingly, demographic models for some regions support asymmetric gene flow from seminatural areas to nearby agricultural landscapes. Collectively, our study demonstrates how seminatural areas can serve as genetic reservoirs of both pollinators and natural enemies for nearby agricultural landscapes, acting as sources for recurrent recolonization and, potentially, contributing to enhancing ecosystem service and crop production resilience in the longer term.

KEYWORDS

agricultural landscapes, conservation genetics, demography, gene flow, genetic diversity, genetic structure, habitat fragmentation, service-providing insects

1 | INTRODUCTION

With more than 80% of the world's land under the direct influence of agricultural systems, conservation strategies that include agricultural landscapes could potentially be important to buffer and connect protected areas, but also in the direct conservation of species that exist there (Garibaldi et al., 2021; Kleijn et al., 2020). Indeed, protecting those species is becoming particularly important in providing ecosystem services such as enhanced crop yield by pollinators (Klein et al., 2007) and control of pest species by natural enemies (Garratt et al., 2011). Advances have been made in transitioning to lower-intensity agriculture and increasingly adopting an ecological intensification approach (Kleijn et al., 2019), with most efforts focused on halting the loss of species. For example, it has been shown that actions such as planting flower strips can increase pollinator richness and abundance (Scheper et al., 2013), and reduced soil tilling regimes improve the survival and numbers of arthropods in agroecosystems (Thorbek & Bilde, 2004). However, in addition to restoring and maintaining species diversity, it is necessary to maintain healthy and viable populations, that is, populations with a sufficient number of individuals to maintain high genetic diversity, which is key to responding to environmental change, and therefore for the long-term persistence of natural populations (Falconer & Mackay, 1996; Frankham, 1995; Spielman et al., 2004). This calls for integrating knowledge of population genetic diversity in the management and conservation of biodiversity (Kardos et al., 2021; Willi et al., 2022).

Although it is increasingly acknowledged that it is important to consider multiple dimensions of biodiversity in conservation, most studies to date focus on simple diversity metrics at the level of species, such as richness and abundance (Cerini et al., 2023; Kardos et al., 2021; Willi et al., 2022). Although species diversity metrics have been core for evaluating the effectiveness of conservation and restoration measures, they cannot provide insights into some of the more complex mechanisms underlying biodiversity responses. Biodiversity comprises many levels of organization, from community composition and functional diversity to species metapopulation dynamics and genetic diversity (e.g., United Nations Convention on Biological Diversity; www.cbd.int). In particular, genetic diversity is an often-neglected aspect with far-reaching implications for long-term conservation (Kardos et al., 2021; e.g., Saccheri et al., 1998; Spielman et al., 2004). More genetically diverse populations may better withstand global change pressures, as these populations have a larger genetic pool from which to adapt to new environmental conditions (Frankham, 2005; Kardos et al., 2021). However, the impact of land-use changes on the genetic diversity of arthropods and the connectivity between populations from seminatural and agricultural landscapes in insect species that provide important ecosystem functions and services are still poorly known (González et al., 2020; Keyghobadi, 2007; Schlaepfer et al., 2018; Webster et al., 2023).

Insects and other arthropods providing ecosystem services are often highly mobile organisms that disperse and forage across habitats in heterogeneous landscapes

(e.g., Jaffé et al., 2016; Lecompte et al., 2016). This implies that their persistence and conservation does not only rely on local habitat quality but also on the landscape-level habitat composition and configuration, in particular, connectivity between suitable habitats (Kremen et al., 2007; Martin et al., 2019). A generally observed pattern is that croplands surrounded by a larger proportion of natural habitats tend to support relatively higher beneficial insect diversity and abundance, which facilitates ecosystem services such as pollination and pest control (Dainese et al., 2019). However, most insect conservation studies focus on a single habitat type (either agricultural or seminatural habitats) and rarely integrate different habitats within the same landscape (Marini et al., 2019). In fact, our understanding of how source-sink metapopulation dynamics work in agricultural systems is still very limited (e.g., Ballare & Jha, 2021; Dreier et al., 2014; Jha, 2015; Ortego, Aguirre, et al., 2015). Many species are able to maintain sustainable populations in agricultural habitats, but other species may not be able to persist in simplified agricultural habitats without continuous immigration from nearby natural or seminatural areas, that is, the agricultural habitats act as population sinks and the (semi-)natural areas as population sources (e.g., Öckinger & Smith, 2007a, 2007b). Either way, the benefits of having bidirectional gene flow between natural and agricultural areas can be important, even when populations would be able to persist in isolation; for example, this would promote long-term persistence of species if one area is subject to a major disturbance, as the remaining populations in neighboring habitats can repopulate and contribute to gene flow toward the disturbed area. However, in what way natural areas contribute to the genetic diversity of ecosystem service providing insect populations in nearby agricultural landscapes and vice versa has, to our knowledge, not yet been explored (Webster et al., 2023).

To improve our understanding of the importance of natural habitats for the conservation of population genetic diversity of key ecosystem service-providing insects, we used a pan-European sampling design to characterize gene flow dynamics between populations of the pest control agent *Coccinella septempunctata* (Coleoptera: Coccinellidae) and the bee pollinator *Andrena flavipes* (Hymenoptera: Andrenidae) in agricultural areas and nearby seminatural landscapes. These are two of the most common ecosystem service providers in Europe, with *A. flavipes* ranking 7th in a recent review of the most common bee crop pollinators in Europe (Kleijn et al., 2015) whereas *C. septempunctata* is a widespread predator of aphids (Jervis & Kidd, 1996; New, 1991), a ubiquitous pest of several crops. To this end, we sampled and genotyped

populations of the two focal species in agricultural and nearby seminatural areas across Europe (i.e., a paired-replicated approach), quantified their respective levels of population genetic diversity and structure, and tested alternative models of gene flow. Specifically, we first (i) tested the hypothesis that populations in seminatural areas sustain larger effective population sizes and, thus, present higher levels of genetic diversity, than those established in agricultural landscapes. We then quantified spatial patterns of genetic structure across the different European regions, and (ii) tested the hypothesis that populations of mobile beneficial insects from seminatural and agricultural landscapes regularly exchange gene flow and show limited genetic structure (e.g., Jaffé et al., 2016; Lecompte et al., 2016). Finally, we used a coalescent-based model-testing approach to evaluate alternative demographic scenarios and (iii) test the hypothesis that seminatural areas play a role as sources of gene flow for populations established in agricultural landscapes (Figure 1). Our study demonstrates how seminatural areas can serve as genetic reservoirs of both pollinators and natural enemies for nearby agricultural landscapes, acting as sources for recurrent recolonization and, potentially, contributing to enhance ecosystem services and crop production resilience in the longer term. These results, general across a network of European sites, directly appeal to practitioners and policy makers—for example, through the European Union's Common Agricultural Policy (CAP)—and emphasize the importance of implementing management actions aimed at preserving or restoring seminatural habitats for the maintenance of source populations of ecosystem service-providing insects in agricultural landscapes.

2 | MATERIALS AND METHODS

2.1 | Population sampling

In spring and summer 2021, we sampled pairs of populations of *C. septempunctata* and *A. flavipes* in agricultural landscapes (i.e., dominated by agriculturally managed land for the last 100 years) and nearby seminatural areas (i.e., dominated by seminatural habitat that was not under agricultural management at least in the last 100 years) separated by <25 km (mean: 11.9 km; range: 4.6–24.0 km; Table 1; Figure 2; Figure S1). The selected species are representative of generalist ecosystem service providers within their guild and are often locally abundant and widely distributed. In seminatural areas, specimens were collected in patches of native vegetation, whereas specimens from agricultural areas were collected




FIGURE 1 (a) Schematic illustrating the hypothesis of unidirectional gene flow from seminatural to nearby agricultural landscapes for populations of the pest predator *Coccinella septempunctata* (shown in panel a) and the pollinator *Andrena flavipes*. Panel (b) shows the four alternative gene flow models tested using *FASTSIMCOAL2*. Parameters include mutation-scaled ancestral effective population sizes (ϑ_{ANC}), contemporary effective population sizes for populations from agricultural (ϑ_{AGR}) and seminatural (ϑ_{NAT}) landscapes, timing of population split (T_{DIV}), and symmetric (m) and unidirectional (m_{AN} and m_{NA}) migration rates.

within crop fields and along paths and field margins. We targeted to sample 8–10 specimens per species and site area and aimed to replicate this sampling scheme in seven European countries: Estonia, Denmark, the Netherlands, Great Britain, Hungary, Switzerland, and Spain (Figure 2; Figure S1). However, low population densities in some areas often reduced our sample sizes and one of the two focal taxa could not be collected in some countries at a minimum sample size of $n = 5$ in agricultural and/or seminatural sites (Table 1). Consequently, our sampling scheme could be replicated in six countries for *C. septempunctata* and four countries for *A. flavipes* and final datasets included 5–10 genotyped specimens per species, country, and area (mean = 8; mode = 8; Table 1). Our pan-European sampling scheme, and the countries selected for our study, frames within the context of a multipartner project funded by the European Union's Horizon 2020 Research and Innovation Programme H2020. We spread the sampling across each site (<500 m of radius around sampling coordinates; Table 1) to avoid collecting closely related individuals (i.e., siblings or half-siblings). Specimens were collected by hand (*C. septempunctata*) or sweep-netting (*A. flavipes*) and preserved at -20°C in 1500 μL ethanol 96% until DNA extraction. We registered spatial coordinates of each sampling plot using a Global Positioning System (GPS). Further details on sampling locations are provided in Table 1.

2.2 | Genomic library preparation and processing

We extracted and purified DNA from each specimen using NucleoSpin Tissue kits (Macherey-Nagel, Düren, Germany). We processed genomic DNA into different genomic libraries using the double-digestion restriction-fragment-based procedure (ddRAD-seq) described in Peterson et al. (2012). In brief, we digested DNA with the restriction enzymes MseI and EcoRI (New England Biolabs, Ipswich, MA, USA) and ligated Illumina adaptors including unique 7-bp barcodes to the digested fragments of each individual. We pooled ligation products, size-selected them between 350 and 450 bp with a Pippin Prep machine (Sage Science, Beverly, MA, USA), and amplified the fragments by PCR with 12 cycles using the iProof™ high-fidelity DNA polymerase (BIO-RAD, Venendaal, The Netherlands). Single-read 201-bp sequencing was performed on an Illumina NovaSeq6000 platform. We used the different programs distributed as part of the *STACKS* v. 1.35 pipeline (Catchen et al., 2013) to filter and assemble our sequences into de novo loci, call genotypes, calculate genetic diversity statistics, and export input files for all downstream analyses (Methods S1). Unless otherwise indicated, all downstream analyses were performed using datasets including only the first single nucleotide polymorphism (SNP) per RAD locus (180-bp length, see Methods S1) and retaining loci with a

TABLE 1 Geographical location, number of genotyped individuals (n), and genetic diversity statistics (π , H_O , and H_E) for populations of *Coccinella septempunctata* and *Andrena flavipes* sampled in agricultural and nearby seminatural landscapes.

| Country | Code | Site | Latitude | Longitude | n | All positions | | | Variant positions | | |
|--------------------------------------|------|--------------|----------|-----------|-----|---------------|--------|--------|-------------------|--------|--------|
| | | | | | | π | H_O | H_E | π | H_O | H_E |
| (a) <i>Coccinella septempunctata</i> | | | | | | | | | | | |
| Estonia | EE | Agricultural | 58.82436 | 23.59144 | 10 | 0.0003 | 0.0002 | 0.0003 | 0.0541 | 0.0299 | 0.0504 |
| Estonia | EE | Seminatural | 58.79168 | 23.51234 | 8 | 0.0003 | 0.0002 | 0.0003 | 0.0537 | 0.0298 | 0.0489 |
| Denmark | DK | Agricultural | 56.29121 | 10.40226 | 8 | 0.0003 | 0.0002 | 0.0003 | 0.0539 | 0.0318 | 0.0489 |
| Denmark | DK | Seminatural | 56.28052 | 10.46685 | 10 | 0.0003 | 0.0002 | 0.0003 | 0.0551 | 0.0310 | 0.0514 |
| Netherlands | NL | Agricultural | 50.79752 | 5.79459 | 8 | 0.0003 | 0.0002 | 0.0003 | 0.0564 | 0.0309 | 0.0517 |
| Netherlands | NL | Seminatural | 50.82924 | 5.87514 | 8 | 0.0003 | 0.0002 | 0.0003 | 0.0546 | 0.0311 | 0.0497 |
| Great Britain | GB | Agricultural | 50.65034 | -2.28968 | 8 | 0.0003 | 0.0002 | 0.0003 | 0.0527 | 0.0288 | 0.0482 |
| Great Britain | GB | Seminatural | 50.64681 | -2.00065 | 8 | 0.0003 | 0.0002 | 0.0003 | 0.0557 | 0.0308 | 0.0509 |
| Hungary | HU | Agricultural | 46.93410 | 19.10340 | 8 | 0.0003 | 0.0002 | 0.0003 | 0.0541 | 0.0297 | 0.0495 |
| Hungary | HU | Seminatural | 47.11210 | 19.28140 | 10 | 0.0003 | 0.0002 | 0.0003 | 0.0565 | 0.0309 | 0.0526 |
| Switzerland | CH | Agricultural | 47.06575 | 7.24068 | 8 | 0.0003 | 0.0002 | 0.0003 | 0.0532 | 0.0326 | 0.0486 |
| Switzerland | CH | Seminatural | 47.19120 | 7.31626 | 8 | 0.0003 | 0.0002 | 0.0003 | 0.0541 | 0.0308 | 0.0493 |
| (b) <i>Andrena flavipes</i> | | | | | | | | | | | |
| Denmark | DK | Agricultural | 56.29121 | 10.40226 | 8 | 0.0003 | 0.0003 | 0.0003 | 0.0608 | 0.0493 | 0.0558 |
| Denmark | DK | Seminatural | 56.27405 | 10.46778 | 6 | 0.0003 | 0.0003 | 0.0003 | 0.0555 | 0.0482 | 0.0505 |
| Netherlands | NL | Agricultural | 50.79752 | 5.79459 | 8 | 0.0004 | 0.0003 | 0.0004 | 0.0711 | 0.0557 | 0.0654 |
| Netherlands | NL | Seminatural | 50.82924 | 5.87514 | 7 | 0.0004 | 0.0003 | 0.0004 | 0.0738 | 0.0599 | 0.0672 |
| Switzerland | CH | Agricultural | 47.06575 | 7.24068 | 8 | 0.0004 | 0.0004 | 0.0004 | 0.0771 | 0.0652 | 0.0710 |
| Switzerland | CH | Seminatural | 47.02291 | 7.31958 | 8 | 0.0004 | 0.0003 | 0.0004 | 0.0777 | 0.0625 | 0.0716 |
| Spain | ES | Agricultural | 37.32987 | -6.19802 | 10 | 0.0004 | 0.0003 | 0.0004 | 0.0800 | 0.0572 | 0.0747 |
| Spain | ES | Seminatural | 37.23660 | -6.18450 | 5 | 0.0004 | 0.0003 | 0.0004 | 0.0766 | 0.0561 | 0.0662 |

Note: Genetic diversity statistics were calculated in STACKS for all positions (polymorphic and nonpolymorphic) and only variant (polymorphic) positions. Average values across loci are presented for nucleotide diversity (π), and observed (H_O) and expected (H_E) heterozygosity.

minimum stack depth ≥ 5 ($m = 5$) and that were represented in at least 75% of the populations ($p = 9$ for *C. septempunctata* and $p = 6$ for *A. flavipes*) and 50% of the individuals within each population ($r = 0.5$).

2.3 | Quantifying genetic structure

We calculated genetic differentiation between each pair of populations using the Weir and Cockerham weighted fixation index (F_{ST}) (Weir & Cockerham, 1984) as implemented in ARLEQUIN v. 3.5 (Excoffier & Lischer, 2010). We determined statistical significance with Fisher's exact tests after 10,000 permutations, applying a false discovery rate (FDR) adjustment (5%, $q < 0.05$) to control for multiple tests. We performed Mantel tests to analyze the association between genetic differentiation (F_{ST}) and geographical distances among populations of each focal species using ZT software with 10,000 permutations (Bonnet

& van de Peer, 2002). We quantified population genetic structure and admixture using the Bayesian Markov chain Monte Carlo clustering method implemented in the program STRUCTURE v. 2.3.3 (Pritchard et al., 2000). We conducted STRUCTURE analyses hierarchically, initially analyzing data from all populations of each taxon jointly and, subsequently, running independent analyses for subsets of populations assigned to the same genetic cluster in the previous hierarchical level analysis (e.g., González-Serna et al., 2019). To make analyses computationally tractable, we ran STRUCTURE using a random subset of 10,000 SNPs. We ran STRUCTURE analyses assuming correlated allele frequencies and admixture and without using prior population information (Hubisz et al., 2009). We conducted 15 independent runs for each value of K (from $K = 1$ to $K = 10$) to estimate the most likely number of genetic clusters with 200,000 MCMC cycles, following a burn-in step of 100,000 iterations. We retained the ten runs having the highest likelihood for




FIGURE 2 Map showing the location of study sites in seven European countries, each including an agricultural (brown dots) and a nearby seminatural (green triangles) plot. Geographical coordinates are given in Table 1 and satellite images for each sampling site are presented in Figure S1. CH, Switzerland; DK, Denmark; EE, Estonia; ES, Spain; GB, Great Britain; HU, Hungary; NL, Netherlands.

each value of K and determined the number of genetic clusters that best describes our data according to log probabilities of the data ($\text{LnPr}(X|K)$) (Pritchard et al., 2000) and the ΔK method (Evanno et al., 2005), as implemented in *STRUCTURE HARVESTER* (Earl & vonHoldt, 2012). We used *CLUMPP* v. 1.1.2 and the Greedy algorithm to align multiple runs of *STRUCTURE* for the same K value (Jakobsson & Rosenberg, 2007) and *DISTRUCT* v. 1.1 (Rosenberg, 2004) to visualize the individuals' probabilities of population membership in bar plots. Complementarily, we used the same datasets of 10,000 random SNPs to perform principal component analyses (PCA) as implemented in the *R* v. 4.2.1 (R Core Team, 2023) package “*adegenet*” (Jombart, 2008). Before running PCAs, we replaced missing data by the mean allele frequency of the corresponding locus estimated across all samples (Jombart, 2008). Results from PCAs were visualized using the *R* package “*scatterplot3d*” (Ligges & Mächler, 2003).

2.4 | Testing alternative models of gene flow

We used the simulation-based approach implemented in *FASTSIMCOAL2* to evaluate alternative migration models and test the hypothesis of asymmetric gene flow between populations from agricultural landscapes and nearby seminatural areas (Excoffier et al., 2013; Figure 1a). Specifically, we tested four alternative migration models for each taxon (*C. septempunctata* and *A. flavipes*) and spatial replicate

(i.e., country), including a null-model considering no gene flow between populations from agricultural and seminatural areas (Model A), a model of symmetric gene flow (Model B), and two models of unidirectional gene flow, one fitting unidirectional gene flow from agricultural to seminatural areas (Model C) and another exclusively considering unidirectional gene flow in the opposite direction (Model D) (Figure 1b). We used the site frequency spectrum (SFS) and *FASTSIMCOAL2* to estimate the composite likelihood of the observed data given a specified model (Excoffier et al., 2013). For each taxon and pair of populations, we calculated a folded joint SFS considering a single SNP per locus. Because we did not include invariable sites in the SFS, we fixed the contemporary effective population size for one of the demes (θ_{AGR} in all cases; see Figure 1b) to enable the estimation of other parameters in *FASTSIMCOAL2* (Excoffier et al., 2013). The effective population size fixed in the model was calculated from the level of nucleotide diversity (π) and estimates of mutation rates per site per generation (μ). Nucleotide diversity (π) was estimated from polymorphic and nonpolymorphic loci using *STACKS* (Table 1). We used the mutation rate per site per generation (2.8×10^{-9}) estimated for *Drosophila melanogaster* (Keightley et al., 2014), which is similar to the spontaneous mutation rate estimated for the butterfly *Heliconius melpomene* (2.9×10^{-9} ; Keightley et al., 2015). To remove all missing data for the calculation of the SFS, minimize errors with allele frequency estimates, and maximize the number of retained SNPs, each population group was downsampled using a Python script written by Qixin He

and available on Dryad (Papadopoulou & Knowles, 2015). Note that this downsampling procedure necessarily results in the number of SNPs retained in the SFS differ across each pair of analyzed populations, as this depends on the number of available samples and the pattern and amount of missing data in each specific dataset (Table 2; e.g., Papadopoulou & Knowles, 2015).

Each model was run 100 independent times considering 100,000–250,000 simulations for the calculation of the composite likelihood, 10–40 expectation-conditional maximization (ECM) cycles, and a stopping criterion of 0.001 (Excoffier et al., 2013; Excoffier & Foll, 2011). We used an information-theoretic model selection approach

based on the Akaike's information criterion (AIC) to compare the set of candidate models and identify the one that is best supported by the data (Burnham & Anderson, 2002; e.g., Ortego et al., 2021; Thome & Carstens, 2016). After the maximum likelihood was estimated for each model in every replicate, we calculated the AIC scores as detailed in Thome and Carstens (2016). AIC values for each model were rescaled (ΔAIC), calculating the difference between the AIC value of each model and the minimum AIC obtained among all competing models (i.e., the best model has $\Delta\text{AIC} = 0$). Point estimates of the different demographic parameters for the best supported model were

TABLE 2 Alternative demographic models (detailed in Figure 1b) tested using FASTSIMCOAL2 for pairs of populations of *Coccinella septempunctata* and *Andrena flavipes* sampled in agricultural and nearby seminatural landscapes from different countries.

| Model | lnL | k | AIC | ΔAIC | ω_i | lnL | k | AIC | ΔAIC | ω_i |
|--------------------------------------|-----------------|----------|------------------|--------------------|-------------|--------------------------------|----------|------------------|--------------------|-------------|
| (a) <i>Coccinella septempunctata</i> | | | | | | | | | | |
| Estonia (EE) (3860 SNPs) | | | | | | Denmark (DK) (5127 SNPs) | | | | |
| A | -4405.22 | 3 | 8816.44 | 108.65 | 0.00 | -5498.61 | 3 | 11,003.21 | 156.03 | 0.00 |
| B | -4354.34 | 4 | 8716.67 | 8.88 | 0.01 | -5423.18 | 4 | 10,854.36 | 7.17 | 0.03 |
| C | -4352.98 | 4 | 8713.97 | 6.17 | 0.04 | -5425.60 | 4 | 10,859.20 | 12.01 | 0.00 |
| D | -4349.90 | 4 | 8707.79 | 0.00 | 0.95 | -5419.59 | 4 | 10,847.18 | 0.00 | 0.97 |
| Netherlands (NL) (2586 SNPs) | | | | | | Great Britain (GB) (2933 SNPs) | | | | |
| A | -2852.51 | 3 | 5711.03 | 79.03 | 0.00 | -3263.30 | 3 | 6532.60 | 47.78 | 0.00 |
| B | -2813.05 | 4 | 5634.09 | 2.10 | 0.21 | -3239.54 | 4 | 6487.07 | 2.25 | 0.16 |
| C | -2813.02 | 4 | 5634.05 | 2.05 | 0.21 | -3238.41 | 4 | 6484.82 | 0.00 | 0.50 |
| D | -2812.00 | 4 | 5631.99 | 0.00 | 0.59 | -3238.81 | 4 | 6485.62 | 0.79 | 0.34 |
| Hungary (HU) (5822 SNPs) | | | | | | Switzerland (CH) (2769 SNPs) | | | | |
| A | -6373.50 | 3 | 12,753.00 | 170.70 | 0.00 | -3046.27 | 3 | 6098.53 | 51.44 | 0.00 |
| B | -6291.05 | 4 | 12,590.09 | 7.78 | 0.02 | -3021.12 | 4 | 6050.24 | 3.15 | 0.15 |
| C | -6289.01 | 4 | 12,586.01 | 3.70 | 0.13 | -3021.22 | 4 | 6050.44 | 3.35 | 0.13 |
| D | -6287.15 | 4 | 12,582.31 | 0.00 | 0.85 | -3019.54 | 4 | 6047.09 | 0.00 | 0.72 |
| (b) <i>Andrena flavipes</i> | | | | | | | | | | |
| Denmark (DK) (3389 SNPs) | | | | | | Netherlands (NL) (6668 SNPs) | | | | |
| A | -5262.58 | 3 | 10,531.16 | 6.09 | 0.03 | -8193.79 | 3 | 16,393.59 | 161.97 | 0.00 |
| B | -5260.36 | 4 | 10,528.71 | 3.64 | 0.11 | -8115.47 | 4 | 16,238.93 | 7.31 | 0.02 |
| C | -5258.54 | 4 | 10,525.07 | 0.00 | 0.66 | -8113.91 | 4 | 16,235.81 | 4.19 | 0.11 |
| D | -5262.40 | 4 | 10,532.81 | 7.74 | 0.01 | -8111.81 | 4 | 16,231.62 | 0.00 | 0.87 |
| Switzerland (CH) (7492 SNPs) | | | | | | Spain (ES) (5623 SNPs) | | | | |
| A | -8787.18 | 3 | 17,580.35 | 169.26 | 0.00 | -5389.38 | 3 | 10,784.75 | 103.39 | 0.00 |
| B | -8703.52 | 4 | 17,415.04 | 3.95 | 0.10 | -5339.27 | 4 | 10,686.54 | 5.17 | 0.06 |
| C | -8701.55 | 4 | 17,411.09 | 0.00 | 0.75 | -5339.14 | 4 | 10,686.27 | 4.91 | 0.07 |
| D | -8703.21 | 4 | 17,414.42 | 3.33 | 0.14 | -5336.68 | 4 | 10,681.36 | 0.00 | 0.86 |

Note: Best-supported scenarios are highlighted in bold. The number of variable single nucleotide polymorphisms (SNPs) retained to calculate the site frequency spectrum is indicated in parentheses. Population codes as described in Table 1.

Abbreviations: AIC, Akaike's information criterion value; k, number of parameters in the model; lnL, maximum likelihood estimate of the model; ΔAIC , difference in AIC value from that of the strongest model; ω_i , AIC weight.

selected from the run with the highest maximum composite likelihood. Finally, we calculated confidence intervals of parameter estimates (based on the percentile method; e.g., de Manuel et al., 2016) from 100 parametric bootstrap replicates by simulating SFS from the maximum composite likelihood estimates and re-estimating parameters each time (Excoffier et al., 2013).

3 | RESULTS

3.1 | Samples and genomic data

Overall, we obtained samples of *C. septempunctata* and *A. flavipes* from six and four countries, respectively, with three countries sampling both species (Table 1). The average number of reads retained per individual after the different quality filtering steps was 3,409,688 (range = 236,373–9,557,006 reads) for *C. septempunctata* (Figure S2) and 2,103,765 (range = 420,468–6,255,835 reads) for *A. flavipes* (Figure S3). On average, this represented 86% (range = 64%–90%) and 73% (range = 60%–81%) of the total number of reads recovered for each individual of *C. septempunctata* and *A. flavipes*, respectively. After filtering loci (see Methods S1), the final dataset including all genotyped populations of *C. septempunctata* contained 10,896 SNPs, with a mean coverage depth of $25\times$ (mode = $35\times$; range = $12\text{--}42\times$) and an average proportion of missing data of 23% (mode = 10%; range = 0%–77%). For *A. flavipes*, the final dataset contained 18,554 SNPs, with a mean coverage depth of $25\times$ (mode = $26\times$; range = $7\text{--}39\times$) and an average proportion of missing data of 27% (mode = 24%; range = 17%–77%).

3.2 | Genetic diversity of populations

Levels of genetic diversity in populations of *C. septempunctata* and *A. flavipes* are presented in Table 1. In *C. septempunctata*, individual genetic diversity (i.e., observed heterozygosity) did not significantly differ among populations sampled in different countries (one-way ANOVA: $F_{5,96} = 0.99$, $p = .423$) or between populations from agricultural landscapes and nearby seminatural areas within any of the countries (all p -values $> .3$). In *A. flavipes*, individual genetic diversity significantly differed among populations sampled in different countries (one-way ANOVA: $F_{3,56} = 5.85$, $p = .002$) and post hoc Tukey's tests revealed that such differences resulted from comparisons involving populations from Denmark, which showed lower levels of genetic diversity than populations from the other countries ($p < .06$ in all cases; Table 1). As in *C. septempunctata*,

individual genetic diversity did not significantly differ between populations of *A. flavipes* from agricultural landscapes and nearby seminatural areas within countries (all p -values $> .3$).

3.3 | Quantifying genetic structure

In *C. septempunctata*, all pairwise F_{ST} values were non-significantly different from zero (Table S1). In contrast, pairwise F_{ST} values in *A. flavipes* ranged between 0 and 0.444, and several pairs of populations showed significant genetic differentiation (Table S2). Except for comparisons involving populations from the Netherlands and Switzerland, all pairs of populations of *A. flavipes* from different countries were significantly differentiated (Table S2). Within most countries, populations of *A. flavipes* from agricultural landscapes and nearby seminatural areas did not show significant genetic differentiation (Table S2). The only exception was the comparison involving the two populations from Denmark, albeit with a small F_{ST} value ($F_{ST} = 0.03$; Table S2). Simple Mantel tests revealed that genetic differentiation (F_{ST}) is positively correlated with geographical distances among populations in *A. flavipes* ($r = 0.96$, $p < .001$, $n = 8$) but not in *C. septempunctata* ($r = 0.00$, $p = 1.000$, $n = 12$). STRUCTURE analyses for *C. septempunctata* showed that $\text{LnPr}(X|K)$ peaked at $K = 2$ and decreased for higher K -values (Figure S4a). For $K = 2$, all individuals and populations of *C. septempunctata* presented a very low probability of assignment ($q < 0.05$) to one of the two genetic clusters (i.e., a “fictive” or “ghost” cluster sensu Guillot et al., 2005; see also Chen et al., 2007), which indicates a lack of genetic structure across analyzed populations of this taxon (Figure 3a). Clustering solutions for higher K -values (from $K = 3$ to $K = 5$) did not reveal either any genetic structure in *C. septempunctata* (Figure S5a). STRUCTURE analyses for *A. flavipes* considering all populations identified the most likely number of clusters as $K = 2$ according to the ΔK criterion and $\text{LnPr}(X|K)$ reached a plateau at the same K value (Figure S4b). For $K = 2$, the two genetic clusters separated the populations of *A. flavipes* from Spain from the rest of European populations (Figure 3b). STRUCTURE analyses at a lower hierarchical level, and excluding Spanish populations, identified the most likely number of clusters as $K = 2$ according to the ΔK criterion and $\text{LnPr}(X|K)$ reached a plateau at the same K value (Figure S4c). In this case, the two genetic clusters separated the populations from Denmark from the populations from the Netherlands and Switzerland, which grouped together (Figure 3b). The new genetic clusters that arise for higher K -values (from $K = 3$ to $K = 5$) were either similarly represented in all populations of *A. flavipes* assigned to the same

genetic cluster for $K = 2$ or were circumscribed to certain populations but showed very low probabilities assignment in all individuals ($q < 0.15$; Figure S5b,c). Principal component analyses (PCA) confirmed the results yielded by STRUCTURE for the two taxa (Figure 4). For *C. septempunctata*, PCA did not reveal any genetic clustering and individuals collected in different countries and sites were interspersed across the three first principal components (Figure 4a). For *A. flavipes*, PCA separated

Spanish populations from the rest of European populations along the first principal component (PC1) (Figure 4b,c). The second principal component (PC2) separated the populations from Denmark from those sampled in the Netherlands and Switzerland, which also tended to segregate along the third principal component (PC3) (Figure 4b). Analyses for *A. flavipes* excluding Spanish populations provided similar results, with PC1 separating populations from Denmark from those sampled in




FIGURE 3 Results of genetic assignments based on the Bayesian method implemented in the program STRUCTURE for populations of (a) *Coccinella septempunctata* and (b) *Andrena flavipes* sampled in agricultural (agr.) and nearby seminatural (nat.) landscapes from different countries. STRUCTURE analyses for *A. flavipes* were performed for all populations jointly (top) and excluding populations from Spain (bottom). Analyses are based on a random subset of 10,000 single nucleotide polymorphisms (SNPs). Each individual is represented by a vertical bar, which is partitioned into K colored segments showing the individual's probability of belonging to the cluster with that color. Thin vertical black lines separate individuals from different populations. Population codes as described in Table 1.




FIGURE 4 Principal component analyses (PCA) of genetic variation for populations of (a) *Coccinella septempunctata* and (b, c) *Andrena flavipes* sampled in agricultural (agr.; dots) and nearby seminatural (nat.; triangles) landscapes from different countries. Principal component analyses for *A. flavipes* were performed for (b) all populations jointly and (c) excluding populations from Spain. Analyses are based on a random subset of 10,000 single nucleotide polymorphisms (SNPs). Population codes as described in Table 1.

the Netherlands and Switzerland, which in this case tended to segregate along PC2 (Figure 4c).

3.4 | Testing alternative models of gene flow

Demographic analyses in FASTSIMCOAL2 supported unidirectional gene flow from populations in seminatural areas to those in agricultural landscapes (Model D; Figure 1b) in five out of six *C. septempunctata* (EE, DK, NL, HU, and CH) and two out of four *A. flavipes* (NL and ES) pairwise comparisons (Table 2). In the rest of the comparisons (*C. septempunctata* in GB and *A. flavipes* in DK and CH), the best supported model was a scenario of unidirectional gene flow from populations in agricultural landscapes to those in nearby seminatural areas (Model C; Table 2). In the case of populations of *C. septempunctata* from Great Britain (GB), the second-best supported model (Model D) was statistically indistinguishable from the best supported model (Model C) ($\Delta\text{AIC} = 0.79$), indicating that the two alternative models of unidirectional gene flow are similarly probable (Table 2). Estimates of migration rates between populations were fairly similar across all pairwise comparisons, ranging from 2.35×10^{-4} to 5.79×10^{-4} in *C. septempunctata* and from 2.27×10^{-4} to 5.67×10^{-4} in *A. flavipes* (Table 3). In both *C. septempunctata* and *A. flavipes*, estimates of gene flow from populations established in agricultural areas to those in nearby seminatural areas (i.e., Model C) were

smaller than estimates obtained when the most supported model was the scenario of unidirectional gene flow in the opposite direction (i.e., Model D) (Table 3).

4 | DISCUSSION

Our data support contrasting patterns of genetic structure and gene flow in two key ecosystem service providing insect species, the natural enemy *C. septempunctata* and the pollinator *A. flavipes*. A panmictic structure characterizes populations of *C. septempunctata* across both local and regional scales. Conversely, populations of *A. flavipes* present a marked structure at a European scale, but a very limited or null genetic differentiation between nearby agricultural and seminatural landscapes within each country. We found no support for the hypothesis of higher levels of genetic diversity in populations from seminatural areas with respect to those established in agricultural landscapes. However, demographic model testing and coalescent-based estimates of migration rates lent partial support to the hypothesis of increased gene flow from seminatural to agricultural landscapes (Figure 1a). For *C. septempunctata*, models considering anisotropic gene flow in the predicted direction received comparatively higher statistical support than the rest of the scenarios across most landscape replicates (i.e., countries), while the evidence on this respect for *A. flavipes* was limited to 50% of the regions. Collectively, these results emphasize the occurrence of gene flow from seminatural to

TABLE 3 Parameters inferred from coalescent simulations with FASTSIMCOAL2 under the most supported demographic model (Model C or Model D, as indicated in Table 2) for pairs of populations of *Coccinella septempunctata* and *Andrena flavipes* sampled in agricultural and nearby seminatural landscapes from different countries.

| Code | θ_{ANC} | | θ_{NAT} | | T_{DIV} | | $m_{\text{NA}}/m_{\text{AN}}$ | |
|--------------------------------------|-----------------------|----------------|-----------------------|----------------|------------------|---------------|-------------------------------|---|
| | Point | 95% CI | Point | 95% CI | Point | 95% CI | Point | 95% CI |
| (a) <i>Coccinella septempunctata</i> | | | | | | | | |
| EE ^D | 6579 | 5637–7291 | 24,889 | 19,358–33,796 | 6661 | 5626–8002 | 5.57×10^{-4} | 4.17×10^{-4} – 7.08×10^{-4} |
| DK ^D | 8191 | 6885–8985 | 82,132 | 67,815–109,399 | 10,385 | 9313–12,554 | 2.81×10^{-4} | 2.34×10^{-4} – 3.41×10^{-4} |
| NL ^D | 5519 | 4383–6360 | 29,795 | 22,444–40,894 | 7090 | 5913–8789 | 5.34×10^{-4} | 4.03×10^{-4} – 7.31×10^{-4} |
| GB ^C | 13,077 | 10,278–16,041 | 113,536 | 88,194–169,780 | 10,098 | 7670–13,554 | 2.35×10^{-4} | 1.95×10^{-4} – 2.91×10^{-4} |
| HU ^D | 8179 | 6804–9029 | 61,597 | 51,126–75,089 | 9830 | 8959–11,936 | 2.76×10^{-4} | 2.35×10^{-4} – 3.12×10^{-4} |
| CH ^D | 5419 | 4551–6369 | 21,203 | 13,913–30,169 | 4445 | 3927–5945 | 5.79×10^{-4} | 3.90×10^{-4} – 8.21×10^{-4} |
| (b) <i>Andrena flavipes</i> | | | | | | | | |
| DK ^C | 118,487 | 59,169–126,548 | 51,303 | 20,647–66,866 | 8338 | 1857–17,100 | 2.27×10^{-4} | 1.67×10^{-4} – 2.86×10^{-4} |
| NL ^D | 13,751 | 11,544–14,635 | 39,466 | 31,734–51,439 | 15,015 | 12,642–18,271 | 4.79×10^{-4} | 3.91×10^{-4} – 5.73×10^{-4} |
| CH ^C | 18,113 | 13,911–21,774 | 127,634 | 99,263–170,344 | 12,537 | 10,399–16,085 | 3.74×10^{-4} | 3.04×10^{-4} – 4.42×10^{-4} |
| ES ^D | 4381 | 3832–4712 | 20,900 | 15,414–26,253 | 4847 | 4376–5551 | 5.67×10^{-4} | 4.46×10^{-4} – 7.42×10^{-4} |

cultivated areas and confirm the importance of preserving seminatural habitats at the landscape level for the maintenance of source populations of ecosystem service-providing insects (Garibaldi et al., 2021).

4.1 | Genetic diversity and structure at regional and landscape scales

Analyses of genetic differentiation (F_{ST}) and structure (STRUCTURE and PCAs) revealed marked differences in the spatial distribution of genomic variation in the two focal taxa (Figures 3 and 4). Populations of *C. septempunctata* were panmictic and presented no evidence for genetic differentiation at both European and landscape scales, a phenomenon likely explained by the long-distance migratory behavior linked to seasonal aggregations characterizing many coccinellids (Hagen, 1962). Long-distance dispersal in this taxon is exemplified by unrestricted gene flow between British and mainland European populations, which indicates the limited effects of seawater as a barrier to dispersal (i.e., The English Channel is 34 km at its narrowest). These results are in line with the findings of a previous microsatellite-based study covering the entire distribution of the species (Lecompte et al., 2016). This study revealed a lack of genetic structure, very limited differentiation ($F_{ST} < 0.05$), and a weak pattern of isolation-by-distance across populations sampled from Portugal to India. Only eastern Asia (China, Japan) and some North African populations (Algeria) were assigned to different genetic clusters and presented a significant genetic differentiation ($F_{ST} = 0.10\text{--}0.30$), which suggests either long-term isolation of these peripheral populations (e.g., in different glacial refugia) or the evolution of reproductive isolation and an incipient process of cryptic speciation (Lecompte et al., 2016).

In contrast, analyses for *A. flavipes* revealed a marked genetic structure and differentiation ($F_{ST} = 0.10\text{--}0.44$) at a European scale, with more peripheral populations from Jutland (DK) and Iberian (ES) peninsulas forming distinct genetic clusters with limited gene flow between them as well as with the rest of analyzed populations (Figures 3 and 4). Likewise, although central European populations from NL and CH (separated by >400 km) clustered together in STRUCTURE analyses (Figure 2; Figure S5) and showed nonsignificant genetic differentiation ($F_{ST} = 0$; Table S2), they tended to segregate in PCAs (Figure 4). These results reveal more limited gene flow in *A. flavipes* than in *C. septempunctata* at a continental scale, which has likely resulted in the development of genetic structure across populations from most studied regions. In contrast, populations from agricultural and nearby seminatural landscapes within countries presented weak genetic differentiation in *A. flavipes* ($F_{ST} < 0.03$). Only the two sites

within Denmark showed significant differentiation, albeit with a very low estimate ($F_{ST} = 0.03$). Remarkably, populations of *A. flavipes* from Denmark showed the lowest levels of genetic diversity compared to the rest of European populations, which suggests that demographic stochasticity, founder effects, genetic drift, and low effective population sizes characterizing leading-edge peripheral populations could have contributed to fine spatial-scale (<5 km) genetic structure between populations from agricultural and nearby seminatural landscapes within Jutland Peninsula (Hampe & Petit, 2005; Lira-Noriega & Manthey, 2014). Weak genetic structure at local and landscape spatial scales has been previously reported in other Andrenidae (e.g., *Andrena fuscipes*: Exeler et al., 2010; *Andrena vaga*: Černá et al., 2013; Exeler et al., 2008), bumblebees (e.g., *Bombus terrestris*; Silva et al., 2020), stingless bees (e.g., *Trigona spinipes*; Jaffé et al., 2016), and carpenter bees (e.g., *Xylocopa virginica*; Ballare & Jha, 2021).

Collectively, these results are in line with the findings of spatially explicit landscape genetic studies suggesting a limited impact of habitat fragmentation and structure on the distribution of spatial patterns of genetic variation in different pollinators (Barbosa et al., 2022; Exeler et al., 2010; Jackson et al., 2018; Jha & Kremen, 2013). Aligned with inferences from analyses of genetic structure, testing of alternative demographic models strongly rejected the scenario of strict isolation (i.e., total lack of gene flow; Model A) between populations from agricultural and natural areas in the two focal taxa (Table 2). Connectivity between nearby areas of different land use is expected to contribute to prevent genetic drift and loss of genetic diversity through time, which can explain similar levels of genetic variation across populations from natural and agricultural landscapes (Table 1).

4.2 | Anisotropic gene flow between agricultural and seminatural landscapes

Asymmetric gene flow has been previously documented linked to passive dispersal in the direction of prevailing winds or ocean currents (e.g., Bertola et al., 2020; Kling & Ackerly, 2021) and associated to differences in the production of propagules in the context of central-peripheral and source-sink metapopulation dynamics (Kirkpatrick & Barton, 1997; e.g., Hauser et al., 2019; Holliday et al., 2012). Our model-based approach provided mixed support to the hypothesis of directional gene flow from populations in seminatural areas to those sustained in nearby agricultural landscapes (Figure 1). Coalescent-based model testing in the natural enemy *C. septempunctata* largely supported unidirectional gene flow in the predicted direction (Model D), suggesting that seminatural areas

genetically subsidize conspecific populations established in nearby croplands. These results are in line with previous capture-mark-recapture and observational studies pointing to the role of seminatural landscapes as population sources for beneficial insects (Öckinger & Smith, 2007a, 2007b). Although analyses in *A. flavipes* consistently supported models fitting asymmetric migration rates over those considering symmetric gene exchange, the scenario of unidirectional gene flow from seminatural to agricultural landscapes was the most supported one in only half of our spatial replicates. Remarkably, the bidirectional migration model received low statistical support in all spatial replicates for both *C. septempunctata* and *A. flavipes*, suggesting that asymmetric gene flow and source-sink dynamics govern the demography of the focal taxa in natural-agricultural mosaic landscapes (Hanski, 1998). Under this scenario, source populations with a net growth supply individuals to sink populations, reducing the chances of extinction of small populations and contributing to increasing the stability and maintaining the levels of genetic diversity of the whole metapopulation (Howe et al., 1991; e.g., Ortego et al., 2008; Saccheri et al., 1998). It must be noted, however, that source-sink dynamics in highly anthropized landscapes are only expected to be established among generalist taxa with relatively high dispersal ability (e.g., Keller et al., 2013; Ortego, García-Navas, et al., 2015), such as the two widespread ecosystem service-providing species analyzed in our study (Jervis & Kidd, 1996; Kleijn et al., 2015; New, 1991). Rare species with more limited dispersal or increased sensitivity to population subdivision (e.g., pools of regionally threatened species) will likely experience dispersal depression, population isolation, reduced levels of genetic diversity, and increased chances of local extinction under the severe fragmentation of natural habitats that characterize most agricultural landscapes (e.g., Jauker et al., 2009; Ortego, Aguirre, et al., 2015; Schtickzelle et al., 2006).

5 | CONCLUSIONS

Collectively, our results support high genetic connectivity between populations established in agricultural and nearby seminatural landscapes, indicating that (i) the latter may serve—at least in some cases—as reservoirs of both natural enemies and pollinators for surrounding agricultural areas, (ii) act as sources for recurrent recolonization, and (iii) potentially contribute to enhance ecosystem services and crop production. This highlights the importance of conserving complex agricultural systems which embed seminatural areas within productive

regions, as those can boost gene flow exchange and contribute to demographic reinforcement of populations of ecosystem service-providing insects and increase the chances that they persist and maintain their important functional role through time.

AUTHOR CONTRIBUTIONS

JO and IB designed the study. IB coordinated the study. MA, AB, TB, SBBC, JMH, AH, DK, CM, FPM, EO, SGP, VS, PFT, FV, EV-A, and IB collected the samples. MT performed the lab work supervised by JO. JO analyzed the data. JO and IB drafted the manuscript. All authors revised the manuscript and contributed to writing.

ACKNOWLEDGMENTS

We thank Claus Rasmussen for help with the identification of sampling sites and bee species recognition in Denmark, Zsolt Józán for identification of bees from Hungary, Roel d'Haese and Carmen Näf for their help with the insect sampling in Switzerland, Gabriella Bishop and Reinier de Vries for field assistance in the Netherlands, Natural England and the National Trust for permission to sample on Studland and Godlingston Heaths, and Lulworth Estate for permission to sample on their land. We also thank two anonymous referees for their constructive and valuable comments on an earlier version of the manuscript. Logistic support was provided by Laboratorio de Ecología Molecular (LEM-EBD) from Estación Biológica de Doñana (CSIC). We also thank Centro de Supercomputación de Galicia (CESGA) and Doñana's Singular Scientific-Technical Infrastructure (ICTS-RBD) for access to computer resources. This study was funded by the project SHOWCASE (SHOWCASing synergies between agriculture, biodiversity, and ecosystem services to help farmers capitalizing on native biodiversity) within the European Union's Horizon 2020 Research and Innovation Programme (grant agreement No. 862480), and by Programa QUALIFICA, Junta de Andalucía, Consejería de Universidad, Investigación e Innovación (project QUAL21020EBD). The collection of samples in Denmark was funded by the Novo Nordisk Challenge Programme (grant NNF20OC0060118).

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

Raw Illumina reads have been deposited at the NCBI Sequence Read Archive (SRA) under BioProject PRJNA1036856. Input files for all analyses (STRUCTURE, PCA, ARLEQUIN, and FASTSIMCOAL2) are available for download from Figshare (<https://doi.org/10.6084/m9.figshare.24525019>).

ORCID

Joaquín Ortego  <https://orcid.org/0000-0003-2709-429X>

Andrés Báldi  <https://orcid.org/0000-0001-6063-3721>

Amelia S. C. Hood  <https://orcid.org/0000-0003-3803-0603>

Flóra Vajna  <https://orcid.org/0000-0003-4451-855X>

Ignasi Bartomeus  <https://orcid.org/0000-0001-7893-4389>

REFERENCES

- Ballare, K. M., & Jha, S. (2021). Genetic structure across urban and agricultural landscapes reveals evidence of resource specialization and philopatry in the Eastern carpenter bee, *Xylocopa virginica* L. *Evolutionary Applications*, 14(1), 136–149. <https://doi.org/10.1111/eva.13078>
- Barbosa, M. D., Jaffé, R., Carvalho, C. S., Lanes, E. C. M., Alves-Pereira, A., Zucchi, M. I., Corrêa, A. S., Ribeiro, M. C., Imperatriz-Fonseca, V. L., & Alves, D. A. (2022). Landscape influences genetic diversity but does not limit gene flow in a Neotropical pollinator. *Apidologie*, 53(4), 48. <https://doi.org/10.1007/s13592-022-00955-0>
- Bertola, L. D., Boehm, J. T., Putman, N. F., Xue, A. T., Robinson, J. D., Harris, S., Baldwin, C. C., Overcast, I., & Hickerson, M. J. (2020). Asymmetrical gene flow in five co-distributed syngnathids explained by ocean currents and rafting propensity. *Proceedings of the Royal Society B: Biological Sciences*, 287(1926), 20200657. <https://doi.org/10.1098/rspb.2020.0657>
- Bonnet, E., & van de Peer, Y. (2002). zt: A software tool for simple and partial Mantel tests. *Journal of Statistical Software*, 7(10), 1–12. <https://doi.org/10.18637/jss.v007.i10>
- Burnham, K. P., & Anderson, D. R. (2002). *Model selection and multimodel inference: A practical information-theoretic approach*. Springer.
- Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). STACKS: An analysis tool set for population genomics. *Molecular Ecology*, 22(11), 3124–3140. <https://doi.org/10.1111/mec.12354>
- Cerini, F., Childs, D. Z., & Clements, C. F. (2023). A predictive timeline of wildlife population collapse. *Nature Ecology & Evolution*, 7, 320–331. <https://doi.org/10.1038/s41559-023-01985-2>
- Černá, K., Straka, J., & Munclinger, P. (2013). Population structure of pioneer specialist solitary bee *Andrena vaga* (Hymenoptera: Andrenidae) in central Europe: The effect of habitat fragmentation or evolutionary history? *Conservation Genetics*, 14(4), 875–883. <https://doi.org/10.1007/s10592-013-0482-y>
- Chen, C., Durand, E., Forbes, F., & François, O. (2007). Bayesian clustering algorithms ascertaining spatial population structure: A new computer program and a comparison study. *Molecular Ecology Notes*, 7(5), 747–756. <https://doi.org/10.1111/j.1471-8286.2007.01769.x>
- Dainese, M., Martin, E. A., Aizen, M. A., Albrecht, M., Bartomeus, I., Bommarco, R., Carvalheiro, L. G., Chaplin-Kramer, R., Gagic, V., Garibaldi, L. A., Ghazoul, J., Grab, H., Jonsson, M., Karp, D. S., Kennedy, C. M., Kleijn, D., da Silva e Silva, F. D., de Groot, G. A., Dudenhöffer, J. H., ... Steffan-Dewenter, I. (2019). A global synthesis reveals biodiversity-mediated benefits for crop production. *Science Advances*, 5(10), eaax0121. <https://doi.org/10.1126/sciadv.aax0121>
- de Manuel, M., Kuhlwilm, M., Frandsen, P., Sousa, V. C., Desai, T., Prado-Martinez, J., Hernandez-Rodriguez, J., Dupanloup, I., Lao, O., Hallast, P., Schmidt, J. M., Heredia-Genestar, J. M., Benazzo, A., Barbujani, G., Peter, B. M., Kuderna, L. F., Casals, F., Angedakin, S., Arandjelovic, M., ... Marques-Bonet, T. (2016). Chimpanzee genomic diversity reveals ancient admixture with bonobos. *Science*, 354(6311), 477–481. <https://doi.org/10.1126/science.aag2602>
- Dreier, S., Redhead, J. W., Warren, I. A., Bourke, A. F. G., Heard, M. S., Jordan, W. C., Sumner, S., Wang, J., & Carvell, C. (2014). Fine-scale spatial genetic structure of common and declining bumble bees across an agricultural landscape. *Molecular Ecology*, 23(14), 3384–3395. <https://doi.org/10.1111/mec.12823>
- Earl, D. A., & vonHoldt, B. M. (2012). STRUCTURE HARVESTER: A website and program for visualizing structure output and implementing the Evanno method. *Conservation Genetics Resources*, 4(2), 359–361. <https://doi.org/10.1007/s12686-011-9548-7>
- Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: A simulation study. *Molecular Ecology*, 14(8), 2611–2620. <https://doi.org/10.1111/j.1365-294X.2005.02553.x>
- Excoffier, L., Dupanloup, I., Huerta-Sanchez, E., Sousa, V. C., & Foll, M. (2013). Robust demographic inference from genomic and SNP data. *PLoS Genetics*, 9(10), e1003905. <https://doi.org/10.1371/journal.pgen.1003905>
- Excoffier, L., & Foll, M. (2011). fastsimcoal: A continuous-time coalescent simulator of genomic diversity under arbitrarily complex evolutionary scenarios. *Bioinformatics*, 27(9), 1332–1334. <https://doi.org/10.1093/bioinformatics/btr124>
- Excoffier, L., & Lischer, H. E. L. (2010). ARLEQUIN suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources*, 10(3), 564–567. <https://doi.org/10.1111/j.1755-0998.2010.02847.x>
- Exeler, N., Kratochwil, A., & Hochkirch, A. (2008). Strong genetic exchange among populations of a specialist bee, *Andrena vaga* (Hymenoptera: Andrenidae). *Conservation Genetics*, 9(5), 1233–1241. <https://doi.org/10.1007/s10592-007-9450-8>
- Exeler, N., Kratochwil, A., & Hochkirch, A. (2010). Does recent habitat fragmentation affect the population genetics of a heathland specialist, *Andrena fuscipes* (Hymenoptera: Andrenidae)? *Conservation Genetics*, 11(5), 1679–1687. <https://doi.org/10.1007/s10592-010-0060-5>
- Falconer, D. S., & Mackay, T. F. C. (1996). *Introduction to quantitative genetics* (3rd ed.). Longman.
- Frankham, R. (1995). Conservation genetics. *Annual Review of Genetics*, 29, 305–327.
- Frankham, R. (2005). Genetics and extinction. *Biological Conservation*, 126(2), 131–140.
- Garibaldi, L. A., Oddi, F. J., Miguez, F. E., Bartomeus, I., Orr, M. C., Jobbagy, E. G., Kremen, C., Schulte, L. A., Hughes, A. C., Bagnato, C., Abramson, G., Bridgewater, P., Carella, D. G., Díaz, S., Dicks, L. V., Ellis, E. C., Goldenberg, M., Huaylla, C. A., Kuperman, M., ... Zhu, C. D. (2021). Working landscapes need at least 20% native habitat. *Conservation Letters*, 14(2), e12773. <https://doi.org/10.1111/conl.12773>
- Garratt, M. P. D., Wright, D. J., & Leather, S. R. (2011). The effects of farming system and fertilisers on pests and natural enemies:

- A synthesis of current research. *Agriculture Ecosystems & Environment*, 141(3–4), 261–270. <https://doi.org/10.1016/j.agee.2011.03.014>
- González, A. V., Gómez-Silva, V., Ramirez, M. J., & Fonturbel, F. E. (2020). Meta-analysis of the differential effects of habitat fragmentation and degradation on plant genetic diversity. *Conservation Biology*, 34(3), 711–720. <https://doi.org/10.1111/cobi.13422>
- González-Serna, M. J., Cordero, P. J., & Ortego, J. (2019). Spatio-temporally explicit demographic modelling supports a joint effect of historical barriers to dispersal and contemporary landscape composition on structuring genomic variation in a red-listed grasshopper. *Molecular Ecology*, 28(9), 2155–2172. <https://doi.org/10.1111/mec.15086>
- Guillot, G., Estoup, A., Mortier, F., & Cosson, J. F. (2005). A spatial statistical model for landscape genetics. *Genetics*, 170(3), 1261–1280. <https://doi.org/10.1534/genetics.104.033803>
- Hagen, K. S. (1962). Biology and ecology of predaceous *Coccinellidae*. *Annual Review of Entomology*, 7, 289–326. <https://doi.org/10.1146/annurev.en.07.010162.001445>
- Hampe, A., & Petit, R. J. (2005). Conserving biodiversity under climate change: The rear edge matters. *Ecology Letters*, 8(5), 461–467. <https://doi.org/10.1111/j.1461-0248.2005.00739.x>
- Hanski, I. (1998). Metapopulation dynamics. *Nature*, 396(6706), 41–49.
- Hauser, S. S., Walker, L., & Leberg, P. L. (2019). Asymmetrical gene flow of the recently delisted passerine black-capped vireo (*Vireo atricapilla*) indicates source-sink dynamics in central Texas. *Ecology and Evolution*, 9(1), 463–470. <https://doi.org/10.1002/ece3.4764>
- Holliday, J. A., Suren, H., & Aitken, S. N. (2012). Divergent selection and heterogeneous migration rates across the range of Sitka spruce (*Picea sitchensis*). *Proceedings of the Royal Society B: Biological Sciences*, 279(1734), 1675–1683. <https://doi.org/10.1098/rspb.2011.1805>
- Howe, R. W., Davis, G. J., & Mosca, V. (1991). The demographic significance of ‘sink’ populations. *Biological Conservation*, 57(3), 239–255. [https://doi.org/10.1016/0006-3207\(91\)90071-g](https://doi.org/10.1016/0006-3207(91)90071-g)
- Hubisz, M. J., Falush, D., Stephens, M., & Pritchard, J. K. (2009). Inferring weak population structure with the assistance of sample group information. *Molecular Ecology Resources*, 9(5), 1322–1332. <https://doi.org/10.1111/j.1755-0998.2009.02591.x>
- Jackson, J. M., Pimsler, M. L., Oyen, K. J., Koch-Uhuad, J. B., Herndon, J. D., Strange, J. P., Dillon, M. E., & Lozier, J. D. (2018). Distance, elevation and environment as drivers of diversity and divergence in bumble bees across latitude and altitude. *Molecular Ecology*, 27(14), 2926–2942. <https://doi.org/10.1111/mec.14735>
- Jaffé, R., Castilla, A., Pope, N., Imperatriz-Fonseca, V. L., Metzger, J. P., Arias, M. C., & Jha, S. (2016). Landscape genetics of a tropical rescue pollinator. *Conservation Genetics*, 17(2), 267–278. <https://doi.org/10.1007/s10592-015-0779-0>
- Jakobsson, M., & Rosenberg, N. A. (2007). CLUMPP: A cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics*, 23(14), 1801–1806. <https://doi.org/10.1093/bioinformatics/btm233>
- Jauker, F., Diekotter, T., Schwarzbach, F., & Wolters, V. (2009). Pollinator dispersal in an agricultural matrix: Opposing responses of wild bees and hoverflies to landscape structure and distance from main habitat. *Landscape Ecology*, 24(4), 547–555. <https://doi.org/10.1007/s10980-009-9331-2>
- Jervis, M., & Kidd, N. (1996). *Insect natural enemies: Practical approaches to their study and evaluation*. Chapman and Hall.
- Jha, S. (2015). Contemporary human-altered landscapes and oceanic barriers reduce bumble bee gene flow. *Molecular Ecology*, 24(5), 993–1006. <https://doi.org/10.1111/mec.13090>
- Jha, S., & Kremen, C. (2013). Urban land use limits regional bumble bee gene flow. *Molecular Ecology*, 22(9), 2483–2495. <https://doi.org/10.1111/mec.12275>
- Jombart, T. (2008). *adeigenet*: A R package for the multivariate analysis of genetic markers. *Bioinformatics*, 24(11), 1403–1405. <https://doi.org/10.1093/bioinformatics/btn129>
- Kardos, M., Armstrong, E. E., Fitzpatrick, S. W., Hauser, S., Hedrick, P. W., Miller, J. M., Tallmon, D. A., & Funk, W. C. (2021). The crucial role of genome-wide genetic variation in conservation. *Proceedings of the National Academy of Sciences of the United States of America*, 118(48), e2104642118. <https://doi.org/10.1073/pnas.2104642118>
- Keightley, P. D., Ness, R. W., Halligan, D. L., & Haddrill, P. R. (2014). Estimation of the spontaneous mutation rate per nucleotide site in a *Drosophila melanogaster* full-sib family. *Genetics*, 196(1), 313–320. <https://doi.org/10.1534/genetics.113.158758>
- Keightley, P. D., Pinharanda, A., Ness, R. W., Simpson, F., Dasmahapatra, K. K., Mallet, J., Davey, J. W., & Jiggins, C. D. (2015). Estimation of the spontaneous mutation rate in *Heliconius melpomene*. *Molecular Biology and Evolution*, 32(1), 239–243. <https://doi.org/10.1093/molbev/msu302>
- Keller, D., van Strien, M. J., Herrmann, M., Bolliger, J., Edwards, P. J., Ghazoul, J., & Holderegger, R. (2013). Is functional connectivity in common grasshopper species affected by fragmentation in an agricultural landscape? *Agriculture Ecosystems & Environment*, 175, 39–46. <https://doi.org/10.1016/j.agee.2013.05.006>
- Keyghobadi, N. (2007). The genetic implications of habitat fragmentation for animals. *Canadian Journal of Zoology*, 85(10), 1049–1064. <https://doi.org/10.1139/z07-095>
- Kirkpatrick, M., & Barton, N. H. (1997). Evolution of a species' range. *American Naturalist*, 150(1), 1–23.
- Kleijn, D., Biesmeijer, K. J. C., Klaassen, R. H. G., Oerlemans, N., Raemakers, I., Scheper, J., & Vet, L. E. M. (2020). Integrating biodiversity conservation in wider landscape management: Necessity, implementation and evaluation. In D. A. Bohan & A. J. Vanbergen (Eds.), *Future of agricultural landscapes, Pt 1* (Vol. 63, pp. 127–159). Academic Press.
- Kleijn, D., Bommarco, R., Fijen, T. P. M., Garibaldi, L. A., Potts, S. G., & van der Putten, W. H. (2019). Ecological intensification: Bridging the gap between science and practice. *Trends in Ecology & Evolution*, 34(2), 154–166. <https://doi.org/10.1016/j.tree.2018.11.002>
- Kleijn, D., Winfree, R., Bartomeus, I., Carvalheiro, L. G., Henry, M., Isaacs, R., Klein, A. M., Kremen, C., M'Gonigle, L. K., Rader, R., Ricketts, T. H., Williams, N. M., Lee Adamson, N., Ascher, J. S., Báldi, A., Batáry, P., Benjamin, F., Biesmeijer, J. C., Blitzer, E. J., ... Potts, S. G. (2015). Delivery of crop pollination services is an insufficient argument for wild pollinator conservation. *Nature Communications*, 6, 7414. <https://doi.org/10.1038/ncomms8414>
- Klein, A. M., Vaissiere, B. E., Cane, J. H., Steffan-Dewenter, I., Cunningham, S. A., Kremen, C., & Tscharntke, T. (2007).

- Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences*, 274(1608), 303–313. <https://doi.org/10.1098/rspb.2006.3721>
- Kling, M. M., & Ackerly, D. D. (2021). Global wind patterns shape genetic differentiation, asymmetric gene flow, and genetic diversity in trees. *Proceedings of the National Academy of Sciences of the United States of America*, 118(17), e2017317118. <https://doi.org/10.1073/pnas.2017317118>
- Kremen, C., Williams, N. M., Aizen, M. A., Gemmill-Herren, B., LeBuhn, G., Minckley, R., Packer, L., Potts, S. G., Roulston, T., Steffan-Dewenter, I., Vázquez, D. P., Winfree, R., Adams, L., Crone, E. E., Greenleaf, S. S., Keitt, T. H., Klein, A. M., Regetz, J., & Ricketts, T. H. (2007). Pollination and other ecosystem services produced by mobile organisms: A conceptual framework for the effects of land-use change. *Ecology Letters*, 10(4), 299–314. <https://doi.org/10.1111/j.1461-0248.2007.01018.x>
- Lecompte, E., Bouanani, M. A., Magro, A., & Crouau-Roy, B. (2016). Genetic diversity and structuring across the range of a widely distributed ladybird: Focus on rear-edge populations phenotypically divergent. *Ecology and Evolution*, 6(15), 5517–5529. <https://doi.org/10.1002/ece3.2288>
- Ligges, U., & Mächler, M. (2003). *scatterplot3d*—An R package for visualizing multivariate data. *Journal of Statistical Software*, 8(11), 1–20. <https://doi.org/10.18637/jss.v008.i11>
- Lira-Noriega, A., & Manthey, J. D. (2014). Relationship of genetic diversity and niche centrality: A survey and analysis. *Evolution*, 68(4), 1082–1093. <https://doi.org/10.1111/evo.12343>
- Marini, L., Bartomeus, I., Rader, R., & Lami, F. (2019). Species-habitat networks: A tool to improve landscape management for conservation. *Journal of Applied Ecology*, 56(4), 923–928. <https://doi.org/10.1111/1365-2664.13337>
- Martin, E. A., Dainese, M., Clough, Y., Baldi, A., Bommarco, R., Gagic, V., Garratt, M. P. D., Holzschuh, A., Kleijn, D., Kovács-Hostyánszki, A., Marini, L., Potts, S. G., Smith, H. G., Al Hassan, D., Albrecht, M., Andersson, G. K. S., Asís, J. D., Aviron, S., Balzan, M. V., ... Steffan-Dewenter, I. (2019). The interplay of landscape composition and configuration: New pathways to manage functional biodiversity and agroecosystem services across Europe. *Ecology Letters*, 22(7), 1083–1094. <https://doi.org/10.1111/ele.13265>
- New, T. R. (1991). *Insects as predators*. NSW University Press.
- Öckinger, E., & Smith, H. G. (2007a). Asymmetric dispersal and survival indicate population sources for grassland butterflies in agricultural landscapes. *Ecography*, 30(2), 288–298. <https://doi.org/10.1111/j.2007.0906-7590.05048.x>
- Öckinger, E., & Smith, H. G. (2007b). Semi-natural grasslands as population sources for pollinating insects in agricultural landscapes. *Journal of Applied Ecology*, 44(1), 50–59. <https://doi.org/10.1111/j.1365-2664.2006.01250.x>
- Ortego, J., Aguirre, M. P., Noguerales, V., & Cordero, P. J. (2015). Consequences of extensive habitat fragmentation in landscape-level patterns of genetic diversity and structure in the Mediterranean esparto grasshopper. *Evolutionary Applications*, 8(6), 621–632. <https://doi.org/10.1111/eva.12273>
- Ortego, J., Aparicio, J. M., Cordero, P. J., & Calabuig, G. (2008). Individual genetic diversity correlates with the size and spatial isolation of natal colonies in a bird metapopulation. *Proceedings of the Royal Society B: Biological Sciences*, 275(1646), 2039–2047.
- Ortego, J., Céspedes, V., Millán, A., & Green, A. J. (2021). Genomic data support multiple introductions and explosive demographic expansions in a highly invasive aquatic insect. *Molecular Ecology*, 30(17), 4189–4203. <https://doi.org/10.1111/mec.16050>
- Ortego, J., García-Navas, V., Noguerales, V., & Cordero, P. J. (2015). Discordant patterns of genetic and phenotypic differentiation in five grasshopper species codistributed across a microreserve network. *Molecular Ecology*, 24(23), 5796–5812. <https://doi.org/10.1111/mec.13426>
- Papadopoulou, A., & Knowles, L. L. (2015). Genomic tests of the species-pump hypothesis: Recent island connectivity cycles drive population divergence but not speciation in Caribbean crickets across the Virgin Islands. *Evolution*, 69(6), 1501–1517. <https://doi.org/10.1111/evo.12667>
- Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double digest RADseq: An inexpensive method for *de novo* SNP discovery and genotyping in model and non-model species. *PLoS One*, 7(5), e37135. <https://doi.org/10.1371/journal.pone.0037135>
- Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus genotype data. *Genetics*, 155(2), 945–959.
- R Core Team. (2023). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Rosenberg, N. A. (2004). DISTRUCT: A program for the graphical display of population structure. *Molecular Ecology Notes*, 4(1), 137–138. <https://doi.org/10.1046/j.1471-8286.2003.00566.x>
- Saccheri, I., Kuussaari, M., Kankare, M., Vikman, P., Fortelius, W., & Hanski, I. (1998). Inbreeding and extinction in a butterfly metapopulation. *Nature*, 392(6675), 491–494.
- Scheper, J., Holzschuh, A., Kuussaari, M., Potts, S. G., Rundlof, M., Smith, H. G., & Kleijn, D. (2013). Environmental factors driving the effectiveness of European agri-environmental measures in mitigating pollinator loss—A meta-analysis. *Ecology Letters*, 16(7), 912–920. <https://doi.org/10.1111/ele.12128>
- Schlaepfer, D. R., Braschler, B., Rusterholz, H. P., & Baur, B. (2018). Genetic effects of anthropogenic habitat fragmentation on remnant animal and plant populations: A meta-analysis. *Ecosphere*, 9(10), e02488. <https://doi.org/10.1002/ecs2.2488>
- Schtickzelle, N., Mennechez, G., & Baguette, M. (2006). Dispersal depression with habitat fragmentation in the bog fritillary butterfly. *Ecology*, 87(4), 1057–1065. [https://doi.org/10.1890/0012-9658\(2006\)87\[1057:ddwhf\]2.0.co;2](https://doi.org/10.1890/0012-9658(2006)87[1057:ddwhf]2.0.co;2)
- Silva, S. E., Seabra, S. G., Carvalheiro, L. G., Nunes, V. L., Marabuto, E., Mendes, R., Rodrigues, A. S. B., Pina-Martins, F., Yurtsever, S., Laurentino, T. G., Figueiredo, E., Rebelo, M. T., & Paulo, O. S. (2020). Population genomics of *Bombus terrestris* reveals high but unstructured genetic diversity in a potential glacial refugium. *Biological Journal of the Linnean Society*, 129(2), 259–272. <https://doi.org/10.1093/biolinnean/blz182>
- Spielman, D., Brook, B. W., & Frankham, R. (2004). Most species are not driven to extinction before genetic factors impact them.

Proceedings of the National Academy of Sciences of the United States of America, 101(42), 15261–15264.

- Thome, M. T. C., & Carstens, B. C. (2016). Phylogeographic model selection leads to insight into the evolutionary history of four-eyed frogs. *Proceedings of the National Academy of Sciences of the United States of America*, 113(29), 8010–8017. <https://doi.org/10.1073/pnas.1601064113>
- Thorbek, P., & Bilde, T. (2004). Reduced numbers of generalist arthropod predators after crop management. *Journal of Applied Ecology*, 41(3), 526–538. <https://doi.org/10.1111/j.0021-8901.2004.00913.x>
- Webster, M. T., Beaufrepaire, A., Neumann, P., & Stolle, E. (2023). Population genomics for insect conservation. *Annual Review of Animal Biosciences*, 11, 115–140. <https://doi.org/10.1146/annurev-animal-122221-075025>
- Weir, B. S., & Cockerham, C. C. (1984). Estimating *F*-statistics for the analysis of population structure. *Evolution*, 38(6), 1358–1370. <https://doi.org/10.1111/j.1558-5646.1984.tb05657.x>
- Willi, Y., Kristensen, T. N., Sgro, C. M., Weeks, A. R., Orsted, M., & Hoffmann, A. A. (2022). Conservation genetics as a management tool: The five best-supported paradigms to assist the management of threatened species. *Proceedings of*

the National Academy of Sciences of the United States of America, 119(1), e2105076119. <https://doi.org/10.1073/pnas.2105076119>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Ortego, J., Albrecht, M., Báldi, A., Bilde, T., Craig, S. B. B., Herrera, J. M., Hood, A. S. C., Kleijn, D., Maurer, C., Molina, F. P., Öckinger, E., Potts, S. G., Settepani, V., Thomsen, P. F., Trillo, M., Vajna, F., Velado-Alonso, E., & Bartomeus, I. (2024). Seminatural areas act as reservoirs of genetic diversity for crop pollinators and natural enemies across Europe. *Conservation Science and Practice*, 6(5), e13080. <https://doi.org/10.1111/csp2.13080>