# Thirty pyrenomycetoid Ascomycota species new to Norway from a workshop in Horten, May 2-5, 2023

Mathias Andreasen<sup>1</sup>, Roger Andersson<sup>2</sup>, Gernot Friebes<sup>3</sup>, Katrin Gilbert<sup>4</sup>, Edvin Johannesen<sup>5</sup>, Karl Soler Kinnerbäck<sup>6</sup>, Patrik Mlčoch<sup>7</sup>, Hermann Voglmayr<sup>8</sup>, Bjørn Wergen<sup>4</sup>, Björn Nordén<sup>1</sup> <sup>1</sup>Norsk institutt for naturforskning NINA, Sognsveien 68, 0855 Oslo, Norway. <sup>2</sup>Swedish University of Agricultural Sciences, Artsdatabanken, Almas allé 8E, Sweden <sup>3</sup>Natural History Museum, Joanneumsviertel, 8010 Graz, Austria <sup>4</sup>Schwarzwälder Pilzlehrschau, Werderstraße 17, 78132 Hornberg, Germany <sup>5</sup>Universitetet i Oslo, Naturhistorisk museum, Fungariet, Sars' gate 1, 0562 Oslo, Norway <sup>6</sup>Hagagatan 34F, 703 40 Örebro, Sweden <sup>7</sup>Department of Botany, Faculty of Science, Palacky University in Olomouc, Šlechtitelů 27, 783 71 Olomouc, The Czech Republic <sup>8</sup>Department of Botany and Biodiversity Research, University of Vienna, Rennweg 14, 1030 Vienna, Austria

Corresponding author: mathias.andreasen@nina.no

deres mikromorfologi, økologi og forekomst i Fennoskandia, noen også med bilder.

Norsk tittel: Tretti nye norske pyrenomycetoide sekksporesopp-arter fra workshop i Horten 2.-5. mai 2023.

Thirty pyrenomycetoid Ascomycota species new to Norway from a workshop in Horten, May 2-5, 2023. Agarica 2024 vol. 44: 3-23. DOI: 10.5617/agarica.11603

# **KEYWORDS**

Ascomycota, bark-inhabiting, woodinhabiting, plant-inhabiting, pyrenomycetes, pyreno-mycetoid fungi

## NØKKELORD

Sekksporesopper, barklevende, vedlevende, plantelevende, pyrenomyceter, pyrenomycetoide sopp

# SAMMENDRAG

Under en workshop med inviterte internasjonale eksperter fant vi 30 pyrenomycetoide sekksporesopp-arter nye for Norge, hvorav 16 nye for Skandinavia og en ny for Europa. Disse presenteres her med korte notater om

## ABSTRACT

During a workshop with invited international experts, we found 30 pyrenomycetoid ascomycete species new to Norway, whereof 16 new to Scandinavia and one new to Europe. These are presented here with short notes on their micromorphology, ecology, and occurrence in Fennoscandia, some also with photos.

# INTRODUCTION

How many species of pyrenomycetoid Ascomycota new to Norway can ten experts find in three days? Previous projects ("Artsprosjekt") in South Norway have recovered at least 215 non-lichenized or weakly lichenized pyrenomycetoid species new to the country (Jordal et al.2014; Nordén 2014; Nordén & Jordal 2014; Nordén et al. 2014; Lechat et al. 2015; Nordén et al. 2015a; Nordén et al. 2015b; Nordén 2016; Jordal et al. 2017; Nordén et al. 2017; Nordén et al. 2017; Nordén & Aptroot 2018; Nordén et al. 2019; Olsen et al. 2019; Nordén & Andreasen 2020; Nordén & Andreasen 2022; Jordal et al. 2022), mainly in temp-



Figure 1. Group photo from the workshop with the participants in front of an apple tree *Malus domestica* infested with mistletoe *Viscum album*. Three species new to Norway growing on mistletoe were found during the workshop. From the left: Mathias Andreasen, Patrik Mlčoch, Karl Soler Kinnerbäck, Gernot Friebes, Edvin Johannesen, Hermann Voglmayr, Roger Andersson, Bjørn Wergen, Katrin Gilbert and Björn Nordén. Photo Mathias Andreasen.

erate broadleaf forest. Due to the many findings, one could imagine that it would become more and more difficult to find new species for the country in this environment. To test this, a workshop was arranged in Horten, S Norway, May 2-5, 2023, inviting international experts to collecting trips during three days and microscopy lab in the evenings. Collection permits were obtained for collection in Nature Reserves.

Pyrenomycetoid Ascomycota is an artificial assemblage of Dothideomycetes, Eurotiomycetes (Chaetothyriomycetidae) and Sordariomycetes, collectively defined by having globose to flask-shaped ascomata with an apical opening. Many are saprotrophs on organic substrates such as wood and other plant material. Many are also lichen-forming, but the workshop concentrated on non-lichenized species, which are considerably less well known than their lichenized counterparts (the number of newly discovered species per biodiversity mapping project tends to be considerably lower in lichenized fungi, Table 2).

## MATERIALS AND METHODS

The presented species were encountered in temperate deciduous forests, wetlands and meadows on short excursions from Horten, Vestfold and Telemark county (Figure 1 & 2). The collection sites are situated below 75 m.a.s.l. and have a cool temperate climate, with the yearly average temperature of 7.5°C (warmest month July average maximum temperature of 23°C, coldest month February average maximum temperature of 2°C), and an average annual precipitation amounting to 952 mm. The bedrock in the area is rich in calcium and sport a high vascular plant diversity in herb-rich deciduous forests and wet-



Figure 2. Pyrenomycetoid Ascomycota collectors in action – experts from Europe gathered in Falkensten naturreservat in Horten, Vestfold og Telemark, May 2023. At this site fungi on twigs and branches appear to have been especially popular objects for study, but also fungi on herbs were studied during the workshop.

lands. Each forenoon and early afternoon were spent on the collection trips, and the rest of the time was spent in the lab. Seven of the reported findings were however collected at other sites in the days following the workshop by two of the participants.

The species were determined by studying the morphology of the sexual morph (teleomorph) or/and asexual morph (anamorph) under the microscope. Species distributions in Scandinavia were checked using the following sources: for Norway Artskart (2023), The Norwegian Mycological Database (Norsk Soppdatabase) NMD (2023), and Aarnæs (2002); for Sweden Eriksson (2014), Westberg et al. (2021) and Artportalen (2023), and for Denmark Atlas of Danish Fungi (2023), along with the international GBIF database

## (GBIF 2023).

Collectors, identifiers and photographers are designated by: BN: Björn Nordén, BW: Bjørn Wergen, EJ: Edvin Johannesen, HV: Hermann Voglmayr, KG: Katrin Gilbert, KSK: Karl Soler Kinnerbäck, MA: Mathias Andreasen, PM: Patrik Mlčoch, RA: Roger Andersson. The reported material will be deposited in Herb. O. Taxonomical nomenclature and systematic classification follows Mycobank.org unless otherwise stated, and should be treated as preliminary in several cases.

### RESULTS

Thirty species are reported as new to Norway (Table 1) and are presented below.

Table 1. Species new to Norway collected during or in immediate connection with the workshop. Known occurrences in neighboring Scandinavian countries are marked with  $(\times)$  in respective column.

Species new to Norway	Sweden	Denmark	Species new to Norway	Sweden	Denmark
Albicollum longisporum			Nemania chestersii	×	×
Anisogramma vepris	×	×	Neocucurbitaria populi		
Cucurbitaria chevalieri			Nitschkia confertula	×	×
Cucurbitaria salicina			Parafenestella pseudosalicis		
Delitschia intonsa	×		Parafenestella salicum		
Diatrypella angulata			Plagiostoma geranii	×	×
Digitodochium amoenum	×	×	Plenodomus visci		×
Hypoxylon julianii	×	×	Pseudohalonectria lignicola		
Lecythothecium duriligni			Saccardoella separans		
Lentithecium aquaticum			Septoriella elongata	×	
Leptosphaeria barriae			Setoseptoria arundinacea	×	×
Leptosphaeria clavispora			Sphaeropsis visci	×	×
Lophidium thyridioides			Splanchnonema loricatum	×	×
Lophodermium herbarum	×	×	Stylonectria wegeliniana	×	×
Massaria conspurcata	×	×	Synfenestella sorbi		
Melanconiella flavovirens	×				

## Albicollum longisporum Voglmayr, J.

Fourn., Tello & Jaklitsch (Sordariomycetes, Xylariales, Xylariaceae), Fig. 3

This species has immersed ascomata and a collar of white pseudostromatic tissue around the ostioles, which is characteristic for the genus Albicollum (Voglmayr et al. 2022). The most characteristic microscopic feature of this species is its helicoid germ slit that is less than 1 µm wide and coiling usually more than two times around the ascospore. Ascospore size of the Norwegian collection  $(43-51 \times 10.5-13)$ µm) fits the range reported for the species  $(21.5-67.5 \times 8.5-19 \,\mu\text{m}; \text{Voglmayr et al. 2022}).$ The Norwegian collection deviates from other collections of *A. longisporum* by the (almost) inamyloid (vs. hemiamyloid) apical ascus ring; however, sequence data confirm the species identification (Voglmayr, unpublished). It occurs on dead branches of various deciduous

shrubs and trees and has so far only been recorded from higher altitudes in the Mediterranean region (Greece, Morocco, Spain), which makes the Norwegian record a remarkable distribution range extension to the north.

Material: Vestfold og Telemark, Tinn, Gjøystdalvegen; EU89 UTM32V: 485108 6650156 (Lat/Long 8.73310 59.98846); on attached twigs of *Populus tremula*; May 9, 2023; leg. BW, KG; det. BW.

*Anisogramma vepris* (Lacroix) Merezhko (Sordariomycetes, Diaporthales, Valsaceae) Syn. *Apioporthe vepris* (Lacroix) Wehm.

This species forms small immersed or erumpent stromata and has 1-septate ascospores measuring  $6-9(10) \times 2-2.5 \ \mu\text{m}$ , often constricted at the septum, and with a short bristle-like mucous appendage to ca 5  $\mu$ m. It grows on dead *Rubus* stems.



Figure 3. *Albicollum longisporum*. Upper photos ascomata. Middle photos of ascospores. Lower photos asci and ascospores in asci, lower far right in lugol. Photos BW.

Material: Vestfold og Telemark, Horten, Falkensten naturreservat; EU89 UTM32V 578950 6589288 (Long/Lat 10.39183 59.43474); on *Rubus idaeus*, May 4 2023, Leg. MA, Det. KSK, MA.

Vestfold og Telemark, Horten, Borrevannet; EU89 UTM32V: 582023 6585510 (Lat/Long 10.44454 59,40024); on *Rubus idaeus*; May 3, 2023; leg. MA; det. KSK, MA.

*Cucurbitaria chevalieri* P.Karst. (Dothideomycetes, Pleosporales, Cucurbitariaceae)

Cucurbitaria chevalieri P.Karst., Fungi Fenn. Exs., Cent. 9: no. 892, 1869. ≡ *Teicho*spora chevalieri (P.Karst.) P.Karst. (as Chevalierii), Bidrag till Kännedom Finlands Natur Folk 23: 66, 1873. [Full synonymy, see https://www.mycobank.org/MB/847906] This species name was validly published on the label of Karsten, Fungi Fenn. Exs. 892, fig. 189 (FH 00785884). The species is characterized by single or clustered erumpent 400-500 µm broad black ascomata with indistinct papillae. The ascospores are light brown, with 3 to 5 transverse septa and one longitudinal septum except in the end cells,  $18-28 \times 8-10$ µm. On old dead wood. This is very likely no true Cucurbitaria (which has been restricted to only two species, only occurring on Berberis; see Jaklitsch et al. 2018). However, in lack of sequence data, the generic classification of the species cannot be evaluated.

Material: Vestfold og Telemark, Horten, Fjugstad naturreservat; EU89 UTM32V 582931 6581330 (Lat/Long 10.45892 59.36253); on *Ulmus glabra* (dead trunk); May 3, 2023; leg. MA; det. PM.

# Cucurbitaria salicina Fuckel (Dothideomy-

cetes, Pleosporales, Cucurbitariaceae), Fig. 4 *Cucurbitaria salicina* was described from *Salix fragilis* and *S. triandra*. It may not belong to *Cucurbitaria* s str. The material fits well with the original description. It forms gregarious ascomata on dead, often decorticated *Salix* twigs. The ascomata are partly embedded into the substrate, globose to somewhat flattened, black. Ascospores (18)18.6- $20.7(21.3) \times (7.9)8.5$ -9.4 µm, light brown to brown, with 5 irregular transverse septations and mostly only one longitudinal septation. Asci 105-125 × 12-13 µm, with 8 uniseriate spores.

See original description in Fuckel (1870). Material: Vestfold og Telemark, Tinn, Skinnarbu; EU89 UTM32V: 461354 6630988 (Lat/Long 8.31098 - 59.81481); on *Salix sp.;* May 7, 2023; leg. BW, KG; det. BW, HV.

*Delitschia intonsa* Luck-Allen (Dothideomycetes, Pleosporales, Sporormiaceae)

This species is fimicolous and appears on dung of various larger animals. Pseudothecia 700-950  $\times$  450-500 um, pyriform, membranous, thoroughly hairy, quite dark brown. Neck up to 200 x 150 µm, semi-coriaceous, black, tuberculate, with white hair. The ascospores are 1-septate, brown, and measure 46-54  $\times$ 14-18 µm. For a full description of the species, see Doveri (2004).

Material: Vestfold og Telemark, Horten, Falkensten naturreservat; EU89 UTM32V 578950 6589288 (Lat/Long 10.39183 59.43474); on deer dung; May 4, 2023; leg. BW, KG; det. BW.

# Diatrypella angulata (Fr.) Ces. & De Not.

Syn. *Diatrypella nigroannulata* (Grev.) Nitschke

This species grows on *Fagus* twigs and has furrowed perithecial necks, protruding from the initially light-colored cushion-shaped stromata. The most characteristic feature is a ring-shaped, dark zone sometimes visible through the bark around each stroma. It also differs from *D. favacea* with which it is synonymized by some authors in the more protruding and furrowed ostiolar necks. The sporebearing part of the ascus is over 60  $\mu$ m long and the ascospores are 5-7(8)  $\mu$ m long. For a



Figure 4. *Cucurbitaria salicina*. Upper photo: ascospores and asci. Lower left photo: cross section of peridium and ascomata. Lower right photos: ascomata on substrate and cross section of ascomata. Photos BW.

full description, se Vasilyeva & Scheuer (1996).

Material: Vestfold og Telemark, Horten, Karljohansvern plante- og dyrefredningsområde, EU89 UTM32V: 584822 6587947 (Lat/Long 10.49478 59.42156); on twig of *Fagus sylvatica*; May 4, 2023; leg. BN; det. HV.

*Digitodochium amoenum* (Nitschke ex Sacc.) Voglmayr, J.Fourn. & Jaklitsch (Sordariomycetes, Xylariales, Xylariaceae)

Syn. Anthostoma amoenum Nitschke ex Sacc.

Digitodochium can be diagnosed by Lopadostoma-like pseudostromata overlain by a white disc and ascospores with a lateral germination pore, 3-4  $\mu$ m diam, conspicuous, usually central. Ascospores (20)21-29.6(33.5) × (8.5)9.2-13(14)  $\mu$ m, aseptate, ellipsoid to narrowly ellipsoid or subcitriform, equilateral, dark brown, with narrowly rounded, frequently slightly pinched ends. The anamorph is highly diagnostic by subhyaline staurosporous conidia consisting of a cylindrical main axis with whorls of lateral branches arising at wide angles from the basal cells of the main axis; conidial masses are produced in yellow-orange acervular conidiomata. The anamorph has been previously recorded from Europe as Digitodochium rhodoleucum, which, however, is a distinct species only known from Japan. This is the first time a holomorphic collection is recorded from Norway; an anamorphic Norwegian collection has already been recorded in Voglmayr et al. (2022) and the species is therefore not included in the sum of species new to Norway. For a full description of the species, see Voglmayr et al. (2022). This species seems to be confined to corticated dead twigs of Fagus sylvatica, and the anamorph-

telomorph connection has only recently been clarified (Voglmayr et al. 2022).

Material: Vestfold og Telemark, Horten, Karljohansvern plante- og dyrefredningsområde, EU89 UTM32V: 584822 6587947 (Lat/Long 10.49478 59.42156); on *Fagus sylvatica*; May 4, 2023; leg. HV; det. HV.

*Hypoxylon julianii* L.E. Petrini (Sordariomycetes, Xylariales, Hypoxylaceae)

This seldom collected species is quite similar to the common *H. rubiginosum* but has smaller perithecia and significantly larger ascospores. The perithecia are spherical,  $(200-)250-320 \ \mu\text{m}$  in diameter and the ascospores measure  $15-20.4 \times 6.2$ -8.4  $\mu\text{m}$ .

Detailed description, see Pyrenomycetes from southwestern France (2004).

Material: Viken, Asker, Vettre; EU89 UTM32V: 582744 6633639 (Lat/Long 10.47607 59.83214); on *Ulmus glabra* (branch on the ground); May 2, 2023; leg. RA; det. RA, conf. BN.

*Lecythothecium duriligni* Réblová & Winka (Sordariomycetes, Chaetosphaeriales, Chaetosphaeriaceae), Fig. 5

This species has embedded single perithecia which grow on wood of various deciduous and very rarely also on coniferous trees. It has unitunicate asci with an inamyloid apical ring, and characteristic spores that are fusiform, up to around 40  $\mu$ m long, 7-septate at maturity, with brown central cells and hyaline end cells, measuring in the studied material (34)35.5-40(41) × (8)8.7-9(9.3)  $\mu$ m.

For a full description of the species, see Reblova & Winka (2001).

Material: Viken, Asker, Spireodden naturreservat; EU89 UTM32V: 583872 6633541 (Lat/Long 10.49615 59.83104); on *Malus domestica*; May 2, 2023; leg. RA; det. RA, BN. *Lentithecium aquaticum* Ying Zhang, J. Fourn. & K.D. Hyde (Dothideomycetes, Pleosporales, Lentitheciaceae)

This species has scattered, immersed ascomata that are lenticular with a flattened base. The rounded papilla protrudes slightly above the wood surface which is stained dark grey. The ascospores measure  $25-30 \times 8-12 \mu m$  and are broadly fusiform with broadly rounded ends, inequilateral in side view, hyaline, twocelled, turning four-celled with age, and are encompassed by a conspicuous sheath, 1.5-3  $\mu m$  thick. For a full description of the species, see Zhang et al. (2008).

Material: Vestfold og Telemark, Horten, Fjugstad naturreservat; EU89 UTM32V 582931 6581330 (Lat/Long 10.45892 59.36253); on *Phragmites australis*; May 3 2023; leg. BW, KG; det. BW.

*Leptosphaeria barriae* Shoemaker (Dothideomycetes, Pleosporales, Leptosphaeriaceae)

This species has small, immersed black ascomata and 6-septate ascospores that measure about  $45-55 \times 5-6 \mu m$ , and grows on old stems of *Lupinus polyphyllus*. Shoemaker (1984) also described several similar species from this host substrate, e.g. *Leptosphaeria castrensis* and *Leptosphaeria wehmeyeri*, with different ascospore features. New phylogenetic analyses show that this species belongs to family Phaeosphaeriaceae (Mlčoch P., unpublished). For a full description, see Shoemaker (1984).

Material: Vestfold og Telemark, Horten, Fjugstad naturreservat; EU89 UTM32V 582931 6581330 (Lat/Long 10.45892 59.36253); on *Lupinus polyphyllus*; May 3, 2023; leg. PM; det. PM, conf. BN.

*Leptosphaeria clavispora* J.H. Mill. & Burton (Dothideomycetes, Pleosporales, Leptosphaeriaceae)

This species has subepidermal, globose pseudothecia with a short papilla. The asco-



Figure 5. *Lecythothecium duriligni*. Upper and lower right photos of ascospores. Middle and lower left photos of ascomata on substrate and cross section of ascomata. Photos BW.

spores are clavate to clavate-fusiform, yellowish-brown, 8- to 10-septate, with third and fourth cells from apex enlarged, constricted at first septum,  $45-65 \times 6.5-8$  um. It grows on herbs of the family Asteraceae. According to new molecular analyses this species belongs to the genus *Ophiobolus* (Mlčoch P., unpublished). Se original description in Miller & Burton (1942)

Material: Viken, Asker, Spireodden naturreservat; EU89 UTM32V: 583872 6633541 (Lat/Long 10.49615 59.83104); on *Centaurea jacea*; May 2, 2023; leg. EJ; det. PM, conf. BN.

*Leptosphaeria modesta* Rabenh. (Dothideomycetes, Pleosporales, Leptosphaeriaceae) Syn. *Nodulosphaeria modesta* (Rabenh.) Munk ex L. Holm

This species has small blacksubcuticular

ascomata and can grow on herbaceous stems of various herbs and the species . It is further characterized by fusoid, hyaline to yellowish, 4- to 5-septate ascospores, 30-36  $\mu$ m long with 2nd cell enlarged. Polar appendages are often present. For detailed description, see Italian microfungi (2021).

Material: Vestfold og Telemark, Horten, Borrevannet; EU89 UTM32V: 582023 6585510 (Lat/Long 10.44454 59.40024); on herbal stem; May 3, 2023; leg. PM; det. PM.

*Lophidium thyridioides* Sacc. & Speg. (Dothideomycetes, Pleosporales, Massariaceae), Fig. 6

Syn. Decaisnella thyridioides (Sacc. & Speg.) M.E.Barr

This species has embedded shiny black ascomata of about 0,7-1 mm in diam, which break through the substrate with an acute

perithecial opening. The muriform ascospores in the studied material measured (31.5)33.5- $38.9(41.5) \times (11.3)12$ - $13.6(14.3) \mu m$  and had 10-12 transverse and 1-3(4) longitudinal septa. The similar *Decaisnella mesascium* has larger spores (>50  $\mu m$  length). For a full description, see Barr (1990).

Material: Vestfold og Telemark, Tinn, Austbygda; EU89 UTM32V: 461354 6630988 (Lat/Long 8.31098 59.81481); on wood of *Salix* sp., May 7, 2023; leg. BW, KG; det. BW.

## *Lophodermium herbarum* (Fr.) Fuckel (Leotiomycetes, Rhytismatales, Rhytismataceae)

This species occurs on dead leaves of *Convallaria majalis*. The ascomata are hystero-thecial. The asci are cylindrical and 8-spored. Ascospores are hyaline, filiform, and measure ca  $55 \times 2$  um. Se original description in Fuckel (1870).

Material: Vestfold og Telemark, Horten, Falkensten naturreservat; EU89 UTM32V 578950 6589288 (Lat/Long 10.39183 59.43474); on *Convallaria majalis*; May 4, 2023; leg. PM; det. PM.

*Massaria conspurcata* Sacc. (Dothideomycetes, Pleosporales, Massariaceae)

This species has immersed black depressed subglobose ascomata 0.6-1 mm diam forming under the bark surrounded by a black stromatic zone. It is well characterized by fourspored asci containing brown narrow ascospores (<16  $\mu$ m wide) with usually paler brown central cells and an occurrence on dead corticated twigs of *Prunus padus* (Voglmayr & Jaklitsch 2011). It is probably not rare, but likely overlooked due to its highly seasonal occurrence in early spring; the material collected was already largely



Figure 6. *Lophidium thyridioides*. Upper left photo of spores. Upper right photo of ascomata on substrate and cross section of ascomata. Lower photos of ascospores in asci. Photos BW.

overmature. For a full description of the species, see Voglmayr & Jaklitsch (2011).

Material: Vestfold og Telemark, Horten, Karljohansvern plante- og dyrefredningsområde; EU89 UTM32V 578822 6587947 (Lat/Long 10.38908 59.42273); on *Prunus padus*; May 4, 2023; leg. HV; det. HV.

*Melanconiella flavovirens* (G.H.Otth) Voglmayr & Jaklitsch (Sordariomycetes, Diaporthales, Melanconiellaceae)

This species forms circular pseudostromata with 1-15(25) ostioles and an ectostroma which is often yellow-green to olive. The spore ends have up to 1.4  $\mu$ m long, distinct, knob-like hyaline appendages (Voglmayr et al. 2012). The ascospores are hyaline, ellipsoid or fusoid, 1-septate with no or slight constriction, have one large and many small guttules in each cell and measure (17)20.5-24.5(30) × (7.0)7.5-8.8(9.8)  $\mu$ m. It occurs on dead corticated twigs and branches of *Corylus avellana*.

Material: Vestfold og Telemark, Horten, Falkensten naturreservat; EU89 UTM32V 578950 6589288 (Lat/Long 10.39183 59.43474); on *Corylus avellana*; May 4, 2023; leg. KSK; det. HV.

*Nemania chestersii* (J.D.Rogers & Whalley) Pouzar (Sordariomycetes, Xylariales, Xylariaceae)

This species has up to 6 cm long, effused, grey to brown stromata with abrupt margins on wood. Ascus annulus  $3 \times 2.5 \,\mu$ m, blue in Lugol's and Melzer's solutions. Ascospores (12)13-17 × 4.5-5.5(6)  $\mu$ m, ellipsoid to navicular or lunate, with narrowly rounded ends, guttulate, brown and without germ slit. The spore wall is characteristically ornamented with around 20 longitudinal ribs which separates this species from other species in the genus. It occurs on decorticated, decayed wood of *Fagus*, *Fraxinus* and *Quercus*. For a full description of the species, see Granmo et

al. 1999.

Material: Vestfold og Telemark, Horten, Karljohansvern plante- og dyrefredningsområde; EU89 UTM32V 578822 6587947 (Lat/Long 10.38908 59.42273); on *Fagus sylvatica*; May 4, 2023: leg. KSK: det. KSK.

*Neocucurbitaria populi* Jaklitsch & Voglmayr (Dothideomycetes, Pleosporales, Cucurbitariaceae), Fig. 7

This species grows on attached twigs of *Populus* sp., the black ascomata often somewhat embedded in remnants of bast tissues. Ascospores muriforme  $(21.9)22.3-25.8 \times (9.7)9.9-11(11.4) \mu m$ , 7 somewhat irregular transverse septations and mostly 1-2 longitudinal septations, dark brown. For a full description, see Jacklitsch et al. (2017).

Material: Vestfold og Telemark, Rjukan, near Vemork, on attached twigs of *Populus* sp.; May 07, 2023; leg. BW, KG; det. BW.

*Nitschkia confertula* (Schwein.) Nannf. (Sordariomycetes, Sordariales, Nitschkiaceae)

The ascomata of this species are seated on a thin black subiculum, are turbinate to pulvinate, and become shallowly to deeply cupulate when dry. The spore-bearing part of the asci measure  $25-35 \times 9-15$  µm and the asci taper into a long, thread-like stalk to ca 30 µm in length. The ascospores measure (7.5)8-10 × 3-4.5 µm and are ovoid to ellipsoidal, occasionally more narrowed toward one end, frequently flattened on one side or subreniform, frequently biguttulate and then sometimes appearing pseudoseptate, initially smoky-grey but becoming golden brown, in mass yellowish-brown, smooth. Original description in Nannfeldt (1975) and a detailed description in Fungi of Great Britain and Ireland (2023).

Material: Vestfold og Telemark, Horten, Fjugstad naturreservat; EU89 UTM32V 582931 6581330 (Lat/Long 10.45892 59.36253); on *Lupinus polyphyllus*; May 3,



Figure 7. Neocucurbitaria populi. Upper and lower left: ascospores in asci. Lower right: ascomata on substrate and cross section of ascomata. Photos BW.

2023; leg. EJ; det. BW.

**Parafenestella pseudosalicis** Jaklitsch & Voglmayr (Dothideomycetes, Pleosporales, Cucurbitariaceae), Fig. 8

There are a few closely related species of *Parafenestella* occurring on dead twigs of *Salix*, often on or near old fruitbodies of *Valsa* sp./*Cytospora* sp. They are difficult to separate without sequence data but we here tentatively report two species based on morphological analyses. *P. pseudosalicis* forms scattered ascomata mostly on twigs where the bark has partly degraded or on remnants of bast tissues, very probably on or at least near other pyrenomycetes, likely *Cytospora* sp.

Ascospores muriforme (25.5)26.2-29.5(31.1)  $\times$  (12.1)12.5-13.5(13.8) µm, dark yellowish brown to dark brown, with 7-10 transverse

and quite irregular three longitudinal septa. For a full description, see Jacklitsch & Voglmayr et al. (2019).

Material: Vestfold og Telemark, Tinn, Skinnarbu; EU89 UTM32V: 461354 6630988 (Lat/Long 8.31098 59.81481); on *Salix* sp.; May 7, 2023; leg. BW, KG; det. BW.

*Parafenestella salicum* Jaklitsch & Voglmayr (Dothideomycetes, Pleosporales, Cucurbitariaceae), Fig. 9

The small ascomata of this species (about 0.4-0.5 mm in diameter) are black, almost globose, and embedded into the substrate (bark and bast tissues). The muriforme ascospores measure  $(31)31.4-34.5(37) \times 13.5-15 \ \mu m$  and are yellowish brown, with 10-15 transverse and 3-4 irregular longitudinal septations. For a full description, see Jacklitsch & Voglmayr et al. (2019).



Figure 8. *Parafenestella pseudosalicis*. Upper left photo of ascomata on substrate. Upper right photo of ascospores. Middle left and lower photos of asci with ascospores. Photos BW.

Material: Vestfold og Telemark, Tinn, Austbygda; EU89 UTM32V: 461354 6630988 (Lat/Long 8.31098 59.81481); on *Salix sp.* on dead *Leucostoma sp.*; May 7, 2023; leg. BW, KG; det. BW.

*Plagiostoma geranii* (Hollós) Sogonov (Sordariomycetes, Diaporthales, Gnomoniaceae)

This species grows on different species of *Geranium*. The perithecia are oblate, sometimes strongly flattened, immersed in the substrate and with a protruding neck which may be curved or bent. Asci ca  $30-40 \times 6-10$  µm, with a refactive apical ring to ca 2 µm diam. Ascospores ca  $13-18 \times 1.5-2.5$  µm, narrowly fusiform, straight or curved, hardly constricted at the ±central septum, hyaline, thin-walled, the cells containing numerous oil droplets, with a short filiform appendage at one or other, or both ends. For details, see

Monod (1983).

Material: Viken, Asker, Spireodden naturreservat; EU89 UTM32V: 583872 6633541 (Lat/Long 10.49615 59.83104); on *Geranium sanguineum*; May 2, 2023; leg. BN, Det. BN & Viken, Asker, Spireodden naturreservat; EU89 UTM32V: 583872 6633541 (Lat/Long 10.49615 59.83104); on *Geranium sanguineum*; May 2, 2023; leg. MA; det. MA.

**Plenodomus visci** (Sacc.) Gruyter, Aveskamp & Verkley (Dothideomycetes, Pleosporales, Leptosphaeriaceae)

Syn. Gibberidea visci Fuckel

This species grows on mistletoe, *Viscum album*. It has subepidermal, later erumpent and densely crowded, black pseudothecia with a shortly papillate ostiole. Asci bitunicate, apically with a small ocular chamber. Ascospores (5-)7-septate, cylindrical to slightly



Figure 9. *Parafenestella salicum*. Upper and lower left photos of ascospores in asci. Middle left photo of ascospores. Lower right photo of ascomata on substrate and cross section of ascomata. Photos BW.

clavate, constricted at the middle, ca 30-40  $\times$  7-8  $\mu m.$  For details, see Holm (1968) and Crous et al. (2020).

Material: Vestfold og Telemark, Horten, Borrevannet; EU89 UTM32V: 582023 6585510 (Lat/Long 10.44454 59.40024); on *Viscum album* on *Tilia cordata;* May 3, 2023; leg. BN; det. BN.

*Pseudohalonectria lignicola* Minoura & T.Muroi (Sordariomycetes, Pseudohalonec-triaceae)

This species occurs in dead wood in freshwater. It is one of the species in the genus with a long perithecial neck. Ascospores up to ca  $70 \times 7 \mu m$ , weakly pigmented, yellow in mass, with 5-7(11) septa. For a detailed description, see Shearer (1989).

Material: Vestfold og Telemark, Horten, Falkensten naturreservat; EU89 UTM32V 578950 6589288 (Lat/Long 10.44454 59,40024); on deciduous wood in water (ditch); May 4, 2023; leg. RA; det. RA, BN.

*Saccardoella separans* M.E. Barr (Sordariomycetes incertae sedis), Fig. 10

This species has small, scattered ascomata fully embedded into the substrate, clypeate, in section often higher than broad. The ascospores measure (59.8)62.8-67.8(79) × (7.5)7.6-8.8(8.9)  $\mu$ m, hyaline, with 19-22 transverse distoseptata, one of the middle cells is remarkably larger than the others, all filled with at least one large oil drop. This species was previously reported only from North America. For a full description, see Barr (1994).

Material: Vestfold og Telemark, Tinn, Hovin Kirke; EU89 UTM32V: 500374 6634643 (Lat/Long 9.00667 59.84943); on branch of *Populus tremula*; May 9, 2023; leg. BW, KG; det. BW.



Figure 10. Saccardoella separans. Upper photos of ascomata on substrate and in cross section. Middle and lower photos of ascospores. Photos BW.

*Septoriella elongata* (Wehm.) Y. Marín & Crous (Dothideomycetes, Pleosporales, Phaeosphaeriaceae)

This species has immersed, globose ascomata. The ascospores are narrowly fusiform, straight or slightly curved, reddish-brown, 10-septate and measure  $44-52 \times 6-7(8) \mu m$ . The fourth cell from the top is enlarged (widened) and slightly longer than adjacent cells. The spores also have a conspicuous sheath which is 2-3  $\mu m$  wide. It is recorded on a variety of grasses. For a full description, see Shoemaker & Babcock (1989) and Felix et al. 2019).

Material: Vestfold og Telemark, Horten, Fjugstad naturreservat; EU89 UTM32V 582931 6581330 (Lat/Long 10.45892 59.36253); on *Phragmites australis*; May 3, 2023; leg. KSK; det. KSK. *Setoseptoria arundinacea* (Sowerby) Kaz. Tanaka & K. Hiray. (Dothideomycetes, Pleosporales Lentitheciaceae)

This species has immersed ascomata and occurs on culms of *Phragmites australis*. The ascospores measure 23-40.5  $\times$  3.5-6 µm and are 1- to 2-septate, hyaline and have a uniform sheath. The material fits the description in Tanaka et al. (2015). *S. arundinacea* may consist of several cryptic species with close morphological resemblance. Another slightly differing collection of a similar species was collected by MA in 2021 and the entry needs to be changed into genus level.

Material: Vestfold og Telemark, Horten, Borrevannet; EU89 UTM32V: 582023 6585510 (Lat/Long 10.44454 59.40024); on *Phragmites australis*; May 3, 2023; leg. KSK; det. KSK.

*Sphaeropsis visci* (Alb. & Schwein.) Sacc. (Dothideomycetes, Botryosphaeriales, Botryosphaeriaceae)

This coelomycete seems to be relatively common on mistletoe *Viscum album*. The black pycnidia occur on leaves and stems and produce oval conidia  $34-43 \times 14-17 \mu m$ (x = 40.1 × 16.7  $\mu m$ , n = 10) with obtuse to rounded apex, obtuse or truncate base. They are moderately thick-walled, initially hyaline, becoming brown, externally smooth-walled and internally finely vertuculose. For a detailed description, see Phillips et al. 2013.

Material: Vestfold og Telemark, Horten, Borrevannet; EU89 UTM32V: 582023 6585510 (Lat/Long 10.44454 59.40024); on *Viscum album*; May 3, 2023; leg. BN, KG; det. BN.

*Splanchnonema loricatum* (Tul. & C.Tul.) M.E.Barr (Dothideomycetes, Pleosporales, Pleomassariaceae)

This species occurs on corticated twigs of *Fagus sylvatica* primarily in the winter season. It is characterised by pseudothecia embedded in bark and by octosporous cylindrical-clavate asci containing three-celled dark brown ascospores of size  $30-50 \times 11-18 \mu m$  that are asymmetric and surrounded by a thick swelling gelatinous sheath. For a full description, see Barr (1982).

Material: Vestfold og Telemark, Horten, Karljohansvern plante- og dyrefredningsområde; EU89 UTM32V 578822 6587947 (Lat/Long 10.38908 59.42273); on *Fagus sylvatica*; May 4, 2023; leg. HV; det. HV.

*Stylonectria wegeliniana* (Rehm) Gräfenhan, Voglmayr & Jaklitsch (Sordariomycetes, Hypocreales Nectriaceae)

This species has semi-immersed orangered to red perithecia and is found on dead twigs of *Ulmus* spp., where it is associated with the pyrenomycete *Hapalocystis bicaudata*, a species that in Norway was first found in 2014 (Nordén et al. 2015). *S. wegeliniana* resembles *S. norvegica* but has larger, verruculose ascospores. For a full description, see Gräfenhan et al. (2011).

Material: Vestfold og Telemark, Horten, Falkensten naturreservat; EU89 UTM32V 578950 6589288 (Lat/Long 10.44454 59,40024); on *Hapalocystis bicaudata/ Ulmus glabra*; May 4, 2023: leg. RA; det. RA.

*Synfenestella sorbi* (P.Karst.) Jaklitsch & Voglmayr (Dothideomycetes, Pleosporales, Cucurbitariaceae), Fig. 11

This species occurs on dead twigs of *Sorbus aucuparia*, where it typically grows in old leucostoma-type fruitbodies of *Valsa* sp. (*Leucostoma* sp.). It has been recorded from Europe and North America so far, and probably the record from Norway is the most northern find in Europe.

Ascospores muriforme (26.6)-29.5(32.2)  $\times$  (12.5)13.9-15.5(16.5) µm, yellowish brown, up to 7 transverse and 2-3 longitudinal septa; Asci 200-220 x 18-20 µm, 4-8 spored. For a full description of the species, see Jaklitsch & Voglmayr (2020).

Material: Vestfold og Telemark, Tinn, Austbygda; EU89 UTM32V: 461354 6630988 (Lat/Long 8.31098 59.81481); on *Sorbus aucuparia* on dead *Leucostoma* sp.; May 7, 2023; leg. BW, KG; det. BW.

# DISCUSSION

Ascomycota is the most abundant and diverse fungal phylum, but its members are still poorly known in Norway. There are new country records of Ascomycota species, and even new species to science, waiting to be discovered in every vegetated habitat. The problem is to find and to identify them, something that requires training and experience.

However, this is not an insurmountable problem, and the question is why so few mycologists are interested in pyrenomycetoid Ascomycota. This group of fungi is very diverse



Figure 11. *Synfenestella sorbi*. Upper and middle left photos: ascomata on substrate and in cross section. Middle center, middle right and lower photos: ascospores and asci. Photos BW.

and present great morphological variation, and the species are easy to collect and store (you just dry them in room temperature, and this doesn't cause them to change).

Compared to other non-lichenized and non lichen associated taxa, the knowledge of pyrenomycetoid fungi is very poor and for instance lichenized Ascomycota are much better known. An illustration of this is that biodiversity mapping projects on lichenized Ascomycota found significantly fewer new species to Norway in comparison (Table 2). We therefore hope that more professional and amateur mycologists will take part in future surveys of poorly studied taxa of non-lichenized Ascomycota.

In addition to the species presented here, at least six species were found to be new to science and these will be taxonomically described by us or other mycologists. It seems clear that not enough attention has been paid to pyrenomycetoid Ascomycota in Southern Norway despite several successful projects. The result from our workshop was similar to previous workshops supporting the need for additional mapping of these megadiverse and interesting fungi.

### ACKNOWLEDGMENTS

We thank the Norwegian Biodiversity Information Centre (Artsdatabanken) for funding the workshop. The Natural History Museum, Oslo, is gratefully acknowledged for the loan of several microscopes. John-Bjarne Jordal is thanked for useful comments on a previous version of the article.

Table 2. Number of species new to Norway identified by completed biodiversity mapping projects on fungi and bryophytes funded by the Norwegian Biodiversity Information Centre (Artsdatabanken). Information on the success of previous fungal and other cryptogam biodiversity mapping projects was gathered from https://www.artsdatabanken.no/Pages/199779/Sopp\_og\_lav) and https://www.artsdatabanken.no/Pages/199785/Moser\_og\_pseudosopper.

Taxon/guild	Proj.	Years	Number of	Comment
	nb.		species new to	
			Norway	
Bitunicate	32-17	2018-2021	100	
Ascomycota on				
bark and wood				
Sordariomycetes	58-12	2013-2015	89	
in broadleaf forest				
Endophytic fungi	23-19	2018-2021	62	
in sapwood			-	
Cortinarius	63-10	2011-2013	60	
Jelly fungi	44-15	2016-2018	57	
Ascomycetes in	56-09	2010-2013	57	Finnmark province
Finnmark	50 05	2010 2015	51	r minimum province
Bryophilous fungi	42-13	2014-2016	32	
Entoloma	50-14	2015-2017	30	
Hypogeous fungi	67-10	2013 2017	29	No exact figure given but
Trypogeous tungi	07-10	2011-2013	2)	less than 30
Lichenized fungi	34-16	2017-2020	20	Rain forest
and their parasites	54-10	2017-2020	29	Kalli lolest
Cortigioid fungi	50.12	2013 2016	25	
conticiola lag	39-12	2013-2010	23	
L acidacida	46.14	2015 2016	22	Of these 18 undescribed
skorpelay - Fase	40-14	2013-2010	25	of these is undescribed
III				
Ramaria	50.00	2010-2012	22	
Hohvalla	45 15	2016-2012	22	
Onhiostoma	51 14	2010-2018	20	
Aluina humanhataa	22.10	2013-2017	14	
Alpine bryophytes	22-19	2019-2022	12	
Bryophytes in	61-10	2011-2013	0	
poorly studied				
nature types	55.00	2010 2012		
Lecideoide	55-09,	2010-2013	5	
lichenized fungi I-	57-12			
	10.10	0010 0000		
<i>Cortinarius</i> on	18-18	2019-2020	4	Svalbard
Svalbard	(1.12	0010 0011		
Levermoser på	61-12	2013-2016	3	
Vestlandet				
Archeorhizo-	4-20	2020-2023	0	Genetic variants found, but
mycetes				no new species described.
Thelephorales	54-09	2010-2013	0	

REFERENCES

- Aarnæs J-O, 2002. Catalogue over macro- and micromycetes recorded for Norway and Svalbard. Fungiflora A/S, Oslo, 412 pp.
- Artportalen, 2023. The Swedish Species Observation System. https://www.artportalen.se/. Accessed September 10, 2023.
- Artskart, 2023. Artsdatabanken, Species Map Service 1.6. https://artskart.artsdatabanken.no/. Accessed September 10, 2023.
- Atlas of Danish Fungi, 2023. Svampeatlas 2.0. https://svampe.databasen.org/. Accessed September 10, 2023.
- Barr ME, 1982. On the Pleomassariaceae (Pleosporales) in North America. Mycotaxon 15: 349-383.
- Barr ME, 1990. North American Flora. Series II.: Part 13: Melanommatales (Loculoascomycetes): 81. New York Botanical Garden, USA.
- Barr ME, 1994. Notes on the Amphisphaeriaceae and related families. Mycotaxon 51: 191-224.
- Crous PW, Wingfield MJ, Schumacher RK, Akulov A, Bulgakov TS, Carnegie AJ, Jurjević Ž, Decock C, Denman S, Lombard L, Lawrence DP, Stack AJ, Gordon TR, Bostock RM, Burgess T, Summerell BA, Taylor PWJ, Edwards J, Hou LW, Cai L, Rossman AY, Wöhner T, Allen WC, Castlebury LA, Visagie CM, Groenewald JZ, 2020. New and interesting fungi. 3. Fungal Systematics and Evolution 6: 157-231.
- Doveri F, 2004. Fungi Fimicoli Italici. A guide to the recognition of Basidiomycetes and Ascomycetes living on feacal material. A.M.B.-Fondazione Centro Studi Micologici, Vicenza.
- Eriksson OE, 2014. Checklist of the nonlichenized ascomycetes of Sweden. Symbolae Botanicae Upsalienses 36: 1-501.
- Marin-Felix Y, Hernández-Restrepo M, Iturrieta-González I, García D, Gené J, Groenewald JZ, Cai L, Chen Q, Quaedvlieg W, Schumacher RK, Taylor PWJ, Ambers C, Bonthond G, Edwards J, Krueger-Hadfield SA, Luangsa-Ard JJ, Morton L, Moslemi S, Sandoval-Denis M, Tan YPc, Thangavel R, Vaghefi N, Cheewangkoon R, Crous PW, 2019. Genera

of phytopathogenic fungi: GOPHY 3. Study in Mycology 94: 109.

- Fuckel KWGL, 1870. Jahrbücher des Nassauischen Vereins für Naturkunde. 23-24: 172.
- Fungi of Great Britain and Ireland, 2023. Fungi of Great Britain and Ireland. https://fungi.myspecies.info/. Accessed September 10,2023.
- GBIF: The Global Biodiversity Information Facility, 2023. https://www.gbif.org/.
- Granmo A, Læssøe T, Schumacher T, 1999. The genus Nemania s.l. (Xylariaceae) in Norden. Sommerfeltia 27(96): 1-99.
- Gräfenhan T, Schroers H-J, Nirenberg HI & Seifert KA, 2011. An overview of the taxonomy, phylogeny, and typification of nectriaceous fungi in Cosmospora, Acremonium, Fusarium, Stilbella, and Volutella. Studies in Mycology 68: 79-113.
- Holm L, 1968. Taxonomic notes on Ascomycetes. VI. On the genus Gibberidea Fuck. and some alleged relatives. Svensk Botanisk Tidskrift 62: 217-247.
- Italian microfungi, 2021. https://italianmicrofungi. org/pleosporales/leptosphaeriaceae/leptosph aeria/leptosphaeria-modesta-.html. Accessed April 16, 2024. Wijesinghe SN, Camporesi E, Wanasinghe DN, Maharachchikumbura SSN, Senanayake IC, Phookamsak R, Hongsanan S, Tibpromma S, Thambugala KM, Luangharn T, McKenzie EHC, Hyde KD 2021. Asian Journal of Mycology 4: 10-18
- Jaklitsch WM, Checa J, Blanco MN, Olariaga I, Tello S & Voglmayr H, 2017. A preliminary account of the Cucurbitariaceae. Studies in Mycology. 90.
- Jaklitsch WM, Checa J, Blanco MN, Olariaga I, Tello S, Voglmayr H, 2018. A preliminary account of the Cucurbitariaceae. Studies in Mycology 90: 71-118.
- Jaklitsch WM & Voglmayr H, 2019. Fenestelloid clades of the Cucurbitariaceae. Persoonia -Molecular Phylogeny and Evolution of Fungi. 44.
- Jaklitsch WM, Voglmayr H, 2020. Fenestelloid clades of the Cucurbitariaceae. Persoonia 44: 1-40.

- Jordal, JB, Holien H, Nordén B, 2022. Melaspilea bagliettoana new to Fennoscandia. Graphis Scripta 34: 1-6.
- Jordal JB, Klepsland JT, Nordén B, 2017. Melaspilea lentiginosula, a species of oceanic pine forests, new to Fennoscandia. Graphis Scripta 29: 33-39.
- Jordal JB, Nordén B, Gaarder G, 2014. Caliciopsis calicioides, a corticolous ascomycete on Populus tremula new to Norway. Agarica 35: 7-12.
- Lechat C, Fournier J, Nordén B, 2015. Stylonectria norvegica (Nectriaceae), a new species from Norway. Ascomycete.org 7: 220-224.
- Miller JH & Gwendolyn B, 1942. Georgia Pyrenomycetes III. Mycologia, vol. 34: 1-7.
- Monod M, 1983. Monographie taxonomique des Gnomoniaceae. Sydowia Beihefte 9: 1-315.
- Nannfeldt JA, 1975. Stray studies in the Coronophorales (Pyrenomycetes) 1-3. Svensk Botanisk Tidskrift 69:49-66.
- Nordén B, 2014. Annulohypoxylon minutellum and Obolarina dryophila (Xylariales), two stromatic pyrenomycetes on oak new to Norway. Agarica 35: 25-28.
- Nordén B, 2016. A corticolous Verrucaria new to Norway. Graphis Scripta 28: 14-16.
- Nordén B, Andersson R, Andreasen M, Aptroot A, Chomnunti P, Friebes G, Jaklitsch W, Johannesen E, Jordal JB, Voglmayr H, 2019. Ascomycetes new to Norway from a workshop in Hordaland, 13-16 May 2019. Agarica 39: 53-59.
- Nordén B & Andreasen M, 2020. Final report Artsprosjekt 23-19 Endophytic fungi in trees. https://www.artsdatabanken.no/Pages/28177 3/Endofyttisk\_sopp\_i\_traer\_br\_small\_23-19 small ?Key=190.
- Nordén B & Andreasen M, 2022. Fremdriftsrapport Artsprosjekt 13-21 Pyrenomycetoid Ascomycota 2022. https://www.artsdatabanken. no/Pages/305728/Pyrenomycetoide\_sekkspo resopper\_i\_planter\_br\_small\_13-21\_small\_ ?Key=190.
- Nordén B, Aptroot A, 2018. Anisomeridium viridescens and Arthopyrenia callunae, two

pyrenolichens new to Fennoscandia. Graphis Scripta 30: 166-169.

- Nordén B, Gardiennet, A, Priou, JP, Döbbeler, P, 2015a. Bryocentria hypothallina (Hypocreales), a new species on Metzgeria furcata. Ascomycete.org 7: 121-124.
- Nordén B, Jäntti M, Jordal JB, 2017. Three species of Sordariomycetes (Ascomycota: Pezizomycotina) new to Norway. Agarica 38: 57-60.
- Nordén B, Jäntti M, Jordal JB, Læssøe T, Voglmayr H, Jaklitsch W, 2017. Twenty species of bitunicate ascomycetes new to Norway. Agarica 38: 47-56.
- Nordén B, Jordal, JB, 2014. Requienella seminuda, a corticolous ascomycete on Fraxinus excelsior new to Norway. Agarica 35: 29-34.
- Nordén B, Læssøe T, Jordal JB, 2014. Chlorostroma vestlandicum sp. nov, a host-specific mycoparasite on Hypoxylon vogesiacum from western Norway. Karstenia 54: 9-14.
- Nordén B, Læssøe T, Jordal JB, Petersen JH, Voglmayr H, Jaklitsch W, 2015. Forty pyrenomycetous fungi belonging to Class Sordariomycetes new to Norway. Agarica 36: 45-56.
- Olsen O, Haug S, Nordén B, 2019. Agonimia globulifera M. Brand, Diederich new to mainland Norway. Graphis Scripta 31: 1-4.
- Phillips A, Alves A, Abdollahzadeh J, Slippers B, Wingfield MJ, Groenewald JZ, Crous PW, 2013. The Botryosphaeriaceae: Genera and species known from culture. Studies in mycology 76. Pyrenomycetes from southwestern France, 2004. http://pyrenomycetes.free.fr /hypoxylon/html/Hypoxylon\_julianii.htm. Accessed April 16, 2024.
- Réblová M & Winka K, 2001. Generic Concepts and Correlations in Ascomycetes Based on Molecular and Morphological Data: Lecythothecium duriligni Gen. et sp. nov. with a Sporidesmium anamorph, and Ascolacicola austriaca sp. nov. Mycologia 93: 478-493.
- Shearer CA, 1989. Pseudohalonectria (Lasiosphaeriaceae), an antagonistic genus from wood in freshwater. Canadian Journal of Botany 67: 1944-1955.

- Shoemaker RA, 1984. Canadian and some extralimital Leptosphaeria species. Canadian Journal of Botany 62: 2688-2729.
- Shoemaker RA & Babcock CE, 1989. Phaeosphaeria. Canadian Journal of Botany 67: 1500-1599.
- Tanaka K, Hirayama K, Yonezawa H, Sato G, Toriyabe A, Kudo H, Hashimoto A, Matsumura M, Harada Y, Kurihara Y, Shirouzu T, Hosoya T, 2015. Revision of the Massarineae (Pleosporales, Dothideomycetes). Studies in Mycology 82: 75-136.
- The Norwegian Mycological Database (Norsk Soppdatabase) NSD, 2023. Norsk Soppdatabase (NSD). https://www.nhm2.uio.no/botanisk /nxd/sopp/nsd\_b.htm. Accessed September 10, 2023.
- Vasilyeva LN & Scheuer C, 1996. Neuere Aufsammlungen stromatisacher Pyrenomyceten aus Österreich, insbesondere der Steiermark. Mitteilungen des Naturwissenschaftlichen Vereines für Steiermark 126: 61-82.
- Voglmayr H & Jaklitsch WM, 2011. Molecular data reveal high host specificity in the phylogenetically isolated genus Massaria (Ascomycota, Massariaceae). Fungal Diversity 46: 133-170.

- Voglmayr H, Rossman AY, Castlebury LA, Jaklitsch WM, 2012. Multigene phylogeny and taxonomy of the genus Melanconiella (Diaporthales). Fungal Diversity 57: 1-44.
- Voglmayr H, Tello S, Jaklitsch WM, Friebes G, Baral H-O, Fournier J, 2022, About spirals and pores: Xylariaceae with remarkable germ loci. Persoonia 49: 58-98.
- Westberg M, Moberg R, Myrdal M, Nordin A, Ekman, S, 2021. Santesson's Checklist of Fennoscandian Lichen-Forming and Lichenicolous Fungi. Uppsala University: Museum of Evolution.
- Zhang Y, Wang HK, Fournier J, Crous PW, Jeewon R, Pointing S, Hyde K, 2008. Towards a phylogenetic clarification of Lophiostoma/Massarina and morphologically similar genera in the Pleosporales. Fungal Diversity 38: 38.