

Interactions among nutrients govern the global grassland biomass–precipitation relationship

Philip A. Fay^{a,1}, Laureano A. Gherardi^{b,1}, Laura Yahdjian^{c,1}, Peter B. Adler^d, Jonathan D. Bakker^e, Siddharth Bharath^f, Elizabeth T. Borer^f, W. Stanley Harpole^{g,h,1}, Erika Hersch-Green^j, Travis E. Huxman^k, Andrew S. MacDougall^l, Anita C. Risch^m, Eric W. Seabloom^f, Sumanta Bagchiⁿ, Isabel C. Barrio^o, Lori Biederman^p, Yvonne M. Bucklev^a, Miguel N. Bugalho^r, Maria C. Caldeira^s, Jane A. Catford^{tu}, QingQing Chen^{g,v}, Elsa E. Cleland^w, Scott L. Collins^x, Pedro Daleo^y, Christopher R. Dickman², Ian Donohue^{aa}, Mary E. DuPre^{bb}, Nico Eisenhauer^{g,cc}, Anu Eskelinen^{g,h,dd}, Nicole Hagenah^{ee}, Yann Hautier^{ff}, Robert W. Heckman^{gg,h,2}, Ian Donohue^{aa}, Jonsdóttirⁱⁱ, Johannes M. H. Knops^{jj}, Ramesh Laungani^{kk}, Jason P. Martina^{ll}, Rebecca L. McCulley^{mm}, John W. Morganⁿⁿ, Harry Olde Venterink^{oo}, Pablo L. Peri^{pp}, Sally A. Power^{qq}, Xavier Raynaud^{rr}, Zhengwei Ren^{ss,tt}, Christiane Roscher^{g,h}, Melinda D. Smith^{uu}, Marie Spohn^w, Carly J. Stevens^{ww}, Michelle J. Tedder^{xx}, Risto Virtanen^d, Glenda M. Wardle^w, and George R. Wheeler^{jzz}

Affiliations are included on p. 8.

Edited by Chao Zhang, University of Helsinki, Helsinki, Finland; received May 31, 2024; accepted March 6, 2025 by Editorial Board Member Donald R. Ort

Ecosystems are experiencing changing global patterns of mean annual precipitation (MAP) and enrichment with multiple nutrients that potentially colimit plant biomass production. In grasslands, mean aboveground plant biomass is closely related to MAP, but how this relationship changes after enrichment with multiple nutrients remains unclear. We hypothesized the global biomass-MAP relationship becomes steeper with an increasing number of added nutrients, with increases in steepness corresponding to the form of interaction among added nutrients and with increased mediation by changes in plant community diversity. We measured aboveground plant biomass production and species diversity in 71 grasslands on six continents representing the global span of grassland MAP, diversity, management, and soils. We fertilized all sites with nitrogen, phosphorus, and potassium with micronutrients in all combinations to identify which nutrients limited biomass at each site. As hypothesized, fertilizing with one, two, or three nutrients progressively steepened the global biomass-MAP relationship. The magnitude of the increase in steepness corresponded to whether sites were not limited by nitrogen or phosphorus, were limited by either one, or were colimited by both in additive, or synergistic forms. Unexpectedly, we found only weak evidence for mediation of biomass-MAP relationships by plant community diversity because relationships of species richness, evenness, and beta diversity to MAP and to biomass were weak or opposing. Site-level properties including baseline biomass production, soils, and management explained little variation in biomass-MAP relationships. These findings reveal multiple nutrient colimitation as a defining feature of the global grassland biomass-MAP relationship.

primary productivity | precipitation | diversity | grasslands

Terrestrial ecosystems across the globe are experiencing changes in mean annual precipitation (MAP), with MAP increasing in some regions and decreasing in others (1). Concurrently, many ecosystems are increasingly enriched with multiple elemental nutrients (2) including nitrogen (N), phosphorus (P), and potassium (K), which frequently colimit plant aboveground biomass production (3–13), a major component of ecosystem primary productivity (14). In grasslands, site-level mean aboveground biomass increases with increasing ecosystem MAP-the biomass-MAP relationship (15-23). At a global scale, the grassland biomass-MAP relationship emerges from across considerable variation in other factors regulating biomass production including, topography, soils, grazing, and other management (24). Theory predicts greater nutrient limitation of biomass production with increasing MAP (17, 19), reflecting higher demand for nutrients required to maintain plant carbon metabolism and water balance (13, 25, 26). Thus, fertilizing with limiting nutrients should result in a biomass-MAP relationship with a steeper slope (Fig. 1) (27-29). However, to what extent the increase in steepness depends on the number or identity of added nutrients or interactions among them is poorly understood. Clarifying the role of multiple nutrient interactions is necessary to forecast how interacting global change drivers—climate change and nutrient enrichment—will affect global patterns in energy flow, primary productivity, and ecosystem services. These processes are critical to societal efforts to mitigate and adapt to the impacts of global change drivers (2, 30, 31).

We propose a new framework—the "Multiple Nutrient Colimitation" Hypothesis predicting how the number of limiting nutrients and interactions among colimiting

Significance

Understanding how multiple interacting nutrients regulate the global relationship between mean annual precipitation and aboveground biomass is crucial for forecasting how ecosystem functioning will be altered by ongoing global changes. We fertilized with nitrogen, phosphorus, and potassium plus micronutrients in all combinations in 71 grasslands representing a global precipitation gradient. The grassland biomass-precipitation relationship became steeper with an increasing number of added nutrients. Increases in steepness corresponded to the form of interaction among added nitrogen and phosphorus. We found weak evidence that variation in plant species diversity mediated changes in the biomassprecipitation relationship. Multiple nutrient colimitation, particularly by nitrogen and phosphorus, is a defining feature of grassland biomass-precipitation relationships, and crucial to predicting grassland responses to global change.

Published April 11, 2025.

¹To whom correspondence may be addressed. Email: philip.fay@usda.gov, lau.gherardi@berkeley.edu, or yahdjian@agro.uba.ar.

²Present address: United States Department of Agriculture Forest Service, Rocky Mountain Research Station, Cedar City, UT 84721.

This article contains supporting information online at https://www.pnas.org/lookup/suppl/doi:10.1073/pnas. 2410748122/-/DCSupplemental.

Form of Nutrient Limitation	Definition	Mean Biomass Response (Arbitrary units)	Predicted Biomass- MAP Relationships
No Limitation	Biomass does not increase with A or B, alone or together.	$\begin{array}{c} 3\\2\\1\\0\\-\hline\\-1\\\hline\\Control \dot{A} & \dot{B} & AB \end{array}$	Biomass
Single	Biomass increases only when one nutrient is added (A) is added, with or without the other (B).	3 2 1 0 	Biomass
	Multiple Nutri	ent Co-limitation	
Additive	Biomass increase with A and B together is the sum of the increases with A and B singly.	$\begin{array}{c} 3 \\ 2 \\ 1 \\ 0 \\ - \frac{1}{2} \\ - \frac{1}{$	Bionass
Subadditive	Biomass increase with A and B together is less than the sum of A and B singly. At least one single nutrient must increase biomass.	3 2 1 0 - - -1 Control À B AB	Blomass
Synergistic	Biomass increase with A and B together is greater than the sum of A and B singly. None, one, or both single nutrients may increase biomass.	$\begin{array}{c} 3 \\ 2 \\ 1 \\ 0 \\ - \frac{1}{9} \\ - \frac{1}{10} \\ \hline \\ \text{Control} \\ \dot{A} \\ \dot{B} \\ \dot{B} \\ \dot{A} \\ \dot{A} \\ \dot{A} \\ \dot{B} \\ \dot{A} \\ \dot{A} \\ \dot{B} \\ \dot{A} \\ \dot{B} \\ \dot{B} \\ \dot{A} \\ \dot{B} \\ \dot{A} \\ \dot{A} \\ \dot{A} \\ \dot{B} \\ \dot{A} \\ \dot{A} \\ \dot{B} \\ \dot{A} \\ \dot{A} \\ \dot{B} \\ \dot{A} \\ \dot{B} \\ \dot{A} \\ \dot{A} \\ \dot{B} \\ \dot{A} \\ \dot{B} \\ \dot{A} \\ \dot{B} \\ \dot{A} \\ \dot{B} \\ \dot{A} \\ \dot{A} \\ \dot{A} \\ \dot{B} \\ \dot{A} \\ \dot{A} \\ \dot{B} \\ \dot{A} \\ \dot{A} \\ \dot{B} \\ \dot{A} \\ $	Blomass

Fig. 1. Conceptual framework for predicting changes in the steepness of the global mean aboveground biomass–MAP relationship in response to fertilization with hypothetically limiting nutrients A, B, and A together with B (AB). A and B represent any two nutrients that potentially limit biomass production. The mean response of biomass across sites to fertilization defines the number of nutrients limiting aboveground biomass and the form of interactions among colimiting nutrients—additive, subadditive, or synergistic. The increase in slope of the grassland biomass–MAP relationship is predicted from the mean biomass response. Limitation forms are generalized from ref. 10. Application of the rubric is detailed in *SI Appendix, Extended Methods*.

nutrients influence the global grassland biomass–MAP relationship. This hypothesis holds that the effects of fertilization on the steepness of the biomass–MAP relationship: 1) increase with the number of added nutrients and attendant reduction in multiple nutrient limitation and 2) corresponds to the form of interaction among multiple nutrients (10). This hypothesis extends previous concepts of the controls on the grassland biomass–MAP relationship (17, 19, 27–29) by explicitly accounting for the number of limiting nutrients and how colimiting nutrients interact (Fig. 1). This idea also builds on previous findings of widespread multiple nutrient colimitation of productivity in grasslands (12, 32) and other ecosystems (10, 33).

Interactions among colimiting nutrients occur in several forms defined by the responses in aboveground biomass when a site is fertilized with nutrients individually and in combination (Fig. 1) (10, 34). Synergistic colimitation is present when the response of biomass to fertilizing with multiple nutrients is greater than the sum of the single nutrient responses and is common in terrestrial ecosystems (10, 34). Additive colimitation occurs when the response to multiple nutrients equals the sum of the single nutrient responses. Subadditive colimitation occurs when the response is less than the sum of the single nutrient responses; this form is uncommon (10, 34) and may reflect intensification of another limiting factor. Single nutrient limitation or no responses to the added nutrients are also possible. The Multiple Nutrient Colimitation hypothesis predicts that fertilizing grasslands with multiple nutrients should cause a synergistic increase in slope of the biomass–MAP relationship across sites where mean biomass production is synergistically colimited, and corresponding responses in slope across sites where colimitation is additive, subadditive, or where limitation is by a single nutrient. However, the correspondence between the increase in slope of the global grassland biomass–MAP relationship and the form of nutrient limitation has not been evaluated, primarily because the multiple nutrient enrichment experiments needed to directly test these effects in grasslands spanning a globally relevant range of MAP have only recently become available (35).

Plant community diversity plays a central role in mediating biomass production (36–42). Thus, variation in plant community diversity is expected to mediate the relationships of MAP and added nutrients to biomass production. For example, with increasing MAP, sites should increase in species richness (43–47) and favor faster-growing, more productive species (48) which may also have higher nutrient requirements (28, 49). At the same time, higher-MAP communities may also be more susceptible to the synergistic effects of adding multiple limiting nutrients, including amplified plant species losses (29, 50, 51), stronger dominance (52–55), or other deviations from the regional species pool (56, 57). Therefore, we predict greater mediation of MAP and nutrient effects on biomass by plant community diversity in synergistically colimited grasslands than in grasslands with other forms of nutrient limitation. Resolving whether the form of nutrient limitation alters mediation of the global Downloaded from https://www.pnas.org by SLU BIBLIOTEKET; SVERIGES LANTBRUKSUNIVERSITETS ULTUNA on May 12, 2025 from IP address 193.10.98.235

grassland biomass–MAP relationship by plant diversity is crucial for forecasting how climate change and eutrophication impact the provision of biomass-related ecosystem services in grasslands.

Here, we test the Multiple Nutrient Colimitation hypothesis by analyzing the relationship of site mean aboveground biomass production (hereafter, "biomass") to site MAP across 71 grasslands (SI Appendix, Table S1) in a global multiple nutrient fertilization experiment, the Nutrient Network (58). These grasslands were distributed across six continents (SI Appendix, Fig. S1) and spanned 12 to 991 g m^{$^{-2}$} aboveground biomass, 167 to 1,823 mm y^{$^{-1}$} MAP, -3.3 to 24.1 °C mean annual temperature, and 0 to 4,241 m elevation (SI Appendix, Table S1). The sites included native and planted grasslands with varying soil textures and soil nutrient contents. Of the 71 sites, 26 were reported by site investigators to be under active management, mostly burning or grazing (17 sites). The form of management was unspecified for the remainder. Management contributes to variation in relationships among precipitation, biomass production, and plant diversity (59). Thus, we evaluated the effects of multiple nutrient fertilization on the biomass-MAP relationship across broadly representative global precipitation gradients and realistic sources of complexity in grassland ecosystem structure and function. Standardized multiple nutrient fertilization treatments were conducted for 4 to 14 y (SI Appendix, Table S1 and Abbreviated *Methods*). All sites fertilized 5 m \times 5 m plots once each year with 10 $g m^{-2}$ each of N, P, and K in factorial combination in a randomized blocks design with at least three replicates. This fertilization rate exceeds global mean deposition rates for these nutrients (0.4 to 1.5 g m⁻² y⁻¹) (60–62). The K treatment included micronutrients in years 1 and 10 (Kµ). Peak live aboveground biomass and plant species cover were determined annually. From the cover, we derived effective species richness (eH), evenness, and beta diversity (βplot). Site MAP was the average of the precipitation accrued between the annual biomass harvests (SI Appendix, Climate Variables). Biomass and the variables derived from cover in each plot were averaged across years by plot to correspond in temporal scale to site MAP (SI Appendix, Extended Methods).

Our analysis addressed three primary research questions about grassland biomass—MAP—nutrient interactions: 1) Does the global biomass—MAP relationship become steeper with increased number of added nutrients? 2) Does the increased steepness of the biomass—MAP relationship correspond to the form of nutrient limitation? 3) Does the form of nutrient limitation alter the mediation of MAP and nutrient effects on biomass by community diversity?

Results

Does the Global Biomass-MAP Relationship Become Steeper with Increased Number of Added Nutrients? Across all 71 sites, the biomass of unfertilized control plots significantly increased with MAP (slope = 0.31, $R^2 = 0.19$, P < 0.0001, *SI Appendix*, Table S2). As hypothesized, fertilization increased the steepness of the biomass-MAP relationship, and this increase was more pronounced when greater numbers of nutrients were added (Fig. 2*A*; MAP × Number of Nutrients P < 0.0001, Table 1). Fertilization with all three nutrients-N, P, and Kµ together increased the slope of the biomass-MAP relationship by 51% compared to the baseline slope for control plots (Fig. 2 *A*, *Inset* and *SI Appendix*, Table S2). The increase in slope was smaller, 33%, for fertilization with pairs of nutrients (NP, NKµ, or PKµ), and smallest, 19%, for fertilization with single nutrients (N, P, or Kµ alone; Fig. 2 *A*, *Inset*).

Fertilizing with greater numbers of nutrients increased the steepness of the biomass-MAP relationship across all sites primarily because slopes increased under fertilization with N (P < 0.0001) and P (<0.0001, Fig. 2B and Table 1). Fertilizing with N alone increased the slope by 26% compared to unfertilized plots (Fig. 2 B, Inset and SI Appendix, Table S2), while fertilizing with P alone increased the slope by only 19%. In contrast, fertilizing with N and P together increased the slope 36%. N and P did not interact with MAP to influence biomass (P = 0.86, Table 1). This signifies that fertilizing with N and P together caused an additive increase in the steepness of the biomass-MAP relationship (Fig. 1). Fertilizing with Kµ together with N and P caused slight increases in the slope of the biomass–MAP relationship (Fig. 2*B* and *SI Appendix*, Table S2) but there was little indication that Kµ interacted with MAP, N, or P (0.06 < P < 0.86, Table 1). These findings support the hypothesis that increasing numbers of nutrients, and particularly fertilizing with N and P, increases the steepness of the grassland biomass-MAP relationship.

Does the Increased Steepness of the Biomass–MAP Relationship Correspond to the Form of Nutrient Limitation? Unexpectedly, across all 71 sites the additive increase in steepness of the global biomass–MAP relationship did not correspond to the effects of N and P fertilization on the mean biomass production across all sites. Adding N and P together synergistically increased mean aboveground biomass (Fig. 2C; N × P P = 0.008, Table 1). Biomass rose43% compared to 23% for adding N alone and 6% for P alone.



Fig. 2. Responses of the global mean biomass—MAP relationship to fertilization with single N, P, and potassium with micronutrients ($K\mu$). (*A*) The biomass–MAP relationships for treatments fertilizing with 0, 1, 2, or 3 nutrients. Inset: the percent increase in linear regression slope relative to unfertilized controls. (*B*) the biomass–MAP relationships for treatments fertilizing with N, P, and K μ in factorial combinations. Nutrient treatments are color-coded as in Panel *C*. (*Inset*) The percent increase in linear regression slopes relative to unfertilized controls for N, P, and NP treatments averaged across levels of K μ . (*C*) Mean ± SE of aboveground biomass across all 71 sites for the factorial N, P, and K μ fertilization treatments. See Table 1 for linear mixed model analyses and *SI Appendix*, Table S2 for linear regression equations.

Table 1. Linear mixed model F statistics, degrees of freedom (dfs), and *P*-values testing the effects of the number of nutrients added, N, P, potassium with micronutrients (K μ), MAP, and their interactions on aboveground biomass across all 71 sites

Mixed model effects	F(dfs)	P-value			
Number (#) of Nutrients					
# of Nutrients	72.5(3,125)	<0.0001			
MAP	27.9(1, 69)	<0.0001			
MAP × # of nutrients	8.4(3,131)	<0.0001			
Factorial nutrient co	ombinations				
Ν	260.7(1,447)	<0.0001			
Р	63.6(1,447)	<0.0001			
N*P	7.0(1,447)	0.0084			
К	0.3(1,447)	0.5604			
N*K	0.1(1,447)	0.7981			
P*K	0.6(1,447)	0.4586			
N*P*K	0.8(1,447)	0.3680			
MAP	28.3(1, 69)	<0.0001			
MAP*N	19.2(1,460)	<0.0001			
MAP*P	4.3(1,460)	0.0376			
MAP*N*P	0.0(1,460)	0.8554			
MAP*K	3.6(1,460)	0.0577			
MAP*N*K	0.2(1,460)	0.6657			
MAP*P*K	0.2(1,460)	0.6919			
MAP*N*P*K	0.2(1,460)	0.6541			

Model effects and statistics for number of nutrients added correspond to Fig. 2*A*, and for factorial nutrient combinations correspond to Fig. 2 *B* and *C*. *P*-values are in bold font when <0.05.

The synergistic mean biomass response corresponded instead to a synergistic increase in the intercept of the global biomass–MAP relationship rather than in the slope (*SI Appendix*, Table S2). However, the absence of Kµ effects on global mean biomass (0.37 < P < 0.80, Table 1) was consistent with the absence of interaction of Kµ with MAP.

Instead, increases in steepness of the biomass–MAP relationship corresponded with the effects of N and P fertilization across sites expressing the same form of limitation by N and P (*SI Appendix*, Table S2). For example, across the 15 sites classified as synergistically colimited by N and P (Fig. 3*D*), adding N and P together caused a synergistic increase in slope (20%, MAP × N × P P = 0.04, Table 2 and *SI Appendix*, Table S2), while fertilization with N or P individually caused little effect on slope (Fig. 3 *D*, *Inset*). The correspondence between increases in slope and form of nutrient limitation continued in sites classified as additively colimited by N and P (21 sites, Fig. 3*C*), single-nutrient limited (13 sites by N, 3 by P, Fig. 3*B*). The correspondence of limitation form with slope response even extended to sites not limited by N or P (15 sites, Fig. 3*A*), where slopes did not respond to adding N or P (Table 2 and *SI Appendix*, Table S2).

The correspondence of increases in slopes of biomass–MAP relationships to the form of nutrient limitation emerged from substantial differences among limitation forms in the strength of the baseline biomass–MAP relationships for unfertilized grassland (MAP × form P < 0.0059, Table 3). For example, across synergistically colimited sites, the baseline unfertilized biomass–MAP relationship was steepest (slope = 0.88) and explained the most variation (R² = 0.49) among the four limitation classes (Fig. 4*A*). In contrast, across No Limitation sites, baseline aboveground biomass was uncorrelated with MAP (*SI Appendix*, Table S2). Baseline biomass–MAP relationships across single nutrient-limited and across additive N-P colimited sites fell between these extremes (slopes = 0.27 to 0.46; R² = 0.27 to 0.37).

Synergistically colimited sites spanned a lower range of MAP (up to 1,013 mm, Fig. 3D) than sites in the other limitation categories. This could explain the steeper baseline biomass-MAP relationship because in drier regions primary production is increasingly controlled by precipitation inputs (63). When we standardized the range of MAP in the limitation categories by excluding sites with >1,013 mm MAP (the maximum of the synergistic sites) from the Additive, Single, and No Limitation categories (Fig. 4B), baseline slopes did not differ among limitation forms (MAP × form P = 0.43, Table 3 and *SI Appendix*, Table S2). Importantly, the correspondence between the form of limitation and the increases in slope with addition of N and P remained (SI Appendix, Fig. S2). Other site-level factors potentially affecting how fertilization with N and P increased the steepness and strength of biomass-MAP relationships, including management, latitude, elevation, mean annual temperature and the texture, and nutrient content of soils did not differ in occurrence or magnitude among limitation categories (SI Appendix, Table S3). These findings show that the effect of fertilization with N and P on the steepness of the biomass-MAP relationship corresponded to the



Mean Annual Precipitation, mm y⁻¹

Fig. 3. Aboveground biomass in relation to MAP for sites classified by form of response to N and P fertilization. (*A*) No Limitation, (*B*) Limited by Single N or P, (*C*) Additive limitation by N and P, (*D*) Synergistic limitation by N and P). Response forms are defined in Fig. 1. N and P treatments are averaged across levels of Kµ fertilization. *Insets* depict the slopes for unfertilized control (C), N, P, and N together with P. *Upper* panels are kernel-smoothed MAP distributions for the sites each form of limitation. See Table 2 for linear mixed model analyses and *SI Appendix*, Table S2 for linear regression equations.

Table 2.	Linear mixed model F statistics ar	nd <i>P</i> -values testing the	effects of N, P,	and MAP, and their	interactions on
abovegr	ound biomass for sites classified b	y form of limitation by	y N and P (Fig. 3)	

	No limita	ition	Singl	e	Additi	ve	Synerg	gistic
Mixed model effects	F(dfs)	P-value	F(dfs)	P-value	F(dfs)	P-value	F(dfs)	P-value
N	4.9(1,281)	0.0278	109.6(1, 89)	<0.0001	204.3(1,133)	<0.0001	55.0(1,96)	<0.0001
Р	0.5(1,281)	0.4598	8.8(1, 89)	0.0039	55.4(1,133)	<0.0001	48.3(1,96)	<0.0001
N*P	0.0(1,281)	0.9862	0.5(1, 89)	0.4739	0.5(1,133)	0.4837	30.4(1,96)	<0.0001
MAP	0.3(1, 13)	0.6091	14.3(1, 14)	0.0020	8.7(1, 19)	0.0081	13.3(1, 13)	0.0030
MAP*N	2.0(1,281)	0.1627	4.6(1,99)	0.0344	8.8(1,143)	0.0035	1.7(1,93)	0.1917
MAP*P	2.3(1,281)	0.1297	0.0(1,99)	0.8339	4.4(1,143)	0.0383	5.0(1,93)	0.0279
MAP*N*P	0.6(1,281)	0.4454	0.2(1,99)	0.6687	0.0(1,143)	0.9207	4.4(1,93)	0.0397

P-values are in bold font when <0.05. N and P effects and interactions are across levels of Kµ.

form of nutrient limitation across variation in other factors that can potentially influence this relationship.

Does the Form of Nutrient Limitation Alter Mediation of MAP and Nutrient Effects on Biomass by Community Diversity? Contrary to our hypothesis, we found little evidence that plant community diversity mediated relationships of MAP to biomass. Structural equation models successfully partitioned indirect ("community mediated") and direct effects of fertilization and MAP on biomass (P > 0.59, Table 4 and SI Appendix, Fig. S3). Across sites in each form of nutrient limitation, MAP was always the largest driver of aboveground biomass. Total (direct + indirect) MAP effects ranged from 0.18 across No Limitation sites to 0.67 across synergistic colimitation sites and were 2 to 3 times greater than total nutrient effects (Fig. 5A and SI Appendix, Table S4). Total MAP effects were almost entirely explained by direct effects (Fig. 5B). Indirect, community-mediated effects of MAP were near 0 (0.08 < P < 0.41) for No Limitation and Synergistic limitation sites and small (5 to 8% of total effects, *P* < 0.045, Fig. 5*C* and *SI Appendix*, Table S4) for Single and Additive limitation sites. Indirect effects of MAP were weak because the individual paths linking the diversity variables to MAP and biomass were either not significant or offsetting (SI Appendix, Fig. S3). In contrast, indirect nutrient effects on biomass ranged from 9 to 30% of total effects (P < 0.045, Fig. 5C and SI Appendix, Table S4) and were largest in Synergistic limitation sites, consistent with large decreases in means of eH, Evenness, and βplot across all sites (*SI Appendix*, Fig. S4).

Discussion

Anthropogenic global changes are causing long-term changes in MAP while concurrently enriching ecosystems with multiple potentially limiting nutrients (1, 2). These changes will have significant consequences for aboveground plant biomass production, a key component of primary productivity, global carbon cycling, and many ecosystem services. Our findings largely supported the

Table 3. Linear mixed model F statistics and *P*-values testing the effects of the form of nutrient limitation (Form), MAP and their interaction on unfertilized aboveground biomass (Fig. 4)

Mixed model	All M	AP	MAP < 1,013 mm		
effects	F(dfs)	P-value	F(dfs)	P-value	
Form	0.7(3, 56)	0.5621	0.3(3, 43)	0.8014	
MAP	14.7(1, 56)	0.0003	12.0(1, 43)	0.0012	
MAP*Form	4.6(3, 56)	0.0059	0.9(3, 43)	0.4260	

P-values are in bold font when <0.05.

Multiple Nutrient Colimitation hypothesis. They demonstrated that the global grassland biomass–MAP relationship became steeper as the number of added nutrients increased (question 1) and that the increased steepness of the biomass–MAP relationship corresponded to the form of nutrient limitation (question 2). However, we found little evidence that the form of nutrient limitation affected mediation of the biomass–MAP relationship by the combined effects of eH, evenness, or β plot (question 3). These findings provide robust evidence that the grassland biomass–MAP relationship is limited by multiple nutrients (17, 19) and reveal the key importance of the number, identity, and interactions among the limiting nutrients.

It is well established that the predominant forms of nutrient limitation of biomass production-single limitation, additive colimitation, synergistic colimitation, or no nutrient limitationare widespread in the world's grasslands (9, 10, 12, 32, 51). These findings build on this foundation by revealing the correspondence of the increased steepness of the biomass-MAP relationship to the mean response to fertilizing with single and multiple nutrients, in particular N and P (Figs. 2 and 3). Steeper biomass-MAP relationships in response to fertilization are consistent with greater nutrient uptake to meet greater demand by plant metabolism (13, 25, 26) and with steeper and more variable within-site temporal biomass-precipitation relationships (64). The present findings extend the generality of our previous findings of widespread site-level synergistic colimitation of grassland biomass production by N and P (12, 32) by including nearly 30 more sites with many fertilized for over a decade (vs. 4 to 7 y), allowing more forms of nutrient interactions to be resolved.

Steeper biomass-MAP relationships in response to fertilizing nutrient-limited sites with N and P indicated that increases across sites in mean water availability increasingly translated into greater plant biomass production (16, 19, 22, 23, 65, 66). However, the biomass-MAP relationships displayed considerable scatter in all fertilization treatments. For example, MAP explained only about 20% of the variation in plant biomass across all 71 sites in the control treatment (Fig. 2B), and fertilization treatments yielded only modest increases in variation explained by MAP (25 to 30%; SI Appendix, Table S2). The relatively high baseline variation seen in our study may be unsurprising for sites spanning multiple continents, large differences in plant species assemblages, and varying management, soils, latitude, and aspects of climate other than MAP (SI Appendix, Tables S1 and S3). However, in the synergistically colimited sites, MAP explained about 50% of the variation in biomass (SI Appendix, Table S2), approached levels previously reported (65). The synergistically colimited grasslands may be more similar to each other in the mechanisms linking biomass production to water availability than are sites in the other limitation forms we identified.



Fig. 4. Baseline global biomass–MAP relationships defined by unfertilized controls across sites classified by form of response to N and P fertilization (Fig. 3). (A) Biomass–MAP relationships across the MAP range spanned by all 71 sites. (B) Biomass–MAP relationships for sites with MAP up to 1,013 mm. See Table 3 for linear mixed models analyses and *SI Appendix*, Table S2 for linear regression equations.

Our findings ran counter to our prediction that plant community change would mediate the biomass-MAP relationship (36, 67, 68). We found weak evidence that combined changes in effective species richness, evenness, or beta diversity mediated the biomass-MAP relationship, despite stronger community mediation of fertilization effects on biomass, which aligns with previous findings that fertilization reduces compositional stability (57, 69, 70) and synergistic global responses in two out of three measures of species diversity in response to fertilization with N and P (SI Appendix, Fig. S4). The weak signal for community diversity mediation of the biomass-MAP relationship is also consistent with findings that large compositional shifts in global grasslands with plant invasion were unable to explain changes in biomass over the last several decades (24). However, including these diversity metrics in our analysis, even if their net effect was small, was still important. SEMs resolved a significant biomass-MAP relationship in sites not limited by N or P (Fig. 5B and SI Appendix, Fig. S3 and Table S4) where analysis not accounting for diversity (Fig. 3A) did not. This finding highlights the importance of accounting for changing plant community attributes when evaluating factors governing the global biomass-MAP relationship.

Several underlying mechanisms may explain the weak community mediation finding. Fertilization effects on species diversity within sites may be poor predictors of responses across larger spatial gradients (71) because the MAP gradient encompasses large, potentially nonlinear diversity changes (47) while within-site

Table 4. Fit statistics for structural equation models fitacross grassland sites assigned to four forms of nutri-ent limitation by nitrogen and phosphorus (Fig. 5)

	Fit statistic				
Kind of limitation	X2 (P-value)	RMSEA	PCF		
	Thresholds				
	<i>P</i> > 0.05	<0.06	>0.05		
No Limitation	1.07 (0.5860)	0.0000	0.6962		
Single	0.25 (0.8803)	0.0000	0.9284		
Additive	0.83 (0.6597)	0.0000	0.8128		
Synergistic	0.06 (0.9711)	0.0000	0.9823		

 χ^2 : Chi-square test. RMSEA:Rootmean square error. PCF:Probability of close fit.

responses are limited by the local species pool. In addition, longer time periods may be required to detect plant community mediation than the 4 to 14 y of observation we had available here (36, 67). Biomass and diversity responses increased through 11 y of fertilization with N, P and K (72), so community mediation may emerge when more sites accumulate more years of fertilization (41). Finally, we did not consider abundance-weighted composition metrics (42) or functional diversity, which decreased with nutrient addition across some Nutrient Network sites (73) and in some instances may better predict ecosystem function than species-based metrics alone (74, 75). However, biomass gains following fertilization can be explained by plant species that persist following fertilization rather than by replacement (53-55, 72). Further analysis of plant compositional and functional dynamics in grasslands differing in form and strength of single and multiple nutrient limitation is a promising area for future research.

We continue to find little evidence for global-average limitation of grassland biomass by Kµ, alone or in combination with N and P, although a few individual sites are Kµ limited (12, 32). The infrequency of K-limitation may reflect several factors. K and micronutrients are broadly abundant in many surface soils, and K is strongly retained by plants compared to N or P (13). Plant tissue K concentration is tightly regulated because of its critical role in metabolite transport and water balance. However, K is also susceptible to loss from the rooting zone by leaching, which may be more prevalent in sites expressing K-limitation (13, 62). We also found no evidence for differences among limitation forms in occurrence or magnitude of site-level factors (SI Appendix, Table S3) that might influence fertilization effects on biomass-MAP relationships. The prevalence of N and P limitation varies with latitude, temperature, and fire frequency (12, 76, 77), and background soil nutrient levels and texture can influence retention and uptake of water and nutrients (78-81). Resolving the contributions of these factors will be aided by the network's growing dataset of responses to multiple nutrient fertilization.

Our findings point to a critical need for better understanding of edaphic mechanisms causing single and multiple nutrient limitation in grasslands. Because water is necessary for biogeochemical processes, mechanisms likely center on ways that water availability influences nutrient availability (82, 83). The number



Fig. 5. Summary of the standardized effects of nutrient addition and MAP on aboveground biomass production from structural equation models fit (Table 4) across grassland sites in each of four forms of nutrient limitation. (*A*) Sankey plot depicting Total effects (direct + indirect) of number of nutrients and MAP (*Left* side) mapped onto each form of nutrient limitation (*Right* side). The widths of the links depict the magnitude of each total effect for each limitation form. (*B*) Direct and (*C*) indirect effects (±SE) of MAP and nutrient addition for each limitation form. Indirect effects represent community mediation of MAP and nutrient effects on biomass, and combine paths through effective species richness, species evenness, and beta diversity (*SI Appendix*, Fig. S3).

of limiting nutrients and the form of interaction between colimiting nutrients may depend on the alignment of water availability and nutrient availability. Water availability interacts with soil parent material, microbial processing, biogeochemical cycling, stoichiometry, and plant uptake of the nutrients (84–88). For example, Vázquez et al. (89) found that synergistic increases in aboveground biomass production resulted in part from enhanced N and P uptake and retention. A comprehensive spatial model incorporating mechanistic drivers of single and multiple nutrient limitation is needed to link with productivity models to predict global scale responses of ecosystem productivity to changing precipitation and eutrophication.

The core findings of this study support our Multiple Nutrient Colimitation hypothesis. The effects of fertilization on the steepness of the global biomass-MAP relationship: 1) increases with the number of added nutrients and attendant reduction in multiple nutrient limitation, and 2) corresponds to the form of interactions among colimiting nutrients, particularly N and P. This critical, globally relevant insight into the regulation of grassland productivity can be exploited to predict the interactive effects of eutrophication and hydrologic intensification on grassland productivity and related ecosystem services. Applying this insight will require predicting the number and form of interaction among multiple limiting nutrients across the world's grasslands, and developing a more general understanding of magnitude and extent to which plant community change mediates grassland productivityprecipitation relationships and of the edaphic mechanisms controlling nutrient limitation.

Abbreviated Methods

Fertilizer Treatments. Each site applied fertilizers following the Nutrient Network standard experimental protocol (58). Nitrogen (as time release urea), phosphorus (as triple super phosphate), and potassium (as K_2SO_4) were hand-spread on 5 m × 5 m plots. The application rate of 10 g m⁻² y⁻¹ of each element was chosen because it is expected to exceed plant demand (58). In year 1 and year 10, 100 g of micronutrients were applied with K as Everris MicromaxTM. Sites had a minimum of three replicates per treatment, but a few sites maintained up to five replicates. In total,

the 4 to 14 y of site-level fertilization treatments (*SI Appendix*, Table S1) yielded 15,204 experimental plot-years.

Biomass and Diversity. Each site used network protocols to measure peak aboveground live plant biomass (in g m⁻² y⁻¹) and the percent cover of each plant species in designated portions of each plot (58). Biomass was sampled by clipping and drying to constant weight (*SI Appendix, Extended Methods*). Species cover was assessed visually to the nearest percent. From species cover, we derived three diversity metrics: the effective number of species (e^H), representing species richness if all species were equally abundant (90); Whitaker's beta (βplot), the ratio of site level species richness to plot-level species richness (91), and Evenness (E), describing the distribution of species relative abundances and the inverse of dominance.

Site MAP. For most (59) sites, MAP was derived from daily precipitation measured at a weather station selected by the local investigator (*SI Appendix*, Table S7). MAP was the mean of precipitation summed from harvest to harvest for the selected years of biomass and cover data. For the remaining 12 sites, we determined MAP from downscaled precipitation estimates (92). We validated the comparability of the two sources of MAP (*SI Appendix*, Figs. S5 and S6). We chose site MAP as the predictor of aboveground biomass after screening 31 other potential site metrics of precipitation, temperature, and evaporative demand (*SI Appendix*, Table S8). This analysis did not consider within-site temporal variability in annual precipitation amounts, which has been examined elsewhere (32, 64).

Statistical Analysis. Analytic approaches were linear mixed models of the interactive effects of the number or identity of nutrients with MAP across all sites and for subsets of sites assigned to one of four nutrient limitation categories. Community mediation of nutrient and MAP effects on biomass were evaluated using structural equation models. Preparation and analysis of the biomass and diversity data are detailed in the *SI Appendix, Extended Methods*.

Data, Materials, and Software Availability. Datasets and code used in this analysis have been deposited in Dryad (10.5061/dryad.vdncjsz50) (93).

ACKNOWLEDGMENTS. This work was conducted within the Nutrient Network (http://www.nutnet.org) experiment, funded at the site-scale by individual researchers. We thank the Minnesota Supercomputer Institute for hosting project data, the University of Minnesota Institute on the Environment for hosting Nutrient Network meetings, and each site investigator for funding their site-level operations. Network coordination and data management were supported by funds from the NSF Research Coordination Network (NSF-DEB-1042132) to E.T.B. and E.W.S., from the Long Term Ecological Research program (NSF-DEB-1831944) to the Cedar Creek LTER, and from the Institute on the Environment (DG-0001-13). The Co-authors acknowledge funding from the United States Department of Agriculture (USDA) (USDA-NIFA-2010-65615-20632) and the Swiss Federal Institute for Forest, Snow, and Landscape Research. The USDA is an equal opportunity employer and provider (P.A.F.). Consejo Nacional de Investigaciones Científicas y Técnicas and Agencia Nacional de Promoción de la Investigación, el Desarrollo Tecnológico y la Innovación, Argentina PICT 2014-3026 (L.Y.). University of Iceland Research Fund (2015), OrkurannsóknasjóÐur Landsvirkjunnar NÝR-09-2017, NÝR-14-2018, NÝR-12-2019 (I.C.B.). Co-Centre award number 22/CC/11103, managed by Science Foundation Ireland, Northern Ireland's Department of Agriculture, Environment and Rural Affairs and UK Research and Innovation, and supported via UK's International Science Partnerships Fund, and the Irish Government's Shared Island initiative (Y.M.B.). Fundação para a Ciência e Tecnologia funding to research units Forest Reseach Center (UIDB/00239/2023) and Research Network in Biodiversity and Evolutionary Biology (UID/BIA/50027/2023) and to Rui Alves for granting access to study site of Companhia das Lezírias (M.N.B. and M.C.C.). European Research Council under the European Union's Horizon 2020 research and innovation programme grant agreement No. [101002987] (J.A.C.). Bush Heritage Australia and the Wangkamadla people for access to the sites (C.R.D. and G.M.W.). iDiv funded by the German Research Foundation DFG-FZT 118, 202548816, and DFG Ei 862/29-1 (N.E.). Academy project no 351089 (A.E.). USDA Forest Service, Rocky Mountain Research Station (R.W.H.). University of Iceland Research Fund (I.S.J.). CEREEP-Ecotron IleDeFrance Centre National de la Recherche Scientifique/Ecole Normale Superior UMS 3194 (X.R.).

Author affiliations: ^aUnited States Department of Agriculture, Agricultural Research Service, Grassland, Soil, and Water Lab, Temple, TX 76502; ^bDepartment of Environmental Sciences, Policy, and Management, University of California, Berkeley, CA 94720; ^cInstituto de Investigaciones Fisiológicas y Ecológicas Vinculadas a la Agricultura-Consejo Nacional de Investigaciones Científicas y Técnicas, Cátedra de Ecología, Facultad de Agronomía, Universidad de Buenos Aires, Ciudad Autónoma de Buenos Aires C1417DSE, Argentina; ^dDepartment of Wildland Resources and the Ecology Center, Utah State University, Logan, UT 84322; ^eSchool of Environmental and Forest Sciences, University of Washington, Seattle, WA 98195; [†]Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, MN 55108; [®]German Centre for Integrative Biodiversity Research, Halle-Jena-Leipzig, Leipzig 04103, Germany; [†]Department of Physiological Diversity, Helmholtz Center for Environmental Research, Leipzig 04318, Germany; [†]Martin Luther University Halle (Saale) 06108, Germany; [†]Department of Ecology and Evolutionary Biology, University of California, Irvine, CA 92697; [†]Department of Integrative Biology, University of Guelph, ON N1G 2W1, Canada; ^mSwitserland; ⁿCentre for Ecologi, Sciences, Indian Institute of Science, Bangalore 560012, India; ^oFaculty of Environmental

- H. Douville et al., "Water cycle changes" in Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, V. Masson-Delmotte et al., Eds. (Cambridge University Press, Cambridge, United Kingdom and New York, NY, 2021), pp. 1055–1210, 10.1017/9781009157896.010.
- 2. J. Rockström et al., A safe operating space for humanity. Nature 461, 472-475 (2009).
- A. J. Bloom, F. S. Chapin III, H. A. Mooney, Resource limitation in plants-an economic analogy. Ann. Rev. Ecol. Sys. 16, 363–392 (1985).
- F. S. Chapin, A. J. Bloom, C. B. Field, R. H. Waring, Plant-responses to multiple environmentalfactors. *Bioscience* 37, 49–57 (1987).
- S. K. Gleeson, D. Tilman, Plant allocation and the multiple limitation hypothesis. Am. Nat. 139, 1322–1343 (1992).
- E. B. Rastetter, G. R. Shaver, A model of multiple-element limitation for acclimating vegetation. Ecology 73, 1157–1174 (1992).
- V. Smil, Phosphorus in the environment: Natural flows and human interferences. Annu. Rev. Energy Environ. 25, 53–88 (2000).
- J. N. Galloway et al., Nitrogen cycles: Past, present, and future. Biogeochemistry 70, 153–226 (2004).
- J. J. Elser et al., Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecosystems. Ecol. Lett. 10, 1135–1142 (2007).
- W. S. Harpole *et al.*, Nutrient co-limitation of primary producer communities. *Ecol. Lett.* 14, 852–862 (2011).
- C. E. Farrior et al., Resource limitation in a competitive context determines complex plant responses to experimental resource additions. Ecology 94, 2505–2517 (2013).

and Forest Sciences, Agricultural University of Iceland, Reykjavík 112, Iceland; ^pDepartment of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, IA 50011; ^qCo-Centre for Climate + Biodiversity + Water, School of Natural Sciences, Trinity College Dublin, Dublin 2, Ireland; ^rCenter for Applied Ecology "Prof. Baeta Neves"-Research Network in Biodiversity and Evolutionary Biology, School of Agriculture, University of Lisbon, Lisbon 1349-017, Portugal; ^sForest Research Centre, Associate Laboratory TERRA, School of Agriculture, University of Lisbon, Lisbon 1349-017, Portugal; ^tDepartment of Geography, King's College London, London WC2B 2BG, United Kingdom, "Fenner School of Environment & Society, Australian National University, Canberra, ACT 2600, Australia; "Senckenberg Museum for Natural History Görlitz, Görlitz D-02826, Germany; "Ecology, Behavior & Evolution Department, University of California San Diego, La Jolla, CA 92103; "Department of Biology, University of New Mexico, Albuquerque, NM 87131; ^yInstituto de Investigaciones Marinas y Costeras, Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Mar del Plata-Consejo Nacional de Investigaciones Científicas y Técnicas, Nacional de Mar del Plata-Consejo Nacional de Investigaciones Clentificas y Tecnicas, Mar del Plata B7600WAG, Argentina; ²Desert Ecology Research Group, School of Life and Environmental Sciences, The University of Sydney, NSW 2006, Australia; ^{a3}Zoology, School of Natural Sciences, Trinity College Dublin, Dublin 2, Ireland; ^{bb}MPG Ranch, Missoula, MT 59833; ^{cc}Institute of Biology, Leipzig University, Leipzig 04103, Germany; ^{dd}Ecology and Genetics Unit, University of Oulu, Oulu FI-90014, Finland; ^{ee}Department of Zoology and Entomology, Mammal Research Institute, University of Pretoria, Pretoria, 2002(4, Couth Africa) [Department of Biology Leipzig Leipzig Leipzig Couth Africa)] 400364, South Africa; fDepartment of Biology, Ecology and Biodiversity Group, Utrecht University, Utrecht 3584 CH, The Netherlands; ^{BE}Department of Biology, University of North Carolina, Chapel Hill, NC 27599; ^{hh}Department of Integrative Biology, University of Texas at Austin, Austin, TX 78712; ^{li}Life and Environmental Sciences, University of Iceland, Reykjavik 102, Iceland; ^{II}Department of Health and Environmental Sciences, Xi'an Jiaotong-Liverpool University, Suzhou 215123, China; ^{Ikk}Department of Environmental Science and Policy, Marist College, Poughkeepsie, NY 12601; ^{II}Department of Biology, Texas State University, San Marcos, TX 78666; ^{mm}Department of Plant and Soil Sciences, University of Kentucky, Lexington, KY 40546; ^{mm}Department of Environment and Genetics, Version and Control (Control Control C La Trobe University, Bundoora, VIC 3083, Australia; ⁰⁰Department of Biology-Wildness, Biodiversity, and Ecosystems Under Change, Vrije Universiteit Brussel, Brussels 1050, Belgium; ppUniversidad Nacional de la Patagonia Austral-Instituto Nacional de Tecnología Agropecuaria-Consejo Nacional de la ratagona dustral-instituto fractiona de l'echologia Agropecuaria-Consejo Nacional de Investigaciones Científicas y Técnicas, Rio Gallegos CP 9400, Santa Cruz, Argentina; ^{qa}Hawkesbury Institute for the Environment, Western Sydney University, Penrith, NSW 2751, Australia; "Sorbonne Université, Université de Paris-Cité, Université Paris-Est Créteil, Institut de Recherche pour le Développement, Centre National de la Recherche Scientifique, Institut National de Recherche pour l'Agriculture, National de la Recherche Scientifique, Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement, Institut d'Ecologie et des Sciences de l'Envrionnement de Paris, Paris 75005, France; ⁵⁵College of Ecology, Lanzhou University, Lanzhou City 730000, China; ¹¹Gansu Gannan Grassland Ecosystem National Observation and Research Station, Maqu County 747300, Gansu Province, China; ¹¹⁰Department of Biology, Colorado State University, Fort Collins, CO 80523; ¹¹⁰Department of Soil and Environment, Swedish University of Agricultural Sciences, Uppsala 75007, Sweden; ¹¹¹Cancaster Environment Centre, Lancaster University, Lancaster LA1 4YQ, United Kingdom; ¹²²Centre for Functional Biodiversity, School of Life Sciences, University of KwaZulu-Natal, Pietermaritzburg 3209, South Africa; ¹¹²Chool of Life and Environmental Sciences ABC Training Centre in Data South Africa; ^{yy}School of Life and Environmental Sciences, ARC Training Centre in Data Analytics for Resources and Environments, The University of Sydney, Sydney, NSW 2006, Australia; and ^{zz}School of Biological Sciences, University of Nebraska-Lincoln, Lincoln, NE 68588

Author contributions: P.A.F., L.A.G., and L.Y. designed research; P.A.F., L.Y., P.B.A., J.D.B., E.T.B., W.S.H., E.H.-G., A.S.M., A.C.R., E.W.S., S. Bagchi, I.C.B., L.B., Y.M.B., M.N.B., M.C.C., J.A.C., Q.Q.C., E.E.C., S.L.C., P.D., C.R.D., I.D., M.E.D., N.E., A.E., N.H., Y.H., R.W.H., I.S.J., J.M.H.K., R.L., J.P.M., R.L.M., J.W.M., H.O.V., P.L.P., S.A.P., X.R., Z.R., C.R., M.D.S., M.S., C.J.S., M.J.T., R.V., G.M.W., and G.R.W. performed research; P.A.F., L.A.G., L.Y., and S. Bharath analyzed data; and P.A.F., L.A.G., L.Y., and T.E.H. wrote the paper with contributions from all authors.

The authors declare no competing interest.

This article is a PNAS Direct Submission. C.Z. is a guest editor invited by the Editorial Board.

Copyright © 2025 the Author(s). Published by PNAS. This article is distributed under Creative Commons Attribution License 4.0 (CC BY).

Although PNAS asks authors to adhere to United Nations naming conventions for maps (https://www.un.org/geospatial/mapsgeo), our policy is to publish maps as provided by the authors.

- 12. P. A. Fay et al., Grassland productivity limited by multiple nutrients. Nat. Plants 1, 15080 (2015).
- 13. J. Sardans, J. Peñuelas, Potassium: A neglected nutrient in global change. Global Ecol. Biogeogr. 24,
- 261-275 (2015).
 A. K. Knapp, J. Briggs, M. D. L. Childers, O. E. Sala, "Estimating aboveground net primaryproduction in grassland- and herbaceous-dominated ecosystems" in *Principles and Standards for Measuring Primary Production*, T. J. Fahey, A. K. Knapp, Eds. (Oxford University Press, New York, 2007), chap. 3, pp. 27-48.
- H. N. Le Houerou, Rain use efficiency: A unifying concept in arid-land ecology. J. Arid Environ. 7, 213–247 (1984).
- O. E. Sala, W. J. Parton, L. A. Joyce, W. K. Lauenroth, Primary production of the central grassland region of the United States. *Ecology* 69, 40–45 (1988).
- D. U. Hooper, L. C. Johnson, Nitrogen limitation in dryland ecosystems: Responses to geographical and temporal variation in precipitation. *Biogeochemistry* 46, 247–293 (1999).
- A. K. Knapp, M. D. Smith, Variation among biomes in temporal dynamics of aboveground primary production. *Science* 291, 481–484 (2001).
- T. E. Huxman *et al.*, Convergence across biomes to a common rain-use efficiency. *Nature* 429, 651–654 (2004).
- P. B. Adler, J. M. Levine, Contrasting relationships between precipitation and species richness in space and time. *Oikos* 116, 221–232 (2007).
- S. Del Grosso *et al.*, Global potential net primary production predicted from vegetation class, precipitation, and temperature. *Ecology* 89, 2117–2126 (2008).
- G. E. Ponce-Campos et al., Ecosystem resilience despite large-scale altered hydroclimatic conditions. Nature 494, 349–352 (2013).

- 23. A. K. Knapp, P. Ciais, M. D. Smith, Reconciling inconsistencies in precipitation-productivity relationships: Implications for climate change. New Phytol. 214, 41-47 (2017).
- A. S. MacDougall et al., Widening global variability in grassland biomass since the 1980s. Nat. Ecol. 24 Evol. 8, 1877-1888 (2024), 10.1038/s41559-024-02500-x.
- N. Mahowald et al., Global distribution of atmospheric phosphorus sources, concentrations and deposition rates, and anthropogenic impacts. Global Biogeochem. Cycles 22, GB4026 (2008).
- T. L. Greaver et al., Key ecological responses to nitrogen are altered by climate change. Nat. Clim. 26. Change 6, 836-843 (2016).
- D. Tilman, D. Wedin, J. Knops, Productivity and sustainability influenced by biodiversity in grassland 27. ecosystems. Nature 379, 718-720 (1996).
- 28 P. B. Reich. The world-wide 'fast-slow' plant economics spectrum: A traits manifesto. J. Ecol. 102. 275-301 (2014).
- W. S. Harpole et al., Addition of multiple limiting resources reduces grassland diversity. Nature 537 29 93-96 (2016).
- O. E. Sala, L. Yahdjian, K. Havstad, M. R. Aguiar, "Rangeland ecosystem services: Nature's supply and 30 humans' demand" in Rangeland Systems, D. D. Briske, Ed. (Springer, 2017), chap. 14, pp. 467-489, 10.1007/978-3-319-46709-2_14.
- M. Migliavacca et al., The three major axes of terrestrial ecosystem function. Nature 598, 468-472 31. (2021)
- 32. O. Carroll et al., Nutrient identity modifies the destabilising effects of eutrophication in grasslands. Ecol. Lett. 25, 754-765 (2022).
- J. J. Elser et al., Shifts in lake N: P stoichiometry and nutrient limitation driven by atmospheric 33 nitrogen deposition. Science 326, 835-837 (2009).
- D. S. Gruner et al., A cross-system synthesis of consumer and nutrient resource control on producer 34. biomass. Ecol. Lett. 11, 740-755 (2008).
- E. T. Borer, J. B. Grace, W. S. Harpole, A. S. MacDougall, E. W. Seabloom, A decade of insights into 35 grassland ecosystem responses to global environmental change. Nat. Ecol. Evol. 1, 0118 (2017).
- M. D. Smith, A. K. Knapp, S. L. Collins, A framework for assessing ecosystem dynamics in response to 36 chronic resource alterations induced by global change. Ecology 90, 3279-3289 (2009).
- 37 M. L. Avolio et al., Changes in plant community composition, not diversity, during a decade of nitrogen and phosphorus additions drive above-ground productivity in a tallgrass prairie. J. Ecol. **102**, 1649-1660 (2014).
- 38. S. E. Koerner et al., Nutrient additions cause divergence of tallgrass prairie plant communities resulting in loss of ecosystem stability. J. Ecol. 104, 1478-1487 (2016).
- Y. Hautier et al., Local loss and spatial homogenization of plant diversity reduce ecosystem multifunctionality. Nat. Ecol. Evol. 2, 50-56 (2018).
- M. L. Avolio et al., Demystifying dominant species. New Phytol. 223, 1106–1126 (2019).
 K. J. Komatsu et al., Global change effects on plant communities are magnified by time and the
- number of global change factors imposed. Proc. Natl. Acad. Sci. U.S.A. 116, 201819027 (2019). J. D. Bakker et al., Compositional variation in grassland plant communities. Ecosphere 14, e4542 42.
- (2023). 43. P. B. Adler, J. Hillerislambers, J. M. Levine, A niche for neutrality. Ecol. Lett. 10, 95-104 (2007).
- E. E. Cleland *et al.*, Sensitivity of grassland plant community composition to spatial vs. temporal variation in precipitation. *Ecology* **94**, 1687–1696 (2013). 44.
- 45 K. R. Wilcox, J. M. Blair, M. D. Smith, A. K. Knapp, Does ecosystem sensitivity to precipitation at the
- site-level conform to regional-scale predictions? Ecology 97, 561-568 (2016). S. Harrison, M. J. Spasojevic, D. J. Li, Climate and plant community diversity in space and time. Proc. 46 Natl. Acad. Sci. U.S.A. 117, 4464-4470 (2020).
- 47. M. D. Smith et al., Richness, not evenness, varies across water availability gradients in grassy biomes on five continents. Oecologia 199, 649-659 (2022).
- 48 K. R. Wilcox et al., Asynchrony among local communities stabilises ecosystem function of metacommunities. Ecol. Lett. 20, 1534-1545 (2017).
- A. Eskelinen, S. Harrison, M. Tuomi, Plant traits mediate consumer and nutrient control on plant 49. community productivity and diversity. Ecology 93, 2705-2718 (2012).
- M. B. Soons et al., Nitrogen effects on plant species richness in herbaceous communities are more widespread and stronger than those of phosphorus. *Biol. Conserv.* **212**, 390–397 50. (2017)
- D. Ward, K. Kirkman, Z. Tsvuura, An African grassland responds similarly to long-term fertilization to the Park Grass experiment. *PLoS One* 12, e0177208 (2017).
- 52 M. D. Smith, A. K. Knapp, Dominant species maintain ecosystem function with non-random species loss. Ecol. Lett. 6, 509-517 (2003).
- E. Ladouceur et al., Linking changes in species composition and biomass in a globally distributed 53 grassland experiment. Ecol. Lett. 25, 2699-2712 (2022).
- A. J. Muehleisen et al., Nutrient addition drives declines in grassland species richness primarily via enhanced species loss. J. Ecol. 111, 552-563 (2023).
- P. A. Wilfahrt et al., Nothing lasts forever: Dominant species decline under rapid environmental change in global grasslands. J. Ecol. 111, 2472-2482 (2023).
- Y. Hautier et al., Eutrophication weakens stabilizing effects of diversity in natural grasslands. Nature 56. 508, 521-525 (2014).
- Q. Chen et al., Multidimensional responses of grassland stability to eutrophication. Nat. Commun. 57. 14, 6375 (2023).
- E. T. Borer et al., Finding generality in ecology: A model for globally distributed experiments. 58. Methods Ecol. Evol. 5, 65-73 (2014).

- 59. Y. Yan et al., Land use intensity controls the diversity-productivity relationship in northern temperate grasslands of China. Front. Plant Sci. 14, 1296544 (2023).
- E. Tipping et al., Atmospheric deposition of phosphorus to land and freshwater. Environ. Sci. Process 60 Impacts 16, 1608-1617 (2014).
- M. Kanakidou et al., Past, present and future atmospheric nitrogen deposition. J. Atmos. Sci. 73, 2039-2047 (2016).
- 62. J. Sardans, J. Peñuelas, Potassium control of plant functions: Ecological and agricultural implications. Plants 10, 419 (2021).
- L. Yahdjian, L. Gherardi, O. E. Sala, Nitrogen limitation in arid-subhumid ecosystems: A meta-63. analysis of fertilization studies. J. Arid Environ. 75, 675–680 (2011).
- 64 S. Bharath et al., Nutrient addition increases grassland sensitivity to droughts. Ecology 101, e02981 (2020)
- O. E. Sala, L. A. Gherardi, L. Reichmann, E. Jobbagy, D. Peters, Legacies of precipitation fluctuations 65 on primary production: Theory and data synthesis. Philos. Trans. R. Soc. Lond. B 367, 3135-3144 (2012).
- 66. H. Yang, S. B. Rood, L. B. Flanagan, Controls on ecosystem water-use and water-use efficiency: Insights from a comparison between grassland and riparian forest in the northern Great Plains. Agric. Forest Meteorol. 271, 22-32 (2019).
- J.A. Langley, B.A. Hungate, Plant community feedbacks and long-term ecosystem responses to multi-factored global change. AoB Plants 6, plu035 (2014).
- P. A. Fay et al., Multiple constraints cause positive and negative feedbacks limiting grassland soil CO2 efflux under CO2 enrichment. Proc. Natl. Acad. Sci. U.S.A. 118, e2008284117 (2020), 10.1073/ pnas.2008284117.
- D. Tilman, J. A. Downing, Biodiversity and stability in grasslands. Nature 367, 363-365 (1994). 69.
- Y. Hautier *et al.*, General destabilizing effects of eutrophication on grassland productivity at multiple spatial scales. *Nat. Commun.* **11**, 5375 (2020). 70.
- D. R. Chalcraft et al., Scale-dependent responses of plant biodiversity to nitrogen enrichment. 71. Ecology 89, 2165-2171 (2008).
- E. W. Seabloom et al., Increasing effects of chronic nutrient enrichment on plant diversity loss and 72 ecosystem productivity over time. Ecology 102, e03218 (2021).
- Q. Chen et al., Change in functional trait diversity mediates the effects of nutrient addition on 73. grassland stability. J. Ecol. 112, 2598-2612 (2024).
- 74 M. W. Cadotte, K. Carscadden, N. Mirotchnick, Beyond species: Functional diversity and the maintenance of ecological processes and services. J. Appl. Ecol. 48, 1079-1087 (2011).
- O. L. Petchey, A. Hector, K. J. Gaston, How do different measures of functional diversity perform? Ecology 85, 847-857 (2004).
- L. W. Walker, J. K. Syers, The fate of phosphorus during pedogenesis. *Geoderma* 15, 1–19 (1976).
 C. L. Turner, J. M. Blair, R. J. Schartz, J. C. Neel, Soil N and plant responses to fire, topography, and 76
- 77
- supplemental N in tallgrass prairie. *Ecology* **78**, 1832–1843 (1997). H. E. Epstein, W. K. Lauenroth, I. C. Burke, Effects of temperature and soil texture on ANPP in the U.S. 78. Great Plains. Ecology 78, 2628–2631 (1997).
- 79. P. B. Hook, I. C. Burke, Biogeochemistry in a shortgrass landscape: Control by topography, soil texture, and microclimate. Ecology 81, 2686-2703 (2000).
- J. Kaye, J. Barrett, I. Burke, Stable nitrogen and carbon pools in grassland soils and variable texture 80 and carbon content. Ecosystems 5, 461-471 (2002).
- M. Spohn, Interactions of nitrogen and phosphorus in plant nutrition-Analysis of a 60-years old 81. field experiment. Plant Soil (2024), 10.1007/s11104-024-06920-3.
- 82 A. Eskelinen, S. P. Harrison, Resource colimitation governs plant community responses to altered precipitation. Proc. Natl. Acad. Sci. U.S.A. 112, 13009-13014 (2015).
- 83 K. J. La Pierre, D. M. Blumenthal, C. S. Brown, J. A. Klein, M. D. Smith, Drivers of variation in aboveground net primary productivity and plant community composition differ across a broad precipitation gradient. Ecosystems 19, 521-533 (2016).
- W. Borken, E. Matzner, Reappraisal of drying and wetting effects on C and N mineralization and fluxes in soils. Global Change Biol. 15, 808-824 (2009).
- S. Manzoni, J. P. Schimel, A. Porporato, Responses of soil microbial communities to water stress: 85. Results from a meta-analysis. Ecology 93, 930-938 (2012).
- L. Augusto, D. L. Achat, M. Jonard, D. Vidal, B. Ringeval, Soil parent material-A major driver of plant 86. nutrient limitations in terrestrial ecosystems. Global Change Biol. 23, 3808-3824 (2017).
- 87 E. Du et al., Global patterns of terrestrial nitrogen and phosphorus limitation. Nat. Geosci. 13, 221-226 (2020).
- D. Tian et al., Global changes alter plant multi-element stoichiometric coupling. New Phytol. 221, 88 807-817 (2019).
- 89. E. Vázquez et al., The synergistic response of primary production in grasslands to combined nitrogen and phosphorus addition is caused by increased nutrient uptake and retention. Plant Soil 490, 371-385 (2023).
- H. Tuomisto, G. Žuquim, G. Cárdenas, Species richness and diversity along edaphic and climatic gradients in Amazonia. *Ecography* 37, 1034–1046 (2014).
- H. Tuomisto, A diversity of beta diversities: Straightening up a concept gone awry. Part 1. Defining 91. beta diversity as a function of alpha and gamma diversity. Ecography 33, 2-22 (2010).
- S. E. Fick, R. J. Hijmans, WorldClim 2: New 1-km spatial resolution climate surfaces for global land 92. areas. Int. J. Climatol. 37, 4302-4315 (2017).
- P.A. Fay et al., Data from: Interactions among nutrients govern the global grassland biomass, precipitation 93. relationship. Dryad. https://doi.org/10.5061/dryad.vdncjsz50. Deposited 14 January 2025.