scientific reports

OPEN



Enhanced understanding of nitrogen fixing bacteria through DNA extraction with polyvinylidene fluoride membrane

Agnieszka Kalwasińska^{®1⊠}, Igor Królikiewicz¹, Sushma Rani Tirkey¹, Attila Szabó^{2,3} & Sweta Binod Kumar¹

The rhizobiota, particularly nitrogen-fixing bacteria, play a crucial role in plant functioning by providing essential nutrients and defense against pathogens. This study investigated the diversity of nitrogen-fixing bacteria in a relatively understudied habitat: technosoils developed from industrial soda production. To analyze the bacterial diversity in the rhizosphere soils of wheat (Triticum aestivum L.) and aster (Tripolium pannonicum Jacq.), regions of the nifH gene were amplified and sequenced from the resident bacterial communities. A polyvinylidene fluoride (PVDF) membrane was employed for metagenomic DNA extraction, enhancing the detection of nitrogen-fixing bacteria. Prior to standard DNA extraction, an enrichment step was conducted in nitrogen-free JMV medium at 26 °C for 24 h, with a modification that replaced soil with the PVDF membrane. This approach enabled a more comprehensive analysis of the rhizosphere bacterial community, revealing that unique amplicon sequence variants (ASVs) in aster and wheat membrane samples accounted for a notable proportion of all ASVs in the dataset (8.5% and 23%, respectively) that were not captured using the standard method. Additionally, our findings demonstrated higher alpha diversity of nitrogen-fixing bacteria in the wheat rhizosphere compared to the aster rhizosphere. In wheat, the dominant genus was Insolitispirillum (38.80%), followed by unclassified genera within Gammaproteobacteria (9.76%) and Rhodospirillaceae (4.74%). In contrast, the aster rhizosphere was predominantly occupied by Azotobacter (95.69%).

Keywords Bacterial diversity, Nitrogen-fixing bacteria, *nif*H, PVDF membrane, Rhizosphere

Polyvinylidene fluoride (PVDF) is a hydrophobic, thermally stable material that possesses great mechanical strength. It is resistant to chemical corrosion, oxidation, and pollution and is low-cost to produce. These qualities make it a popular membrane material in various industrial fields^{1,2}. PVDF membranes are integral parts of seawater desalination, and wastewater microfiltration processes^{3,4}. However, biofilm formation contributes to the blocking of the pores, reduction of filtration efficiency, shortens the lifespan of the membrane and increases energy consumption in membrane technologies in wastewater treatment applications. It has been reported that the PVDF membrane possesses a high affinity towards various bacteria due to its high hydrophobicity, roughness, and the accumulation of trace organic nutrients at the solid-liquid interface^{5,6}.

The adhesion of bacteria to the membrane results from a combination of bacterial and material characteristics. Bacteria, like other microorganisms, have a charged cell wall due to acid-based groups on their surface. Their cell walls are mostly hydrophobic, a crucial factor in strong adherence to the material's hydrophobic surfaces⁷. However, hydrophobicity isn't the sole factor influencing adhesion. Surface charge, roughness, chemical composition, pH, temperature, and ionic strength of the membrane surface and surrounding environment can significantly impact bacterial attachment⁸.

The results of research by Kumar et al.⁵ indicate that the bacterial community recovered from the PVDF membrane dipped in sample water was more diverse compared to the community obtained directly from the same water samples. This discovery served as the foundation for developing distinct sampling tools i.e. membrane-

¹Department of Environmental Microbiology and Biotechnology, Faculty of Biological and Veterinary Sciences, Nicolaus Copernicus University in Toruń, Lwowska 1, Toruń 87-100, Poland. ²Department of Aquatic Sciences and Assessment, Swedish University of Agricultural Sciences, Uppsala 750 07, Sweden. ³Institute of Aquatic Ecology, HUN-REN Centre for Ecological Research, Karolina út 29, Budapest 1113, Hungary. ^{\Box}email: kala@umk.pl based kits in the form of strips for the detection of *E. coli*⁹ and *Vibrio* species (*V. cholerae*, *V. parahaemolyticus*, *V. campbellii*, *V. harveyi*, and *V. proteolyticus*) in water samples¹⁰.

In this study, we aimed to evaluate the effectiveness of PVDF membranes in the pre-treatment step to enhance bacterial detection in soil environments. DNA extraction from soil generally involves isolating genetic material from microorganisms within the soil matrix using a combination of physical, chemical, and enzymatic methods. Common approaches include direct methods, which lyse soil cells to maximize yield, and indirect methods, which separate cells before lysis to minimize contaminants, such as humic acids, that can interfere with downstream analyses¹¹. The use of a PVDF membrane, which concentrates bacteria and allows them to proliferate before conventional DNA extraction, may improve the detection of rare taxa present in small quantities within the soil. To assess bacterial diversity, amplicon sequencing of the *nif*H gene on the Illumina platform was employed. Our focus was specifically on nitrogen-fixing bacteria from the rhizosphere, which have been extensively studied for their unique ability to fix atmospheric nitrogen and convert it into a plant-available form, offering potential to reduce agriculture's chemical dependence¹²⁻¹⁴.

Recent microbiological research has been extensively conducted in northern Poland to investigate how environmental parameters influence microbial abundance, structure, diversity, and, in particular, specific guilds involved in the nitrogen cycle in saline soils affected by the soda industry¹⁵. Our findings indicated that salinity had less impact on microorganisms responsible for nitrogen fixation and denitrification compared to the nitrifying guild. Notably, the diversity of nitrogen-fixing communities had not been previously explored in this region. Therefore, a secondary goal of our study was to uncover the diversity of nitrogen-fixing bacteria in technosoils from Inowrocław. The research hypotheses were as follows: (i) Soil DNA extraction using a PVDF membrane during the enrichment step is more efficient, resulting in higher bacterial diversity compared to the standard method without enrichment. (ii) Nitrogen-fixing bacteria in soda-impacted technosoils form a unique community when compared to those in other saline soils.

Materials and methods

Sampling and experimental design

Aster (*Tripolium pannonicum* (Jacq.)), and wheat (*Triticum aestivum* L.) plants were collected in April 2023 from two distinct locations: a saline wasteland for Aster, and an arable field for wheat (Fig. 1).

The samples were obtained in Inowrocław, Poland, close to the soda lime repository ponds of the CIECH Soda company with four replicates for each species. Plants were placed in sterile polyethylene bags and stored in a portable travel refrigerator at 4 °C before laboratory analyses. The time from sampling to analysis did not exceed 90 min. Soil samples were collected from the root surface of each plant using a sterile spatula, and then transferred into sterile Eppendorf tubes for subsequent studies (four replicates per root system of the given species).

The experiment consisted of three stages: pretreatment, in which samples were enriched with bacterial cells; genomic DNA extraction; amplification and next-generation sequencing (NGS) analysis. Additionally, soil basic parameters, including ECe, pH, Corg, CaCO₃, Ntot, and C/N were characterized.

Soil characteristics

The soil samples for physicochemical analysis were dried and sieved through a 2.0 mm mesh screen and immediately stored at 4 $^{\circ}$ C for further analysis. Soil properties were analysed as described in Hulisz et al.¹⁶.



Fig. 1. Outlook of the sampling sites. Figure 1. A - saline wasteland site with aster (*Tripolium pannonicum* Jacq.), W - arable field site with wheat (*Triticum aestivum* L.).

Pretreatment, PVDF membrane preparation

Five grams of rhizosphere soil and four pieces of PVDF membrane were placed into 45 ml of sterile JMV medium in Simax bottles (100 mL) to enrich samples in nitrogen-fixing bacteria during the 24-hour incubation at 20 °C. The second set of bottles with 5 g of soil without any liquid served as a control.

The nitrogen-free JMV medium¹⁷ had the following composition (g L⁻¹): mannitol, 5.0; K₂HPO₄, 0.6; KH₂PO₄, 1.8; MgSO₄.7H₂O, 0.2; NaCl, 10; CaCl₂.2H₂O, 0.02; micronutrient solution 2 mL (CuSO₄.5H₂O, 0.04; ZnSO₄.7H₂O, 0.12; H₃BO₃, 1.40; Na₂MoO₄.2H₂O, 1.0; MnSO₄.H₂O, 1.175); bromothymol blue 2 mL (5 g L⁻¹ in 0.2 N KOH); vitamin solution 1 mL (biotin, 0.1 gL⁻¹). The pH was adjusted with KOH to 8.5 ± 0.2. The JMV medium was selected based on our prior experience, which demonstrated its effectiveness in the isolation of a diverse range of putative nitrogen-fixing bacteria from technogenic soil¹⁸. The JMV medium is highly selective for nitrogen-fixing bacteria due to several factors: a nitrogen-free environment, mannitol as a carbon source (which is favoured by nitrogen-fixing bacteria), an alkaline pH that supports exopolysaccharide production and stress resistance, essential micronutrients (especially molybdenum for nitrogenase activity), and salinity. This makes it particularly useful in saline or degraded soils, where other microbes may struggle due to osmotic stress, nutrient imbalances, or high pH^{17,18}.

A polyvinylidene fluoride (PVDF) solution in dimethyl formamide (20% w/w) was prepared through slow dissolution by heating at 45 °C overnight. Based on previous studies, this concentration was selected as the optimal one for maximizing bacterial attachment⁵. The viscous polymeric solution was then cast onto a sheet of non-woven polyester fabric (Novatexx 2470, Freudenberg Filtration Technologies, Weinheim, Germany), taped to a glass plate, and evenly spread using a 100 μ m casting roller (MeSep, Kraków, Poland). The membrane was immediately immersed in water and left to air-dry. Once dried, the membrane was cut into small pieces (~ 1 cm²), autoclaved, and prepared for use in the experiment.

Genomic DNA extraction

A volume of 250 μ l of the 10-fold soil suspension in JMV medium (control samples) or four pieces of PVDF membrane (membrane samples), retrieved from the JMV medium, were used for DNA extraction following the Qiagen protocol for soil samples (DNeasy PowerSoil Kit, Qiagen). These samples were designated as C_A or C_W (control aster and control wheat, respectively) and M_A or M_W (membrane aster and membrane wheat, respectively). The experiment was performed in four replicates for each of the two types of rhizosphere.

NGS analysis

Metataxonomic analysis of nitrogen-fixing biota was evaluated by amplifying the *nifH* gene with the following primers: 5' TGCGAYCCSAARGCBGACTC 3' (PolF forward, without overhangs) and 5' ATSGCCATCATYTCRCCGGA 3' (PolR reverse, without overhangs) according to Poly et al.¹⁹.

The libraries were prepared following the Illumina Support Centre (ISC) protocol with a minor modification $(2 \times \text{Phanta Max Master Mix}, \text{Vazyme Biotech}, \text{Nanjing City}, \text{China was applied instead of Kapa HifiHot Start Ready mix}). Sequencing was performed by the Biobank (University of Łódź, Poland) on a MiSeq platform (Illumina, San Diego, CA, USA) using MiSeq Reagent Kit v2 (500 cycles) in paired-end mode (2 × 250 bp).$

Data analysis of *nif*H gene amplicons with a yield of 1,626,305 raw reads was processed using the DADA2 R package²⁰based on the DADA2 Pipeline Tutorial 1.16^{21} . The filter and trim step was performed using the filterAndTrim command with maxN =0, maxEE = c(2,2), trimLeft = c(20,20), truncQ = 2, rm.phix = TRUE, compress = TRUE, multithread = FALSE parameters. The nifH dada2 v2.05 database²² has been used as a reference dataset in the taxonomy assignment. Data was normalized using subsampling to 20,896 reads per sample.

Statistical analyses

The rarefaction curves for the obtained reads, canonical correspondence analysis (CCA), principal coordinate analysis (PCoA), analysis of similarities (ANOSIM), and calculations of alpha diversity metrics, were done using vegan in R v. $4.1.0^{23}$. Differences in the α -diversity metrics between groups of samples were tested with analysis of variance (ANOVA) using Past v. 3.08^{24} . In CCA the nitrifying biota at the ASV level was used as the dependent variable. The DNA extraction method and plant species were used as independent variables. The permutation test was applied to test the significance of constraints in the CCA model. Venn diagrams were generated in R using the limma package.

Results

Rhizosphere samples and sequencing efficiency

The most crucial physicochemical properties of the rhizosphere soil are presented in Table S1. This soil is classified as Mollic gleysol¹⁶, originating from alluvial sands and sandy loams. It is influenced by aeolian processes and the influx of technogenic materials rich in iron and $CaCO_3$ from highly saline groundwater. The soil from the wasteland (aster rhizosphere) exhibited higher salinity (ECe 20.01 dS/m) compared to the soil from the arable field (wheat rhizosphere, ECe 0.86 dS/m). Both soils were characterized by a slightly alkaline pH (7.2–7.5), a high content of calcium carbonate (23.5 and 18.5 mg/g, respectively), and a similar C to N ratio (9–11). This ratio is typical for agricultural soils, indicating rapid organic matter mineralization.

Next-generation sequencing of amplicons from PCR products of the *nif*H gene resulted in 1,626,305 raw reads. Out of these, 424,033 reads were high-quality sequences. The mean read depth for the membrane samples was significantly higher at 65.6 Mbases (\pm 9.2 Mb) compared to the control samples at 51.0 Mbases (\pm 5.5 Mb) (t-test, t = 3.62, *p* = 0.005). The rarefaction curves of the *nif*H samples flattened at around 10,000 sequences in all types of samples, suggesting that high coverage of the sample's diversity was captured (Fig.S1).

Target	Variable	Df	Chi ²	F	Pr(>F)
nifH	DNA extraction method	1	0.225	0.839	0.686
	Plant species	1	0.987	3.685	0.001***
	Residual	11	2.947		

Table 1. Significance of parameters used in this study, affecting bacterial community composition inrhizosphere soil samples, as revealed by CCA. *p < 0.05; ** p < 0.01; ***p < 0.001.

.....



Fig. 2. Venn diagrams presenting the number of unique elements (ASVs) in the rhizosphere of aster (A) and wheat (W) in control (C) and membrane (M) samples.

Target			Sobs	Shannon-Wienner H'	Simpson 1-D
nifH	Aster	С	30 ± 16	1.8 ± 0.6	0.730 ± 0.175
		М	44 ± 27	1.9 ± 0.5	0.747 ± 0.162
	Wheat	С	143 ± 60	2.8 ±0.7	0.830 ± 0.093
		М	144 ± 47	2.9 ± 0.9	0.785 ± 0.192

Table 2. Alpha diversity metrics in rhizosphere soil samples. Differences between control and membrane samples were not significant (p > 0.05).

DNA extraction method and plant species as potential drivers of community structure in rhizosphere soil

The results obtained from CCA revealed that, among the two factors that might have influenced the structure of nitrogen-fixing bacteria in the rhizosphere, plant species composition had a significant impact (Table 1).

The method of DNA extraction did not affect the composition of the nitrogen-fixing bacteria. This was supported by the PCoA analysis (Fig. S2) and the ANOSIM results. PCoA showed a clear separation of samples based on plant species but no separation based on the DNA extraction method. The ANOSIM statistic was R -0.2 (p-value 0.971) for aster and R -0.1 (p-value 0.6) for wheat.

Venn diagrams (Fig. 2) unveiled distinctions in the number of amplicon sequence variants (ASVs) between control and PVDF membrane treatment for *nif*H gene amplicons.

The median value of the unique elements in aster membrane samples was significantly higher than in control samples (22 vs. 11; Mann-Whitney U = 98.5, p = 0.026). The number of unique elements in wheat rhizosphere samples did not differ between membrane samples and controls (mean value 84 vs. 81; t-test: t = 0.12, p = 0.903). Unique ASVs in aster and wheat membrane samples constituted 8.5% and 23%, respectively, of all ASVs in the dataset. Further analysis showed that there were no significant differences in diversity metrics in soil rhizosphere samples between the methods of DNA extraction (Table 2).

Structure of bacterial communities - nitrogen-fixing bacteria

At the phylum level, nitrogen-fixing bacteria were dominated by reads assigned to Pseudomonadota (Fig. 3; Table S2). The Aster rhizosphere exhibited a nearly complete dominance of this phylum compared to the wheat rhizosphere (median 99.91% versus 86.11%). Representatives of the phyla Thermodesulfobacteriota and Bacillota were present in both types of rhizosphere; however, they were more abundant in the wheat rhizosphere compared to the aster (median 8.20% versus 0.08%, respectively, and 0.87% versus 0.00%). Phylotypes assigned to Bacteroidota, Cyanobacteria, Verrucomicrobiota, Desulfobacterota, and Elusimicrobiota were exclusively present in the wheat rhizosphere.

In terms of class-level taxonomy, the majority of reads in aster samples belonged to Gammaproteobacteria (median value of 99.48%). In wheat rhizosphere samples, Alphaproteobacteria was the most abundant class (51.48%), with Desulfuromonadia ranking second (6.65%). Representatives of other top ten bacterial classes, namely Alphaproteobacteria, Betaproteobacteria, and Desulfuromonadia, were more abundant in the wheat rhizosphere compared to the aster rhizosphere. Phylotypes assigned to Bacteroidia, Clostridia, Desulfobulbia, and Desulfobacteria were exclusively detected in the wheat rhizosphere.

The rhizosphere of Aster was predominantly occupied by *Azotobacter* reads, accounting for a median value of 95.69%. The second most abundant genus was *Pseudomonas*, constituting 2.02% of the community. In the wheat rhizosphere, the most prevalent genus was *Insolitispirillum* with 38.80%, followed by unclassified ASVs within the class Gammaproteobacteria (9.76%) and the family *Rhodospirillaceae* (4.74%), respectively.

The rare, putative nitrogen-fixing bacteria, constituting less than 0.01% of the relative abundance in a given sample and found exclusively in the membrane samples, are presented in Table 3.

Discussion

Assessing nitrogen-fixing microbial community composition through amplicon sequencing of the nifH gene is a widely used technique in microbial ecology. However, several inherent challenges and limitations can affect the accuracy and reliability of the results. The most important encompass primer bias, PCR amplification bias, and sequencing errors^{25,26}.

Rare species are often present in very low abundances (< 0.01% RA), making their detection highly dependent on the sequencing depth. Insufficient sequencing depth may fail to capture these low-abundance taxa, leading to an incomplete picture of the community diversity²⁷. During PCR amplification, rare species may be disproportionately underrepresented compared to more abundant taxa. This is due to the stochastic nature of PCR, where low-abundance sequences are less likely to be amplified consistently across cycles, especially when starting from a small number of template molecules²⁸. The results of the sequencing of the *nif*H gene appeared successful in capturing the diversity of the putative nitrogen fixers. This indicates that the sequencing approach was well-designed and performed reliably. The method of DNA isolation, as indicated by CCA, PCoA, ANOSIM, and analysis of alpha diversity metrics, did not significantly affect the overall community composition.



Fig. 3. Structure of the putative nitrogen-fixing bacterial communities at the phylum (**a**), class (**b**) and genus (**c**) levels in the rhizosphere of aster (A) and wheat (W) in control (C) and membrane (M) samples.

Target	ASV	Taxonomy
Aster	ASV402	Parazoarcus
	ASV455	unclassified Hydrogenophilales
	ASV467	Geobacter
	ASV487	unclassified Geminicoccaceae
	ASV522	Caenispirillum
	ASV143	Halorhodospira
	ASV366	Parazoarcus
	ASV469	unclassified Anaerolineae
Wheat	ASV435	Leptolyngbya
	ASV461	Citrifermentans
	ASV470	Anabaena
	ASV474	Geobacter
	ASV477	unclassified Desulfuromonadaceae
	ASV493	unclassified Chromatiales
	ASV496	Geobacter
	ASV524	unclassified Elusimicrobiota
	ASV525	unclassified Terasakiellaceae
	ASV411	Methyloferula
	ASV441	Candidatus 'Dadabacteria'
	ASV491	Methylocystis
	ASV492	Desulfovibrio
	ASV513	Desulfopila
	ASV515	Desulfovibrio
	ASV516	Geobacter
	ASV517	Geobacter

Table 3. Rare (< 0.01% RA) nitrogen-fixing bacteria exclusively present in membrane samples of aster and wheat in Inowrocław technosoil.

.....

However, Venn diagrams revealed notable differences in the number of bacterial ASVs uniquely detected in samples where metagenomic DNA was isolated using a membrane. In wheat membrane samples, unique ASVs accounted for 23% of all ASVs in the dataset, suggesting that the PVDF membrane method enabled the detection of a substantial portion of the bacterial community not captured by the standard method. Similarly, in aster membrane samples, unique ASVs constituted 8.5% of all ASVs. Although less pronounced than in wheat samples, this still highlights the membrane method's ability to uncover additional bacterial diversity compared to conventional approaches.

Alpha diversity metrics, such as the Shannon and Simpson indices, evaluate both the richness and evenness of taxa²⁹. If the unique ASVs detected in membrane samples are primarily rare and of low abundance, their contribution to these metrics may be minimal. In contrast, control samples with a more even distribution of taxa, despite having fewer unique ASVs, could exhibit similar or even higher alpha diversity values. In this study, both membrane and control samples were dominated by a few highly abundant taxa, which likely constrained the ability of alpha diversity metrics to capture differences arising from rare, unique ASVs. Consequently, the additional diversity detected in membrane samples may have been masked by the dominance of a few taxa. Furthermore, both CCA and PCoA analyses are heavily influenced by the most abundant taxa, as these methods aim to explain the largest variation in the dataset³⁰. Rare taxa, which contribute minimally to overall variation, have a limited impact on ordination results. This limitation is common in community ecology, as rare taxa typically represent a small fraction of the data matrix and do not significantly influence the patterns highlighted by ordination methods.

It is well established that nonspecific interactions or without the involvement of any particular stereospecific macromolecular binding sites mediate the attachment between membrane and bacteria. Therefore, the attachment is not substrate-specific which leads to the easy attachment of various kinds of bacteria on the membrane surface³¹. The deposition and accumulation of microorganisms onto the membrane surface involves a series of mechanisms causing irreversible attachment of the bacteria at the surface and inside the pores of the membrane³². Secondly, the amount of samples collected for any microbial studies comprises a very small amount which might not represent all types of bacteria⁵. Therefore, the membrane helps to concentrate/enrich different types of bacteria from the surrounding sample. Subsequently, it results in the isolation of bacteria according to several studies conducted previously^{5,33}. In those specific studies, the PVDF (20% W/W) membrane effectively and successfully isolated diversified bacteria from seawater samples. However, the studies were based on biochemical methods and identification (16 S rRNA) of bacteria recovered from the membrane compared to the surrounding water samples. The present study was an approach to validate those findings with the help

of molecular-based studies and whether this applies to soil samples too. Interestingly, the results of the present study accompany the fact that membranes especially PVDF (20% W/W) can concentrate diverse bacteria from the soil samples.

Plants profoundly influence the soil microbiome through root exudates, which are complex mixtures of sugars, amino acids, carboxylic acids, and secondary metabolites. These compounds serve as signalling molecules, attractants, stimulants, or inhibitors that shape the microbial community structure. The rhizobiome, or the microbial community in the rhizosphere, is shaped by plant-specific exudates, whose composition is under genetic control and varies across plant species^{34,35}. Additionally, the rhizobiome is influenced by soil type and geographical distance³⁶.

Aster, a halophytic plant, thrives in natural and anthropogenic environments near saline water sources, such as coastal areas or soda factory waste ponds^{37,38}. Halophytes like aster possess salinity-responsive genes and proteins that enable survival in high-salt environments. Their rhizospheres host plant growth-promoting rhizobacteria (PGPR) which enhance salinity tolerance and mitigate environmental stress³⁹.

In saline technogenic soils, the nitrogen-fixing bacterial community in the aster rhizosphere was characterized by a predominance of halotolerant bacteria, including members of the *Azotobacter* genus. These bacteria exhibit adaptations such as increased energy capacity to withstand saline conditions, which otherwise suppress microbial biomass and nitrogenase activity^{40,41}. Previous studies from the same area identified a halotolerant strain, *Azotobacter chroococcum*W4ii, with plant growth-promoting properties, suggesting a vital role for such bacteria in supporting aster growth in saline technogenic soils¹⁸. Rare putative nitrogen-fixing bacteria, including *Parazoarcus*⁴², *Geobacter*⁴³, and *Halorhodospira*⁴⁴, were identified using the membrane-based approach. Despite their low abundance, these bacteria appear to be key members of the aster rhizosphere community, given their documented roles in nitrogen fixation and adaptation to saline environments.

Wheat, a crop with moderate sensitivity to salinity, experiences significant reductions in growth and yield under saline conditions. This sensitivity arises from ionic and osmotic stress that disrupts water uptake and nutrient balance⁴⁵. Various strategies, including salt-tolerant endophytes and genetic modifications, have been employed to enhance wheat's salinity tolerance^{46,47}. Wheat requires a balanced supply of nutrients, particularly nitrogen, for optimal growth. Over-application of nitrogen fertilizers, a common practice in wheat cultivation, alters soil microbial activity and ecological functions⁴⁸. This dynamic impacts the rhizosphere microbiome, with wheat root exudates potentially attracting diverse nitrogen-fixing bacteria to compensate for nitrogen deficiencies⁴⁹.

The nitrogen-fixing bacterial communities in the rhizospheres of aster and wheat differed significantly. The aster rhizosphere, dominated by halotolerant Azotobacter and Pseudomonas species, reflected its adaptation to high-salinity environments. In contrast, wheat exhibited a broader spectrum of nitrogen-fixing bacteria, likely influenced by its nutrient demands and sensitivity to saline stress. This distinction underscores the role of root exudation in shaping rhizosphere composition, though further research is needed to elucidate how domestication and metabolic specialization drive these interactions⁵⁰. Among the nitrogen-fixing bacteria in wheat rhizosphere soil, Insolitispirillum emerged as a dominant genus. Though relatively understudied⁵¹, its prevalence suggests a significant ecological role within the *Rhodospirillaceae* family, which includes many known nitrogen-fixing species⁵². Rare taxa in wheat rhizosphere revealed with the improved detection method assigned to the genera Anabaena and Leptolyngbya are both well-known diazotrophic cyanobacteria⁴⁷ that convert atmospheric nitrogen into ammonium, making it available for plant uptake. Their ability to form biofilms may also help to retain moisture and nutrients in saline soils. Additionally, Methyloferula and Methylocystis, methane-oxidizing bacteria capable of nitrogen fixation⁴⁸, were identified. These bacteria may play a dual role in the rhizosphere by contributing to both nitrogen and carbon cycling, further supporting soil health and plant growth. At the broader taxonomic level, Gammaproteobacteria and Alphaproteobacteria were the dominant diazotrophic bacteria in coastal saline soils⁴⁷, a trend mirrored in the rhizospheres of both plants. However, variations in salinity levels significantly influenced community structure. For instance, higher salinity in our samples (approximately three times higher than coastal saline soils) likely shifted diazotrophic bacterial dominance, as observed in the relative abundance of Betaproteobacteria and Burkholderia in Polish arable soils⁴⁸.

Conclusions

The findings of this study confirm that PVDF membranes are an effective tool for gaining insights into nitrogenfixing communities and rare taxa. The interaction between plants and their rhizosphere microbiomes is highly dynamic, shaped by root exudates, soil conditions, and environmental stressors. In saline soils, aster supports a specialized, halotolerant nitrogen-fixing community, while wheat relies on a more diverse set of nitrogen-fixing bacteria to meet its nutrient needs.

The presence of rare nitrogen fixers, including cyanobacteria and methane-oxidizing bacteria, highlights the complex ecological interactions that sustain plant health and soil fertility. Further research is needed to understand how plant-specific factors, domestication, and environmental pressures influence the assembly of these microbial communities. Future studies should also focus on optimizing sequencing and analytical approaches to better integrate the contributions of rare taxa into broader ecological analyses.

PVDF membranes offer significant advantages for microbial diversity studies, particularly in challenging environments. Their unique properties, such as high chemical resistance and durability, may enable efficient microbial cell capture in extreme conditions, including highly saline and alkaline soils.

Data availability

Sequencing data were deposited in NCBI under accession number PRJNA1049880 (https://www.ncbi.nlm.nih. gov/bioproject/PRJNA1049880/).

Received: 17 January 2025; Accepted: 25 April 2025 Published online: 08 May 2025

References

- Ji, J., Liu, F., Hashim, N. A., Abed, M. M. & Li, K. Poly (vinylidene fluoride) (PVDF) membranes for fluid separation. *React. Funct. Polym.* 86, 134–153 (2015).
- 2. Ruan, L. et al. Properties and applications of the B phase poly(vinylidene fluoride). Polymers 10, 228 (2018).
- 3. Dallaev, R. et al. Brief review of PVDF properties and applications potential. Polymers 14 (22), 4793 (2022).
- 4. Xiang, S. et al. Fabrication of PVDF/EVOH blend Hollow fiber membranes with hydrophilic property via thermally induced phase process. Sep. Purif. Technol. 301, 122031 (2022).
- Kumar, S. B., Sharnagat, P., Manna, P., Bhattacharya, A. & Haldar, S. Enhanced bacterial affinity of PVDF membrane: its application as improved sea water sampling tool for environmental monitoring. *Environ. Sci. Pollut Res. Int.* 24 (6), 5831–5840 (2017).
- Zhao, G. & Chen, W. N. Biofouling formation and structure on original and modified PVDF membranes: role of microbial species and membrane properties. RSC Adv. 7, 37990–38000 (2017).
- 7. Krasowska, A. & Sigler, K. How microorganisms use hydrophobicity and what does this mean for human needs? Front. Cell. Infect. Microbiol. 4, 112 (2014).
- 8. AlSawaftah, N., Abuwatfa, W., Darwish, N. & Husseini, G. A. A review on membrane biofouling: prediction, characterization, and mitigation. *Membr* 12 (12), 1271 (2022).
- 9. Kumar, S. B., Shinde, A. H., Mehta, R., Bhattacharya, A. & Haldar, S. Simple, one-step dye-based kit for bacterial contamination detection in a range of water sources. *Sens. Actuators Chem.* 276, 121–127 (2018).
- Kumar, S. B., Shinde, A. H., Behere, M. J., Italia, D. & Haldar, S. Simple, rapid and on spot dye-based sensor for the detection of Vibrio load in shrimp culture farms. Arch. Microbiol. 203(6), 3525–3532 (2021). (2021).
- 11. Wydro, U. & Soil Microbiome Study Based on DNA Extraction. Rev. Water, 14(24), 3999 (2022).
- 12. 12, Kuan, K. B., Othman, R., Abdul Rahim, K. & Shamsuddin, Z. H. Plant growth-promoting rhizobacteria inoculation to enhance vegetative growth, nitrogen fixation and nitrogen remobilisation of maize under greenhouse conditions. *PLoS One* 24,11(3), e0152478 (2016).
- 13. Singh, R. K. et al. Diversity of nitrogen-fixing rhizobacteria associated with sugarcane: a comprehensive study of plant-microbe interactions for growth enhancement in *Saccharum* spp. *BMC Plant. Biol.* **20**, 1–21 (2020).
- 14. Shi, Z. et al. Screening of high-efficiency nitrogen-fixing bacteria from the traditional Chinese medicine plant *Astragalus mongolicus* and its effect on plant growth promotion and bacterial communities in the rhizosphere. *BMC Microbiol.* **23** (1), 292 (2023).
- 15. Kalwasińska, A. et al. Technogenic soil salinisation, vegetation, and management shape microbial abundance, diversity, and activity. Sci. Total Environ. 905, 167380 (2023).
- 16. Hulisz, P., Pindral, S., Kobierski, M. & Charzynski, P. Technogenic layers in organic soils as a result of the impact of the soda industry. *Eurasian Soil. Sci.* 51, 1133–1141 (2018).
- Baldani, J. I., Reis, V. M., Videira, S. S., Boddey, L. H. & Baldani, V. L. D. The Art of isolating nitrogen-fixing bacteria from nonleguminous plants using N-free semi-solid media: a practical guide for microbiologists. *Plant. Soil.* 384, 413–431 (2014).
- Binod Kumar, S., Kalwasińska, A., Brzezinska, M. S. & Wróbel, M. Application of halotolerant Azotobacter chroococcum W4ii isolated from technosoils to mitigate salt stress in wheat plant. Open Res. Eur. 3(76), 76 (2023). (2023).
- Poly, F., Monrozier, L. J. & Bally, R. Improvement in the RFLP procedure for studying the diversity of NifH genes in communities of nitrogen fixers in soil. Res. Microbiol. 152 (1), 95–103 (2001).
- 20. Callahan, B. J. et al. DADA2: High-resolution sample inference from illumina amplicon data. Nat. Methods. 13 (7), 581-583 (2016).
- 21. DADA2 Pipeline Tutorial 1.16. https://benjjneb.github.io/dada2/tutorial.html
- Moynihan, M. A. & Reeder, C. F. moyn413/nifHdada2: v2.0.5. Zenodo, (2023). https://zenodo.org/records/7996213
- 23. Oksanen, J. et al. Vegan: Community Ecology Package. R package version 2.5-6 (2019). https://github.com/vegandevs/vegan
- 24. Hammer, Ø. Paleontological statistics software package for education and data analysis. Palaeontol. Electron. 4, 9 (2001).
- Poretsky, R., Rodriguez-R, L. M., Luo, C., Tsementzi, D. & Konstantinidis, K. T. Strengths and limitations of 16S rRNA gene amplicon sequencing in revealing Temporal microbial community dynamics. *PloS One* 9(4), e93827 (2014).
- Smith, D. P. & Peay, K. G. Sequence depth, not PCR replication, improves ecological inference from next generation DNA sequencing. *PloS One* 9(2), e90234 (2014).
- 27. Pinto, A. J. & Raskin, L. PCR biases distort bacterial and archaeal community structure in pyrosequencing datasets. *PLoS One*. 7 (8), e43093 (2012).
- 28. Chiu, C. H. & Chao, A. Estimating and comparing microbial diversity in the presence of sequencing errors. PeerJ 1, 4:e1634 (2016).
- 29. Verniest, F. & Greulich, S. Methods for assessing the effects of environmental parameters on biological communities in long-term ecological studies A literature review. *Ecol. Model.* **414**, 108732 (2019).
- 30. Mallevialle, J., Odendaal, P. E. & Wiesner, M. R. Water Treatment Membrane Processes (McGraw-Hill, 1996).
- Rosales, A. B. et al. Minimizing bacterial adhesion on membrane: multiscale characterization of surface modifications. J. Membr. Sci. 684, 121867 (2023).
- 32. Kumar, S. B., Shinde, A. H., Mehta, R., Bhattacharya, A. & Haldar, S. Simple, one-step dye-based kit for bacterial contamination detection in a range of water sources. *Sens. Actuators B Chem.* **276**, 121–127 (2028).
- Kumar, S. B., Trivedi, H., Baraiya, N. R. & Haldar, S. An improved device with an affinity membrane to collect depth specific contamination free water for environmental assessment. *Analyst* 143(3), 662–669 (2018). (2018).
- 34. Hu, L. et al. Root exudate metabolites drive plant-soil feedbacks on growth and defense by shaping the rhizosphere microbiota. *Nat. Commun.* 9 (1), 2738 (2018).
- 35. Zhao, X. et al. Introduction of *Panax notoginseng* into pine forests significantly enhances the diversity, stochastic processes, and network complexity of nitrogen-fixing bacteria in the soil. *Front Microbiol.*;16:1531875 (2025). (2025).
- Qin, K., Dong, X., Jifon, J. & Leskovar, D. I. Rhizosphere microbial biomass is affected by soil type, organic and water inputs in a bell pepper system. *Appl. Soil. Ecol.* 138, 80–87 (2019).
- Lazarus, M. & Wszałek-Rożek, K. Two rare halophyte species: Aster tripolium L. and Plantago maritima L. on the Baltic Coast in Poland-their resources, distribution and implications for conservation management. *Biodiv Res. Conserv.* 41(1), 51–60 (2016). (2016).
- Karasińska, W., Nienartowicz, A., Kunz, M., Kamiński, D. & Piernik, A. Resources and dynamics of halophytes in agricultural and industrial landscapes of the Western part of Kujawy, central Poland. *Ecol. Quest.* 32 (4), 7–25 (2021).
- Etesami, H. & Glick, B. R. Halotolerant plant growth-promoting bacteria: Prospects for alleviating salinity stress in plants. *Environ. Exp. Bot.* 178, 104124 (2020). (2020).
- Severin, I., Confurius-Guns, V. & Stal, L. J. Effect of salinity on nitrogenase activity and composition of the active diazotrophic community in intertidal microbial Mats. Arch. Microbiol. 194 (6), 483–491 (2012).
- Rath, K. M., Maheshwari, A., Bengtson, P. & Rousk, J. Comparative toxicities of salts on microbial processes in soil. Appl. Environ. Microbiol. 82 (7), 2012–2020 (2016).

- 42. Boden, R., Hutt, L. P. & Rae, A. W. Reclassification of *Thiobacillus aquaesulis* (Wood & Kelly, 1995) as *Annwoodia aquaesulis* gen. Nov., comb. Nov., transfer of *Thiobacillus* (Beijerinck, 1904) from the *Hydrogenophilales* to the Nitrosomonadales, proposal of *Hydrogenophilalia* class. Nov. Within the 'proteobacteria', and four new families Within the orders *Nitrosomonadales* and *Rhodocyclales. Int. J. Syst. Evol. Microbiol.* **67** (5), 1191–1205 (2017).
- Ortiz-Medina, J. F., Poole, M. R., Grunden, A. M. & Call, D. F. Nitrogen fixation and ammonium assimilation pathway expression of *Geobacter sulfurreducens* changes in response to the anode potential in microbial electrochemical cells. *Appl. Environ. Microbiol.* 26 (4), e0207322 (2023).
- Tsuihiji, H., Yamazaki, Y., Kamikubo, H., Imamoto, Y. & Kataoka, M. Cloning and characterization of nif structural and regulatory genes in the purple sulfur bacterium, *Halorhodospira halophila. J. Biosci. Bioeng.*; 101, 263–270 (2006). (2006).
- 45. Duarte-Delgado, D. et al. Transcriptome profiling at osmotic and ionic phases of salt stress response in bread wheat uncovers traitspecific candidate genes. *BMC Plant. Biol.* 16 (20(1)), 428 (2020). (2020).
- 46. Bouzouina, M., Kouadria, R. & Lotmani, B. Fungal endophytes alleviate salt stress in wheat in terms of growth, ion homeostasis and osmoregulation. J. Appl. Microbiol. 130 (3), 913–925 (2021).
- Yu, Y., Wu, Y. & He, L. A wheat WRKY transcription factor TaWRKY17 enhances tolerance to salt stress in Transgenic Arabidopsis and wheat plant. *Plant. Mol. Biol.* 113 (4–5), 171–191 (2023).
- 48. Zhou, J. et al. Consistent effects of nitrogen fertilization on soil bacterial communities in black soils for two crop seasons in China. *Sci. Rep.* **7** (1), 3267 (2017).
- Yang, Y. et al. Effect of nitrogen management on wheat yield, water and nitrogen utilization, and economic benefits under ridgefurrow cropping system with supplementary irrigation. Agronomy 13 (7), 1708 (2023).
- 50. Yue, H. et al. Plant domestication shapes rhizosphere Microbiome assembly and metabolic functions. *Microbiome* 11 (1), 70 (2023).
- Koziaeva, V. V., Sorokin, D. Y., Kolganova, T. V. & Grouzdev, D. S. Magnetospirillum sulfuroxidans Sp. nov., capable of sulfurdependent lithoautotrophy and a taxonomic Reevaluation of the order *Rhodospirillales*. Syst. Appl. Microbiol. 46 (3), 126406 (2023).
- Madigan, M., Cox, S. S. & Stegeman, R. A. Nitrogen fixation and nitrogenase activities in members of the family. *Rhodospirillaceae J. Bacteriol.* 157, 73–78 (1984).
- 53. Charpy, L. et al. Dinitrogen-fixing cyanobacteria in microbial Mats of two shallow coral reef ecosystems. *Microb. Ecol.* **59** (1), 174–186 (2010).
- 54. Mäkipää, R. et al. Methanotrophs are core members of the Diazotroph community in decaying Norway Spruce logs. *Soil. Biol. Biochem.* **120**, 230–232 (2018).
- 55. Song, Y. et al. The diversity and structure of diazotrophic communities in the rhizosphere of coastal saline plants is mainly affected by soil physicochemical factors but not host plant species. *Front. Mar. Sci.* **9**, 1100289 (2022).
- Wolińska, A. et al. Metagenomic analysis of some potential nitrogen-fixing bacteria in arable soils at different formation processes. *Microb. Ecol.* 73, 162–176 (2017).

Acknowledgements

This study received funding from the European Union's Horizon 2020 Research and Innovation Programme (Grant agreement 101038072) and EF OBSIDIAN Soil Science, Microbiology, Agricultural Genetics, and Food Quality https://soil-micro.umk.

Author contributions

AK - conceptualization, investigation, funding acquisition, writing - original draft, methodology, validation, visualization, writing - review and editing, supervision, resources, project administration; IK -writing - original draft, software, formal analysis, data curation; SRT - formal analysis, writing - original draft; AS -formal analysis, writing - original draft; SBK -methodology, writing - original draft.

Declarations

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at https://doi.org/1 0.1038/s41598-025-00173-5.

Correspondence and requests for materials should be addressed to A.K.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2025