

Commodity risk assessment of *Alnus cordata* and *Alnus glutinosa* specimen trees from the UK

EFSA Panel on Plant Health (PLH) | Antonio Vicent Civera | Paula Baptista | Anna Berlin | Elisavet Chatzivassiliou | Jaime Cubero | Nik Cunniffe | Eduardo de la Peña | Nicolas Desneux | Francesco Di Serio | Anna Filipiak | Paolo Gonthier | Beata Hasiów-Jaroszewska | Hervé Jactel | Blanca B. Landa | Lara Maistrello | David Makowski | Panagiotis Milonas | Nikos T. Papadopoulos | Hanna Susi | Dirk Jan van der Gaag | Jane Debode | Charles Manceau | Christer Sven Magnusson | Juan A. Navas-Cortes | Christos Kritikos | Maria Kormpi | Dimitrios Papachristos | Chrysavgi Reppa | Olaf Mosbach Schulz | Ciro Gardi | Monia Federica Lombardo | Antigoni Akrivou | Spyridon Antonatos | Despoina Beris | Roel Potting

Correspondence: plants@efsa.europa.eu

The declarations of interest of all scientific experts active in EFSA's work are available at <https://open.efsa.europa.eu/experts>

Abstract

The European Commission requested the EFSA Panel on Plant Health to prepare and deliver risk assessments for commodities listed in Commission Implementing Regulation (EU) 2018/2019 as 'high risk plants, plant products and other objects'. This Scientific Opinion covers the plant health risks posed by the following commodities: *Alnus cordata* and *A. glutinosa* as specimen trees (from 7 to 25 years old) in pots imported into the EU from the UK. A list of pests potentially associated with the commodities was compiled. The relevance of each pest was assessed based on evidence following defined criteria. Three pests were selected for further evaluation: one EU-protected zone quarantine pest (*Entoleuca mammata*), one EU quarantine pest (*Phytophthora ramorum* (non-EU isolates)) and one non-quarantine pest (*Phytophthora siskiyouensis*). For the selected pests, the risk mitigation measures implemented in the UK and specified in the technical dossier were evaluated. For these pests, an expert judgement is given on the likelihood of pest freedom taking into consideration the risk mitigation measures acting on the pest, including uncertainties associated with the assessment. In the assessment of risk, the age of the plants was considered, as larger trees are more likely to be infested mainly due to longer time grown in the field. In addition, large canopies and root systems are more difficult to inspect, thereby making the detection of pests more challenging on large trees. The degree of pest freedom varies between the pests evaluated, with *E. mammata* being the pest most frequently expected on imported *Alnus* spp. specimen trees. Expert Knowledge Elicitation indicated, with 95% certainty, that between 9905 and 10,000 per 10,000 *Alnus* spp. specimen trees would be free from *E. mammata*.

KEY WORDS

alder, *Alnus*, commodity risk assessment, European Union, plant health, plant pests

CONTENTS

Abstract.....	1
1. Introduction	3
1.1. Background and Terms of Reference as provided by European Commission	3
1.1.1. Background	3
1.1.2. Terms of Reference.....	3
1.2. Interpretation of the Terms of Reference	3
2. Data and Methodologies.....	5
2.1. Data provided by DEFRA of the UK	5
2.2. Literature searches performed by EFSA.....	6
2.3. Methodology.....	6
2.3.1. Commodity data.....	7
2.3.2. Identification of pests potentially associated with the commodity	7
2.3.3. Listing and evaluation of risk mitigation measures	7
2.3.4. Expert Knowledge Elicitation.....	8
3. Commodity Data	8
3.1. Description of the commodity.....	8
3.2. Description of the production areas.....	9
3.3. Production and handling processes	11
3.3.1. Growing conditions.....	11
3.3.2. Source of planting material	11
3.3.3. Production cycle	11
3.3.4. Post-harvest processes and export procedure.....	13
4. Identification of Pests Potentially Associated With the Commodity	13
4.1. Selection of relevant EU-quarantine pests associated with the commodity.....	13
4.2. Selection of other relevant pests (non-quarantine in the EU) associated with the commodity.....	15
4.3. List of potential pests not further assessed.....	15
4.4. Summary of pests selected for further evaluation	15
5. Risk Mitigation Measures	15
5.1. Risk mitigation measures applied in the UK.....	16
5.2. Evaluation of the current measures for the selected pests including uncertainties	16
5.2.1. Overview of the evaluation of <i>Entoleuca mammata</i>	17
5.2.2. Overview of the evaluation of <i>Phytophthora ramorum</i>	17
5.2.3. Overview of the evaluation of <i>Phytophthora siskiyouensis</i>	18
5.2.4. Outcome of Expert Knowledge Elicitation.....	19
6. Conclusions.....	22
Glossary	22
Abbreviations	22
Requestor.....	23
Question number.....	23
Copyright for non-EFSA content.....	23
Panel members	23
Map disclaimer	23
References.....	23
Appendix A.....	25
Appendix B	57
Appendix C	66
Appendix D.....	67

1 | INTRODUCTION

1.1 | Background and Terms of Reference as provided by European Commission

1.1.1 | Background

The Plant Health Regulation (EU) 2016/2031,¹ on the protective measures against pests of plants, has been applied from December 2019. Provisions within the above Regulation are in place for the listing of 'high risk plants, plant products and other objects' (Article 42) on the basis of a preliminary assessment, and to be followed by a commodity risk assessment. A list of 'high risk plants, plant products and other objects' has been published in Regulation (EU) 2018/2019.² Scientific opinions are therefore needed to support the European Commission and the Member States (MSs) in the work connected to Article 42 of Regulation (EU) 2016/2031, as stipulated in the terms of reference.

1.1.2 | Terms of Reference

In view of the above and in accordance with Article 29 of Regulation (EC) No 178/2002,³ the Commission asks EFSA to provide scientific opinions in the field of plant health.

In particular, EFSA is expected to prepare and deliver risk assessments for commodities listed in the relevant Implementing Act as 'high risk plants, plant products and other objects'. Article 42, paragraphs 4 and 5, establishes that a risk assessment is needed as a follow-up to evaluate whether the commodities will remain prohibited, removed from the list and additional measures will be applied or removed from the list without any additional measures. This task is expected to be ongoing, with a regular flow of dossiers being sent by the applicant required for the risk assessment.

Therefore, to facilitate the correct handling of the dossiers and the acquisition of the required data for the commodity risk assessment, a format for the submission of the required data for each dossier is needed.

Furthermore, a standard methodology for the performance of 'commodity risk assessment' based on the work already done by Member States and other international organisations needs to be set.

In view of the above and in accordance with Article 29 of Regulation (EC) No 178/2002, the Commission asked EFSA to provide scientific opinion in the field of plant health for *Alnus cordata* (Loisel.) Duby, *A. glutinosa* (L.) Gaertn and *A. incana* (L.) Moench plants from the United Kingdom (UK) taking into account the available scientific information, including the technical dossier provided by the UK.

1.2 | Interpretation of the Terms of Reference

The EFSA Panel on Plant Health (hereafter referred to as 'the Panel') was requested to conduct a commodity risk assessment of *A. cordata*, *A. glutinosa* and *A. incana* plants from the UK following the Guidance on commodity risk assessment for the evaluation of high-risk plant dossiers (EFSA PLH Panel, 2019) and the protocol for commodity risk assessment as presented in the EFSA standard protocols for scientific assessment (EFSA PLH Panel, 2024a, 2024b; Gardi et al., 2024), taking into account the available scientific information, including the technical information provided by the UK.

In the Dossier, requests were made for the following commodity types: graftwood, bare-root plants, cell-grown plants, rooted plants in pots and large specimen trees in pots.

The commodities graftwood, bare-root plants, cell-grown plants and rooted plants in pots were addressed in another Opinion (EFSA PLH Panel, 2025), while in this Opinion, only large specimen trees (from 7- to 25-year-old trees) of *A. cordata* and *A. glutinosa* are considered for evaluation.

After the consultation with the Commission, the Panel was informed that the commodity to be evaluated could be up to 25 years old and grown in bare soil for up to 9 years, updating the growing condition described in the Dossier.

In accordance with the Agreement on the withdrawal of the United Kingdom of Great Britain and Northern Ireland from the European Union and the European Atomic Energy Community, and in particular Article 5(4) of the Windsor Framework in conjunction with Annex 2 to that Framework, for the purposes of this Opinion, references to the UK do not include Northern Ireland.

¹Regulation (EU) 2016/2031 of the European Parliament of the Council of 26 October 2016 on protective measures against pests of plants, amending Regulations (EU) 228/2013, (EU) 652/2014 and (EU) 1143/2014 of the European Parliament and of the Council and repealing Council Directives 69/464/EEC, 74/647/EEC, 93/85/EEC, 98/57/EC, 2000/29/EC, 2006/91/EC and 2007/33/EC. OJ L 317, 23.11.2016, pp. 4–104.

²Commission Implementing Regulation (EU) 2018/2019 of 18 December 2018 establishing a provisional list of high risk plants, plant products or other objects, within the meaning of Article 42 of Regulation (EU) 2016/2031 and a list of plants for which phytosanitary certificates are not required for introduction into the Union, within the meaning of Article 73 of that Regulation C/2018/8877. OJ L 323, 19.12.2018, pp. 10–15.

³Regulation (EC) No 178/2002 of the European Parliament and of the Council of 28 January 2002 laying down the general principles and requirements of food law, establishing the European Food Safety Authority and laying down procedures in matters of food safety. OJ L 31, 1.2.2002, pp. 1–24.

The EU-quarantine pests that are regulated as a group in the Commission Implementing Regulation (EU) 2019/2072⁴ were considered and evaluated separately at species level.

Annex II of Implementing Regulation (EU) 2019/2072 lists certain pests as non-European populations or isolates or species. These pests are regulated quarantine pests. Consequently, the respective European populations, or isolates, or species are non-regulated pests.

Annex VII of the same Regulation, in certain cases (e.g. point 32) makes reference to the following countries that are excluded from the obligation to comply with specific import requirements for those non-European populations, or isolates, or species: Albania, Andorra, Armenia, Azerbaijan, Belarus, Bosnia and Herzegovina, Canary Islands, Faeroe Islands, Georgia, Iceland, Liechtenstein, Moldova, Monaco, Montenegro, North Macedonia, Norway, Russia (only the following parts: Central Federal District (Tsentrally federalny okrug), Northwestern Federal District (SeveroZapadny federalny okrug), Southern Federal District (Yuzhny federalny okrug), North Caucasian Federal District (Severo-Kavkazsky federalny okrug) and Volga Federal District (Privolzhsky federalny okrug), San Marino, Serbia, Switzerland, Türkiye, Ukraine and the UK (except Northern Ireland⁵). Those countries are historically linked to the reference to 'non-European countries' existing in the previous legal framework, Directive 2000/29/EC.

Consequently, for those countries,

- (i) any pests identified, which are listed as non-European species in Annex II of Implementing Regulation (EU) 2019/2072 should be investigated as any other non-regulated pest;
- (ii) any pest found in a European country that belongs to the same denomination as the pests listed as non-European populations or isolates in Annex II of Implementing Regulation (EU) 2019/2072, should be considered as European populations or isolates and should not be considered in the assessment of those countries.

Pests listed as 'Regulated Non-Quarantine Pest' (RNQP) in Annex IV of the Commission Implementing Regulation (EU) 2019/2072, and deregulated pests [i.e. pests which were listed as quarantine pests in the Council Directive 2000/29/EC and were deregulated by Commission Implementing Regulation (EU) 2019/2072] were not considered for further evaluation.

In case a pest is at the same time regulated as an RNQP and as a protected zone quarantine pest, in this Opinion, it should be evaluated as quarantine pest.

In its evaluation, the Panel:

- checked whether the provided information in the technical dossier (hereafter referred to as 'the Dossier') provided by the applicant (UK, Department for Environment Food and Rural Affairs – from this point onwards referred to as 'DEFRA') was sufficient to conduct a commodity risk assessment. When necessary, additional information was requested to the applicant;
- selected the relevant Union quarantine pests and protected zone quarantine pests [as specified in Commission Implementing Regulation (EU) 2019/2072⁶, from this point onwards referred to as 'EU quarantine pests'] and other relevant pests present in the UK and associated with the commodity;
- assessed the effectiveness of the measures described in the Dossier for those Union quarantine pests for which no specific measures are in place for the importation of the commodity from the UK and other relevant pests present in the UK and associated with the commodity;
- did not assess the effectiveness of measures for Union quarantine pests for which specific measures are in place for the import of the commodity from the UK in Commission Implementing Regulation (EU) 2019/2072 and/or in the relevant legislative texts for emergency measures and if the specific country is in the scope of those emergency measures. The assessment was restricted to whether or not the applicant country implements those measures.

Risk management decisions are not within EFSA's remit. Therefore, the Panel provided a rating based on expert judgement on the likelihood of pest freedom for each relevant pest given the risk mitigation measures proposed by DEFRA of the UK.

⁴Commission Implementing Regulation (EU) 2019/2072 of 28 November 2019 establishing uniform conditions for the implementation of Regulation (EU) 2016/2031 of the European Parliament and the Council, as regards protective measures against pests of plants, and repealing Commission Regulation (EC) No 690/2008 and amending Commission Implementing Regulation (EU) 2018/2019. OJ L 319, 10.12.2019, pp. 1–279.

⁵In accordance with the Agreement on the withdrawal of the United Kingdom of Great Britain and Northern Ireland from the European Union and the European Atomic Energy Community, and in particular Article 5(4) of the Protocol on Ireland/Northern Ireland in conjunction with Annex 2 to that Protocol, for the purposes of this Opinion, references to Member States include the United Kingdom in respect of Northern Ireland.

⁶Commission Implementing Regulation (EU) 2019/2072 of 28 November 2019 establishing uniform conditions for the implementation of Regulation (EU) 2016/2031 of the European Parliament and the Council, as regards protective measures against pests of plants, and repealing Commission Regulation (EC) No 690/2008 and amending Commission Implementing Regulation (EU) 2018/2019, OJ L 319, 10.12.2019, pp. 1–279.

2 | DATA AND METHODOLOGIES

2.1 | Data provided by DEFRA of the UK

The Panel considered all the data and information in the Dossiers provided by DEFRA of the UK in July 2023. The Dossier is managed by EFSA.

The structure and overview of the Dossier is shown in **Table 1**. The number of the relevant section is indicated in the Opinion when referring to a specific part of the Dossier.

TABLE 1 Structure and overview of the Dossier.

Dossier section	Overview of contents	Filename
1	Technical dossiers	Alnus cordata commodity information final.pdf Alnus glutinosa commodity information final.pdf
2	Pest list	Alnus_Pest_List_Final_DEFRA.xlsx
3	<i>Alnus cordata</i> and <i>A. glutinosa</i> plant distribution	A_cordata_distribution.pdf A_glutinosa_distribution.pdf
4	List of plants produced in the nurseries	Alnus_cordata_producers_sample_product_list.xlsx Alnus_glutinosa_producers_sample_product_list.xlsx
5	Additional information: Information from DEFRA on <i>Phytophthora siskiyouensis</i> outbreaks in the UK	RE Info on Phytophthora siskiyouensis in UK.pdf

The data and supporting information provided by DEFRA of the UK formed the basis of the commodity risk assessment. **Table 2** shows the main data sources used by DEFRA of the UK to compile the Dossier (Dossier Sections 1 and 2).

TABLE 2 Databases used in the literature searches by DEFRA of the UK.

Database	Platform/link
Aphids on World Plants	https://www.aphidsonworldplants.info/
Beetles of Britain and Ireland	https://www.coleoptera.org.uk/
Biological Records Centre	https://www.brc.ac.uk/
British Bugs	https://www.britishbugs.org.uk/gallery.html
Butterflies and Moths of North America	https://www.butterfliesandmoths.org/
CABI Crop Protection Compendium	https://www.cabi.org/cpc/
CABI Plantwise Knowledge Bank	https://www.plantwise.org/knowledgebank/
CABI Publishing	https://www.cabi.org/what-we-do/publishing/
Checklist of Aphids of Britain	https://influentialpoints.com/aphid/Checklist_of_aphids_in_Britain.htm
Encyclopedia of Life	https://eol.org/
EPPO Global Database	https://gd.eppo.int/
Fauna Europaea	https://www.gbif.org/dataset/90d9e8a6-0ce1-472d-b682-3451095dbc5a
Forest research	https://www.forestryresearch.gov.uk/
Fungi of Great Britain and Ireland	https://fungi.myspecies.info/
Global Biodiversity Information Facility	https://www.gbif.org/
Global Taxonomic Database of Gracillariidae (Lepidoptera)	https://www.gbif.org/dataset/98fb9418-8215-4575-abfb-07a30b81acf
National Collection of Plant Pathogenic Bacteria (NCPPB)	https://ncppb.fera.co.uk/ncppbresult.cfm
Nature Spot	https://www.naturespot.org.uk/
Natural History Museum (NHM)	https://data.nhm.ac.uk/dataset/hosts
NBN Atlas	https://species.nbnatlas.org/
NEMAPLEX	https://nemplex.ucdavis.edu/
Plant Parasites of Europe – leafminers, galls and fungi	https://bladmineerders.nl/
Pyrenomycetes from southwestern France	https://pyrenomycetes.free.fr/
Scalenet	https://scalenet.info/
Spider Mites Web	https://www1.montpellier.inra.fr/CBGP/spmweb/
The Sawflies (Symphyta) of Britain and Ireland	https://www.sawflies.org.uk/
Thrips-iD	https://www.thrips-id.com/en/
UK Beetles	https://www.ukbeetles.co.uk/
UK Moths	https://ukmoths.org.uk/
UK Plant Health Information Portal	https://planthealthportal.defra.gov.uk/

2.2 | Literature searches performed by EFSA

Literature searches in different databases were undertaken by EFSA to complete a list of pests potentially associated with the genus *Alnus*. The following searches were performed successively: (i) a general search to identify pests reported on the genus *Alnus* in the databases, and subsequently (ii) a tailored search to identify whether the pests identified by the first search are present or not in the UK. The searches were run on 30 January 2024. No language, date or document type restrictions were applied in the search strategy.

The Panel used the databases indicated in [Table 3](#) to compile the list of pests associated with the genus *Alnus*. In Web of Science, the literature search was performed using a specific, ad hoc established search string ([Appendix B](#)). The string was run in 'All Databases' with no range limits for time or language filters. The methodology is further explained in [Section 2.3.2](#).

TABLE 3 Databases used by EFSA for the compilation of the pest list associated with *Alnus* spp.

Database	Platform/link
Aphids on World Plants	https://www.aphidsonworldsplants.info/C_HOSTS_AAIntro.htm
CABI Crop Protection Compendium	https://www.cabi.org/cpc/
Database of Insects and their Food Plants	https://www.brc.ac.uk/dbif/hosts.aspx
Database of the World's Lepidopteran Hostplants	https://www.nhm.ac.uk/our-science/data/hostplants/search/index.dsml
EPPO Global Database	https://gd.eppo.int/
EUROPHYT	https://webgate.ec.europa.eu/europhyt/
Global Biodiversity Information Facility	https://www.gbif.org/
Google Scholar	https://scholar.google.com/
Leafminers	https://www.leafmines.co.uk/html/plants.htm
Nemaplex	https://nemaplex.ucdavis.edu/Nemabase2010/PlantNematodeHostStatusDDQuery.aspx
Plant Parasites of Europe	https://bladmineerders.nl/
Plant Pest Information Network	https://www.mpi.govt.nz/news-and-resources/resources/registers-and-lists/plant-pest-information-network/
Plant Viruses Online	https://www1.biologie.uni-hamburg.de/b-online/e35/35tmv.htm#Range
Scalenet	https://scalenet.info/associates/
Spider Mites Web	https://www1.montpellier.inra.fr/CBGP/spmweb/advanced.php
USDA ARS Fungal Database	https://fungi.ars.usda.gov/
Web of Science: All Databases (Web of Science Core Collection, CABI: CAB Abstracts, BIOSIS Citation Index, Chinese Science Citation Database, Current Contents Connect, Data Citation Index, FSTA, KCI-Korean Journal Database, Russian Science Citation Index, MEDLINE, SciELO Citation Index, Zoological Record)	Web of Science https://www.webofknowledge.com
World Agroforestry	https://www.worldagroforestry.org/treedb2/speciesprofile.php?Spid=1749
The American Phytopathological Society	https://www.apsnet.org/Pages/default.aspx

Additional documents were retrieved when developing the Opinion. The available scientific information, including previous EFSA opinions on the relevant pests and diseases and the relevant literature and legislation (e.g. Regulation (EU) 2016/2031; Commission Implementing Regulations (EU) 2018/2019; (EU) 2018/2018 and (EU) 2019/2072) were taken into account.

2.3 | Methodology

When developing the Opinion, the Panel followed the EFSA Guidance on commodity risk assessment for the evaluation of high-risk plant dossiers (EFSA PLH Panel, [2019](#)). In the first step, pests potentially associated with the commodity in the country of origin (EU-regulated pests and other pests) that may require risk mitigation measures are identified. The EU non-regulated pests not known to occur in the EU were selected based on evidence of their potential impact in the EU. After the first step, all the relevant pests that may need risk mitigation measures were identified. In the second step, if applicable, the implemented risk mitigation measures for each relevant pest are evaluated. In the final step, a conclusion on the pest freedom status of the commodity for each of the relevant pests, if any, is determined and uncertainties identified using expert judgements.

Pest freedom was assessed by estimating the number of infested/infected units out of 10,000 exported units of large specimen trees in pots from 7 to 25 years old.

2.3.1 | Commodity data

Based on the information provided by DEFRA of the UK, the characteristics of the commodity are summarised in Section 3 of this Opinion.

2.3.2 | Identification of pests potentially associated with the commodity

To evaluate the pest risk associated with the importation of the commodity from the UK, a pest list was compiled. The pest list is a compilation of all identified plant pests reported as associated with all species of *Alnus* based on information provided in the Dossier Sections 1 and 2 and on further literature searches performed by the Panel. The search strategy and search syntax were adapted to each of the databases listed in [Tables 3](#), according to the options and functionalities of the different databases and CABI keyword thesaurus.

The scientific name of the host plant (i.e. *Alnus*) was used when searching in the European and Mediterranean Plant Protection Organisation (EPPO) Global database (EPPO GD, [online](#)) and CABI Crop Protection Compendium (CABI, [online](#)). The same strategy was applied to the other databases ([Table 3](#)) excluding EUROPHYT and Web of Science. The notifications of interceptions associated with *Alnus* spp. from the whole world to the EU were investigated on EUROPHYT from 1995 to May 2020 and TRACES-NT from May 2020 to January 2025. To check whether *Alnus* spp. can act as a pathway, all notifications (all origins) for *Alnus* spp. were evaluated. For each selected pest, it was checked if there were any notification records for UK (all commodities).

The search query used for Web of Science Databases was designed combining English common names for pests and diseases, terms describing symptoms of plant diseases and the scientific and English common names of the commodity and excluding pests which were identified during searches in other databases. The established search string is detailed in [Appendix B](#) and was run on 30 January 2024.

The titles and abstracts of the scientific papers retrieved were screened and the pests associated with *Alnus* genus were included in the pest list. The pest list was eventually further updated with other relevant information (e.g. EPPO code per pest, taxonomic information, categorisation and distribution) useful for the selection of the pests relevant for the purposes of this Opinion.

The compiled pest list includes all pests reported as hosted by *Alnus* genus ([Appendix D](#)).

The relevance of EU-quarantine pests was first assessed ([Section 4.1](#)), followed by an assessment of the relevance of any other plant pests ([Section 4.2](#)).

2.3.3 | Listing and evaluation of risk mitigation measures

All proposed risk mitigation measures were listed and evaluated. When evaluating the likelihood of pest freedom at origin, the following types of potential infestation/infection sources for *A. cordata* and *A. glutinosa* in nurseries were considered (see also [Figure 1](#)):

- Pest entry from surrounding areas,
- Pest entry with new plants/seeds,
- Pest spread within the nursery.




FIGURE 1 Conceptual framework to assess likelihood that plants are exported free from relevant pests (Source: EFSA PLH Panel, [2019](#)).

Information on the biology, estimates of likelihood of entry of the pest into the nursery and spread within the nursery, and the effect of the measures on a specific pest is summarised in pest data sheets compiled for each pest selected for further evaluation (Appendix A).

2.3.4 | Expert Knowledge Elicitation

To estimate the pest freedom of the commodities, an Expert Knowledge Elicitation (EKE) was performed following EFSA guidance (Annex B.8 of EFSA Scientific Committee, 2018).

The specific question for EKE was defined as follows: 'taking into account (i) the risk mitigation measures listed in the Dossier, and (ii) other relevant information (reported in the specific pest datasheets), how many of 10,000 large specimen trees in pots from 7 to 25 years old will be infested/infected with the relevant pest/pathogen when arriving in the EU?'. The uncertainties associated with the EKE were taken into account and quantified in the probability distribution applying the semi-formal method described in Section 3.5.2 of the EFSA-PLH Guidance on quantitative pest risk assessment (EFSA PLH Panel, 2018). Finally, the results were reported in terms of the likelihood of pest freedom. The lower 5% percentile of the uncertainty distribution reflects the opinion that pest freedom is with 95% certainty above this limit.

3 | COMMODITY DATA

3.1 | Description of the commodity

The commodity consists of the following type of deciduous plant of *A. cordata* (Loisel.) Duby (common name: Italian alder; family: Betulaceae) and *A. glutinosa* (L.) Gaertn. (common name: alder, common alder, black alder; family: Betulaceae) (Figure 2) to be imported from UK to EU as large specimen trees in pots from 7 to 25 years old (Table 4). These trees may be field grown in bare soil for up to 9 years (see Section 3.3.3 for more details on growing conditions).

TABLE 4 Type of *Alnus cordata* and *A. glutinosa* specimen trees to be exported to the EU (Dossier Section 1).

Type of plant	Age	Diameter	Height/length
Large specimen trees in pots	7–25 years	80–600 mm	Up to 12 m




FIGURE 2 Specimen trees for *Alnus cordata* (A) and *A. glutinosa* (B) growing in 80 litre containers in a row on gravel (Source: Dossier Section 1).

According to ISPM 36 (FAO, 2019), the commodity can be classified as 'rooted plants in pots'. According to the Dossier Section 1, the expected trade volume for *A. cordata* and *A. glutinosa* is estimated to be 350 specimen trees per year.

The growing media is virgin peat or peat-free compost (a mixture of coir, tree bark, wood fibre, etc.) (Dossier Sections 1) complying with the requirements for growing media as specified in the Annex VII of the Commission Implementing Regulation 2019/2072.

Rooted plants in pots can be exported at any time in the year to fulfil consumer demand, but more usually from September/October to April/May. Rooted plants in pots may be exported with leaves, depending on the timing of the export and the life cycle of the species, in any period of the year (Dossier Section 1).

3.2 | Description of the production areas

According to the dossier, specific nurseries contributed to the technical information provided by the applicant (Figure 3), but this does not exclude the possibility that other nurseries in the UK may be willing to export *A. cordata* and *A. glutinosa*.

large specimen trees in pots from 7 to 25 years old to the EU in the future. Such nurseries would need to meet the import requirements set out in any subsequent EU legislation, as would the nurseries that have contributed technical information to the dossier.

All nurseries are registered as professional operators with the UK NPPO, either by the Animal and Plant Health Agency (APHA) in England and Wales, or by the Scottish Government, and are authorised to issue UK plant passports and phytosanitary certificates for export (Dossier Section 1).

Approximately 20% of the nurseries likely to export to the EU also sell plants within the UK to final users as ornamental plants, e.g. to the Local Authorities/Landscape Architects (Dossier Section 1). Trade of all plant types will mainly be to Northern Ireland and Republic of Ireland. Most of the nurseries also produce plants for the local market, and there is no distancing between production areas for the export and the local market.




FIGURE 3 Production areas of *Alnus cordata* and *A. glutinosa* in UK for export to the EU (Dossier Section 1). The coordinates of nursery production facilities, provided by the applicant, were used to generate the custom map.

The minimum and maximum sizes of nurseries growing *A. cordata* and *A. glutinosa* for export are as follows: for container grown stock, a minimum of 8 ha and a maximum of 150 ha; for field-grown stock intended for bare root plants, the maximum size is 325 ha.

The exporting nurseries cultivate a variety of other plant species (Dossier Section 4). The production area for *A. glutinosa* plants is approximately 0.1%–4% of the total nurseries area, while the production area for *A. cordata* plants is around 1%–4%. The commodities grown at the nurseries will vary depending on the nursery (Dossier Section 1).

The surrounding areas of exporting nurseries are predominately rural, mainly characterised by arable farmland with some pasture for livestock and small areas of woodland. Arable crops are rotated in line with good farming practices and could include oilseed rape (*Brassica napus*), wheat (*Triticum* spp.), barley (*Hordeum vulgare*), turnips (*Brassica rapa* subsp. *rapa*), potatoes (*Solanum tuberosum*) and maize (*Zea mays*) (Dossier Section 1).

The pasture is predominantly composed of ryegrass (*Lolium* spp.) (Dossier Section 1). Woodlands tend to be a standard UK mixed woodland, with a range of UK native trees such as oak (*Quercus robur*), pine (*Pinus* spp.), poplar (*Populus* spp.), ash (*Fraxinus* spp.), sycamore (*Acer pseudoplatanus*), holly (*Ilex* spp.), Norway maple (*Acer platanoides*) and field maple (*Acer campestre*) (Dossier Section 1).

Shelter boundary hedges are mostly alders (*A. glutinosa* or *A. cordata*) with hawthorn (*Crataegus monogyna*) and field maple (*A. campestris*) interspersed. There is also Western red cedar (*Thuja plicata*) along neighbours' boundaries.

It is not possible to identify what plant species are growing within the gardens of private dwellings. The nearest woodland to one of the nursery borders the boundary fence, and its composition is as per the description above.

3.3 | Production and handling processes

3.3.1 | Growing conditions

As the plants are intended for outdoor cultivation, only early growth stages are normally maintained under protection, such as young plants/seedlings that are vulnerable to climatic conditions including frost. The commodity to be exported should therefore be regarded as outdoor grown. Growth under protection is primarily to protect against external climatic conditions rather than protection from pests. The early stages of plants grown under protection are maintained in plastic polytunnels, or in glasshouses which typically consist of a metal or wood frame construction and glass panels (Dossier Section 1).

3.3.2 | Source of planting material

The starting material is a mix of seeds and seedlings depending on the nursery. *Alnus cordata* seeds purchased in the UK are not covered by The Forest Reproductive Material (Great Britain) Regulations 2002 (legislation.gov.uk). *Alnus glutinosa* seeds purchased in the UK are certified under The Forest Reproductive Material (Great Britain) Regulations 2002 (legislation.gov.uk). *Alnus cordata*, *A. glutinosa* seedlings sourced in the UK are traded with UK Plant Passports; a small percentage of plants may be obtained from the EU (The Netherlands); seedlings originating from the EU countries are certified with phytosanitary certificates (Dossier Section 1). None of the nurseries expected to export *A. glutinosa* to the EU produce plants from grafting. Only one of the nurseries expected to export *A. cordata* to the EU produces plants from grafting. This nursery has mother plants of *A. cordata* on site, but as these are the only species produced by grafting, there are no mother plants of other *Alnus* species present. All other growers use only seed and seedlings.


3.3.3 | Production cycle

The growing conditions are as follows [as defined in Annex 1 of ISPM 36 (FAO, 2019)]:

- field grown (up to 9 years);
- field grown in containers (from 7 to 25 years).

The commodity production stages, and the phenology of the crop associated are reported in [Table 5](#).

TABLE 5 Commodity production stages (planting, grafting, budding) and the phenology of the crop (including flowering, leaf drop), and harvesting periods (lifting).



*Rooted plants in pots can be planted at any time of year, with less common periods (light grey) and most common period (dark grey).

Planting. Rooted plants in pots can be planted at any time of year, though late-autumn winter is most common.

Growing. Specimen trees are grown in EU-compliant growing media in pots for their whole life, or initially field grown in bare soil (for no more than 9 years) before being lifted, root-washed to remove any soil, and subsequently potted in EU-compliant growing media. The trees are grown on racks with no substrate below them (on gravel or on a geotex root barrier (geotex 1000)) and are sold either as root balls or in peat-free container bags. Plants will be exported as a containerised tree in an air-pot system ([Figure 4A](#)), or in polypropylene woven pots ([Figure 4B](#)).

Any plants in pots with organic growing medium being exported from UK to the EU need to meet the requirements for growing media in EU Regulation 2019/2072, Annex VII.

In the production or procurement of plants, the use of growing media is assessed for the potential to harbour and transmit plant pests. Growers use virgin peat or peat-free compost, which is a mixture of coir, tree bark, wood fibre, etc. This compost is heat-treated by commercial suppliers during production to eliminate pests and diseases. It is supplied in sealed bulk bags or shrink-wrapped bales and stored off the ground on pallets, these are completely hygienic and free from contamination. Where delivered in bulk, compost is kept in a dedicated bunker, either indoors, or covered by tarpaulin outdoors, and with no risk of contamination with soil or other material (Dossier Section 1).

Soil testing might also be carried out to ensure pest freedom ahead of export.



FIGURE 4 (A) Example of open pot (air-pot system) with growing media; (B) polypropylene woven pots.

Grafting. Most of the nurseries expected to export to the EU do not use grafting in the production of *A. cordata* and *A. glutinosa*. Where it does occur, grafting is done indoors, and two different methods are used:

- side-spliced grafting is usually undertaken in late winter or early spring before bud break;
- whip and tongue grafting is normally undertaken in March or early April.

Pruning. To ensure a good root architecture, trees are regularly pruned (at least once per year) and transplanted (every 3–5 years).

Irrigation. The irrigation is done on a need basis and could be overhead, subirrigation or drip irrigation. Rainfall catchment reservoir water is used for irrigation. All water is passed through a sand filtration system to remove contaminants and is contained in storage tanks prior to use. All mains water supply used meets the UK standard Water Supply (Water quality) regulation 2016 and the WHO/EU potable water standards, (Drinking water Directive (98/83/EC and the revised Drinking Water Directive 2020/2184) which includes a total freedom from both human and plant pathogens (Article 2-(7)). All mains water supply conducting pipework fully complies with the UK Water Supply (Water Fittings) regulations of 1999 and the amendments of 2019. Irrigation water used is not stored in any open tanks where airborne contamination could take place and is entirely isolated from any outside exposure.

Regardless of the source of the water used to irrigate, the nurseries contributing information to this dossier declared that they have never experienced the introduction of a pest/disease because of contamination of the water supply.

Growers are required to assess water sources, irrigation and drainage systems used in the plant production for the potential to harbour and transmit plant pests. Water is routinely sampled and sent for analysis. No quarantine pests have been found so far (Dossier Section 1).

Hygiene measures. All nurseries have plant hygiene, housekeeping rules and practices in place, which are communicated to all relevant employees. The rules will be dependent on the plants handled and the type of business but will refer to growing media, water usage, weed management, tools and visitors.

General hygiene measures are undertaken as part of routine nursery production, including disinfection of tools and equipment between batches/ lots. Tools are disinfected after operation on a stock and before being used on a different plant species. The tools are dipped and wiped with a clean cloth between trees to reduce the risk of pathogens and pests transfer between subjects. Potassium peroxymonosulfate and sodium chloride were reported as the most commonly used disinfectant. Growers keep records allowing traceability for all plant material handled (Dossier Section 1).

Growers must assess weeds and volunteer plants for the potential to host and transmit plant pests and have an appropriate programme of weed management in place at the nursery (Dossier Section 1). Growing areas are kept clear of non-cultivated herbaceous plants. In access areas, non-cultivated herbaceous plants are kept to a minimum and only exist at nursery boundaries. Non-cultivated herbaceous plants grow on less than 1% of the nursery area (Dossier Section 1). The predominant species is rye grass (*Lolium*). Other species may include dandelions (*Taraxacum officinale*), hairy bittercress (*Cardamine hirsute*), Common daisy (*Bellis perennis*), Creeping cinquefoil (*Potentilla reptans*) and bluebells (*Hyacinthoides non-scripta*), present in a low amount.

Plant material is regularly monitored for plant health issues. This monitoring is carried out by trained nursery staff via regular crop walking and records kept of this monitoring. Qualified agronomists also undertake regular crop walks to verify

the producer's assessments. Curative or preventative actions are implemented together with an assessment of phytosanitary risk. Unless a pest can be immediately and definitively identified as non-quarantine, growers are required to treat it as a suspect quarantine pest and notify the competent authority (Dossier Section 1).

Pest and disease pressures vary from season to season. Chemical treatments are reported to be applied when required and depend on the situation at that time (disease pressure, growth stage, etc., and environmental factors) (Dossier Section 1).

To reduce the number of overwintering sites for pest and disease, waste materials (i.e. leaves, prunings and weeds) are all removed from the nurseries.

There are no specific measures/treatments against soil pests. However, the containerised young plants are grown in trays on top of protective plastic membranes to prevent contact with soil. Membranes are regularly refreshed when needed. Alternatively, young plants may be grown on raised galvanised steel benches stood on gravel as a barrier between the soil and bench feet and/or concreted surfaces (Dossier Section 1).

3.3.4 | Post-harvest processes and export procedure

The UK NPPO carries out inspections and testing (where required by the country of destination's plant health legislation) to ensure all requirements are fulfilled and a valid phytosanitary certificate with the correct additional declarations is issued.

Large plants can be transported on ISPM 15 certified pallets, or individually in pots for larger containers. Plants are transported by lorry (size dependent on load quantity) (Figure 5) (Dossier Section 1).



FIGURE 5 Large specimen trees transported by lorry.

4 | IDENTIFICATION OF PESTS POTENTIALLY ASSOCIATED WITH THE COMMODITY

The search for potential pests associated with *Alnus* spp. (and if available specific information with pests associated with *Alnus* spp. including *A. cordata* and *A. glutinosa*) retrieved 2743 pest species (for search string see Appendix B, for pest list see Appendix D).

4.1 | Selection of relevant EU-quarantine pests associated with the commodity

The EU listing of Union quarantine pests and protected zone quarantine pests (Commission Implementing Regulation (EU) 2019/2072) is based on assessments concluding that the pests can enter, establish, spread and have potential impact in the EU.

The 20 EU-quarantine species that are reported to use *Alnus* spp. as a host were evaluated (Table 6) for their relevance of being included in this Opinion.

The relevance of an EU-quarantine pest for this Opinion was based on evidence that:

- the pest is present in the UK;
- the commodity is a host of the pest;
- one or more life stages of the pest can be associated with the specified commodity.

Pests that fulfilled all criteria are selected for further evaluation.

Of the 20 EU-quarantine pest species evaluated, two pests (*Entoleuca mammata* and *Phytophthora ramorum*) were selected for further assessment.

TABLE 6 Overview of the evaluation of the 20 (protected zone) EU-quarantine pest species known to use *Alnus* species as host plants for their relevance for this Opinion.

No.	Pest name according to EU legislation*	EPPO code	Group	Pest present in the UK	<i>Alnus</i> confirmed as a host	Pest can be associated with the commodity (NA = not assessed)	Pest relevant for the opinion
1	<i>Acleris senescens</i>	ACLRSE	Insect	No	Gilligan and Epstein (2014)	NA	No
2	<i>Aleurocanthus spiniferus</i>	ALECSN	Insect	No	Dubey and Ko (2012)	NA	No
3	<i>Anoplophora chinensis</i>	ANOLCN	Insect	No	Lim et al. (2014), Sjöman et al. (2014)	NA	No
4	<i>Anoplophora glabripennis</i>	ANOLGL	Insect	No	Sjöman et al. (2014)	NA	No
5	<i>Choristoneura conflictana</i>	ARCHCO	Insect	No	Ciesla and Kruse (2009), Prentice (1966)	NA	No
6	<i>Choristoneura rosaceana</i>	CHONRO	Insect	No	Ferguson (1975), Prentice (1966)	NA	No
7	<i>Cryphonectria parasitica</i> **	ENDOPA	Fungi	Yes	Uncertain	NA	No
8	<i>Euwallacea fornicatus</i> sensu lato	XYLBFO	Insect	No	Eskalen et al. (2013), USDA (Online)	NA	No
9	<i>Entoleuca mammata</i>	HYPOMA	Fungi	Yes	Callan (1998)	Yes	Yes
10	<i>Grapevine flavescence dorée phytoplasma</i>	PHYP64	Phytoplasma	No	Malembic-Maher et al. (2020), Mehle et al. (2011), Radonjic et al. (2013), Scalenet (Online)	NA	No
11	<i>Lopholeucaspis japonica</i>	LOPLJA	Insect	No	Batsankalashvili et al. (2017), Shrewsbury et al. (2013), EPPO (online)	NA	No
12	<i>Lycorma delicatula</i>	LYCMDE	Insect	No	Barringer and Ciafré (2020), Park et al. (2009), CABI (online)	NA	No
13	<i>Monochamus guttulatus</i>	MONCGU	Insect	No	Anisimov and Bezborodov (2021)	NA	No
14	<i>Oemona hirta</i>	OEMOHI	Insect	No	Plant-SyNZ (online)	NA	No
15	<i>Phymatotrichum omnivorum</i>	PHMPOM	Fungi	No	Anonymous (1960)	NA	No
16	<i>Phytophthora ramorum</i> (non-EU isolates)	PHYTRA	Chromista	Yes	O'Hanlon et al. (2016)	Yes	Yes
17	<i>Popillia japonica</i>	POPIJA	Insect	No	Fleming (1972); Regione Lombardia Servizio Fitosanitario (Online)	NA	No
18	<i>Pseudopityophthorus pruinosis</i>	PSDPPR	Insect	No	Atkinson (online)	NA	No
19	<i>Spodoptera ornithogalli</i>	PRODOR	Insect	No	Brito et al. (2019), Palmer (1987)	NA	No
20	<i>Trirachys sartus</i>	AELSSA	Insect	No	Farashiani et al. (2001)	NA	No

*Commission Implementing Regulation (EU) 2019/2072. **There is only one host record of *C. parasitica* on *Alnus* sp. (Turchetti et al., 1991). This host record could not be verified, and therefore, this pathogen was not considered as relevant for this commodity.

4.2 | Selection of other relevant pests (non-quarantine in the EU) associated with the commodity

The information provided by the UK, integrated with the search performed by EFSA, was evaluated in order to assess whether there are other relevant pests potentially associated with the *A. cordata* and *A. glutinosa* present in the exporting country. For these potential pests that are non-regulated in the EU, pest risk assessment information on the probability of entry, establishment, spread and impact is usually lacking. Therefore, these pests were also evaluated to determine their relevance for this Opinion based on evidence that:

- a. the pest is present in the UK;
- b. the pest is (i) absent or (ii) has a limited distribution in the EU;
- c. *Alnus* spp. is a host of the pest;
- d. one or more life stages of the pest can be associated with the traded commodity of *Alnus* spp.;
- e. the pest may have an impact in the EU.

For non-regulated species with a limited distribution in the EU (i.e. present in one or a few EU member states), they should also satisfy at least one of the following conditions for the pest to be selected for further evaluation:

- official phytosanitary measures have been adopted in at least one EU MS;
- any other reason justified by the working group (e.g. recent evidence of presence).

Based on the information collected, 2723 potential pests (non-EU quarantine) known to be associated with *Alnus* spp. were evaluated for their relevance to this Opinion.

Species were excluded from further evaluation when at least one of the conditions listed above (a-e) was not met. Details can be found in the pest list (Appendix D). Of the evaluated pests not regulated in the EU, one pest (*Phytophthora siskiyouensis*) was selected for further evaluation because it met all the selection criteria. More information on this species can be found in the pest datasheets (Appendix A).

4.3 | List of potential pests not further assessed

From the list of pests not selected for further evaluation, the Panel highlighted four species (Appendix C) for which there was uncertainty on one of the criteria to be selected for further evaluation in this Opinion. A specific justification of the inclusion in this list is provided for each species in Appendix C.

4.4 | Summary of pests selected for further evaluation

Three pests that were identified to be present in UK and having potential for association with *A. cordata* and *A. glutinosa* plants designated for export to the EU, were selected for further evaluation (Table 7). The efficacy of the risk mitigation measures applied to the commodity were evaluated for these selected pests.

TABLE 7 List of relevant pests selected for further evaluation.

No.	Current scientific name	EPPO code	Taxonomic information	Group	Regulatory status
1	<i>Entoleuca mammata</i>	HYPOMA	Xylariales, Xylariaceae	Fungi	EU protected zone Quarantine Pest
2	<i>Phytophthora ramorum</i> (non-EU isolates)	PHYTRA	Peronosporales, Peronosporaceae	Chromista	EU Quarantine Pest
3	<i>Phytophthora siskiyouensis</i>	PHYTSK	Peronosporales, Peronosporaceae	Chromista	Not regulated in EU

5 | RISK MITIGATION MEASURES

For the selected pests (Table 7), the Panel evaluated the likelihood that they could be present in the *A. cordata* and *A. glutinosa* nurseries by evaluating the possibility that the commodity in the export nurseries is infested/infected either by:

- introduction of the pest from the environment surrounding the nursery;
- introduction of the pest with new plants/seeds;
- spread of the pest within the nursery.

The information used in the evaluation of the effectiveness of the risk mitigation measures is summarised in pest data sheets (Appendix A).

5.1 | Risk mitigation measures applied in the UK

With the information provided by the UK (Dossier Sections 1, 2, 3, & 4), the Panel summarised the risk mitigation measures (Table 8) that are implemented in the production nursery.

TABLE 8 Overview of implemented risk mitigation measures for *Alnus cordata* and *A. glutinosa* plants designated for export to the EU from the UK.

No.	Risk mitigation measure	Implementation in the UK
1	Registration of production sites	All nurseries are registered as professional operator with the UK NPPO, by the APHA for England and Wales, or with SASA for Scotland, and are authorised to issue UK plant passports (Dossier Section 1).
2	Certification of propagation material	<i>Alnus cordata</i> and <i>A. glutinosa</i> seeds purchased in the UK are certified under The Forest Reproductive Material (Great Britain) Regulations 2002 (legislation.gov.uk); seedlings sourced in the UK are certified with UK Plant Passports; a small percentage of plants may be obtained from EU (Netherlands); seedlings from the EU countries are certified with phytosanitary certificates. (Dossier Section 1).
3	Origin and treatment of growing media	In the production or procurement of these plants, the use of growing media is assessed for the potential to harbour and transmit plant pests. Growers most commonly use virgin peat or peat-free compost, which is a mixture of coir, tree bark, wood fibre, etc. The compost is heat-treated by commercial suppliers during production to eliminate pests and diseases. It is supplied in sealed bulk bags or shrink-wrapped bales and stored off the ground on pallets, these are completely hygienic and free from contamination. Where delivered in bulk, compost is kept in a dedicated bunker, either indoors, or covered by tarpaulin outdoors, and with no risk of contamination with soil or other material (Dossier Section 1).
4	Surveillance, monitoring and sampling	Inspections are carried out at least once a year as part of the Quarantine Surveillance programme (Great Britain uses the same framework for its surveillance programme as the EU). Surveillance is based on visual inspection with samples taken from symptomatic material, and where appropriate, samples are also taken from asymptomatic material (e.g., plants, soil, watercourses) (Dossier Section 1).
5	Hygiene measures	According to the Dossier Section 1, all the nurseries have plant hygiene and housekeeping rules and practices in place, which are communicated to all relevant employees.
6	Irrigation water quality and/or treatments	Growers are required to assess water sources, irrigation and drainage systems used in the plant production for the potential to harbour and transmit plant pests. Rainwater that is collected is sand filtrated. Water is routinely sampled and sent for analysis. No quarantine pests have been found (Dossier Section 1).
7	Application of pest control products	Crop protection is achieved using a combination of measures including approved plant protection products, biological control or physical measures. Plant protection products are only used when necessary and records of all plant protection treatments are kept (Dossier Section 1).
8	Washing of the roots	Specimen trees (up to 25 years old) are grown in soil for up to 9 years, root washed and placed in pots (based on the request letter of DEFRA to the EU Commission on September 25th, 2023).
9	Inspections and management of plants before export	The UK NPPO carries out inspections and testing where required by the country of destination's plant health legislation, to ensure all requirements are fulfilled and a valid phytosanitary certificate with the correct additional declarations is issued. Separate to any official inspection, plant material is checked by growers for plant health issues before dispatch.

5.2 | Evaluation of the current measures for the selected pests including uncertainties

The relevant risk mitigation measures acting on the selected pests were identified. Factors reducing the efficacy of the measures were documented. All the relevant information including the related uncertainties deriving from the limiting factors used in the evaluation are summarised in the pest datasheets provided in Appendix A. Based on this information, an expert judgement has been given for the likelihood of pest freedom of the commodity taking into consideration the risk mitigation measures acting on the pest and their combination. An overview of the evaluation of the selected pests is given in the sections below (Sections 5.2.1–5.2.3). The outcome of EKE on pest freedom after the evaluation of the proposed risk mitigation measures is summarised in Section 5.2.4.

5.2.1 | Overview of the evaluation of *Entoleuca mammata*

Overview of evaluation of <i>Entoleuca mammata</i> for specimen trees					
Rating of the likelihood of pest freedom	Pest free with few exceptional cases (based on the median)				
Percentile of the distribution	5%	25%	Median	75%	95%
Proportion of pest-free plants	9905 out of 10,000 plants	9952 out of 10,000 plants	9973 out of 10,000 plants	9987 out of 10,000 plants	9997 out of 10,000 plants
Proportion of infested plants	3 out of 10,000 plants	13 out of 10,000 plants	27 out of 10,000 plants	48 out of 10,000 plants	95 out of 10,000 plants
Summary of the information used for the evaluation	<p>Possibility that the pest could become associated with the commodity</p> <p><i>Entoleuca mammata</i> is present in the UK, although not widely distributed. <i>Entoleuca mammata</i> has been reported on various <i>Alnus</i> spp., including <i>A. incana</i>, <i>A. crispa</i>, <i>A. rugosa</i>, <i>A. sinuata</i> and <i>A. tenuifolia</i> (for references see Appendix A.1.1). Given the fact that it has been reported in several <i>Alnus</i> spp. the Panel assumes that <i>A. cordata</i> and <i>A. glutinosa</i> can be a host of <i>E. mammata</i>. The pathogen can naturally spread with ascospores dispersed by air currents from the surrounding natural environment. Furthermore, mechanical wounds are expected to be present and may represent infection courts. Altogether, this suggests that the association of <i>E. mammata</i> with the commodity is possible.</p> <p>The reasoning of EKE values in the previous Opinion of <i>Alnus</i> spp. were considered and because of the similarity of the commodities, the production systems, the locations of the nurseries, the Panel also considered the EKE scenarios from the previous Scientific opinion on <i>Betula pendula</i> and <i>B. pubescens</i> from the UK (EFSA PLH Panel, 2024a, 2024b) for <i>A. cordata</i> and <i>A. glutinosa</i>.</p> <p>As a result of this evaluation, the final estimated values for <i>Alnus</i> spp. specimen trees were based on the values for <i>Betula</i> spp. specimen trees, and <i>Alnus</i> spp. and adjusted by considering also the differences in the age of the plants.</p> <p>Pest control measures applied during production</p> <p>Pest control measures applied during the production of <i>Alnus</i> plants include: (a) registration of production sites; (b) the use of certified plant material; (c) surveillance, monitoring and sampling; (d) hygiene measures; (e) irrigation water testing; (f) washing of the roots of the field grown plants (up to 9 years old); (g) application of pest control products; (h) inspection and management of plants before export.</p> <p>Evaluation of control measures</p> <p>In general, the measures taken by the growers are effective against this pathogen. The following critical points were identified:</p> <ul style="list-style-type: none"> – The exposure time of large specimen trees can be up to 25 years. – Early infections are difficult to be detected by visual inspections. <p>Main uncertainties</p> <ul style="list-style-type: none"> – The pest pressure in the surrounding environment (the presence and density of infected host plants). The level of susceptibility of <i>Alnus</i> spp. to the pathogen. – Whether symptoms on <i>Alnus</i> spp. are recognisable and may be promptly detected. 				

5.2.2 | Overview of the evaluation of *Phytophthora ramorum*

Overview of evaluation of <i>Phytophthora ramorum</i> specimen trees					
Rating of the likelihood of pest freedom	Pest free with some exceptional cases (based on the median)				
Percentile of the distribution	5%	25%	Median	75%	95%
Proportion of pest-free plants	9920 out of 10,000 plants	9955 out of 10,000 plants	9975 out of 10,000 plants	9988 out of 10,000 plants	9997 out of 10,000 plants
Proportion of infected plants	3 out of 10,000 plants	12 out of 10,000 plants	25 out of 10,000 plants	45 out of 10,000 plants	80 out of 10,000 plants
Summary of the information used for the evaluation	<p>Possibility that the pest could become associated with the commodity</p> <p><i>Phytophthora ramorum</i> is present in most regions of the UK, but it is more often reported in wetter, western regions. <i>Phytophthoraramorum</i> has a wide host range, including <i>A. cordata</i> as natural host, whereas <i>A. glutinosa</i> was reported to be susceptible hosts following artificial inoculation (for references see Appendix A.2).</p> <p>The possible entry of <i>P. ramorum</i> from the surrounding environment may occur through wind, water and soil containing propagules on feet of animals/humans entering the field (if any). The pathogen can also enter with new seedlings of <i>Alnus</i> spp. and new plants of other species used for plant production in the nurseries.</p> <p>The reasoning of EKE values in the previous Opinion of <i>Alnus</i> spp. were considered and because of the similarity of the commodities, the production systems, the locations of the nurseries, the Panel also considered the EKE scenarios from the previous Scientific opinion on <i>Betula pendula</i> and <i>B. pubescens</i> from the UK (EFSA PLH Panel, 2024a, 2024b) for <i>A. cordata</i> and <i>A. glutinosa</i>.</p>				

(Continues)

(Continued)

As a result of this evaluation, the final estimated values for *Alnus* spp. specimen trees were based on the values for *Betula* spp. specimen trees, and *Alnus* spp. and adjusted by considering also the differences in the age of the plants.

Pest control measures applied during production

Pest control measures applied during the production of *Alnus* plants include: (a) registration of production sites; (b) the use of certified plant material; (c) surveillance, monitoring and sampling; (d) hygiene measures; (e) irrigation water testing; (f) washing of the roots of the field grown plants (up to 9 years old); (g) application of pest control products; (h) inspection and management of plants before export.

Evaluation of control measures

In general, the measures taken by the growers are effective against this pathogen. The following critical points were identified:

- The exposure time of large specimen tree can be up to 25 years trees.
- The commodity is grown for up to 9 years in the field, before being transplanted into pots with growing media.
- The washing of the roots removes (parts of) the soil and the pathogen present in the soil, but it does not remove the pathogen that may occasionally be present in the roots.
- Leaves could be present on the commodity at the time of export increasing the probability that the oomycete could be present.
- Early infections are difficult to be detected by visual inspections.

Main uncertainties

- The efficiency of the hygiene measures especially concerning the cleaning of the machinery and the possible movement of soil within the nursery.
- Whether symptoms on *Alnus* spp. are recognisable and may be promptly detected.
- Effect or efficiency of fungicide treatments against the pathogen.
- The susceptibility of *Alnus* spp. to the pathogen.
- The pest pressure in the surrounding environment (the presence and density of infected host plants).

5.2.3 | Overview of the evaluation of *Phytophthora siskiyouensis*

Overview of evaluation of *Phytophthora siskiyouensis* specimen trees

Rating of the likelihood of pest freedom	Almost always pest free (based on the median)				
Percentile of the distribution	5%	25%	Median	75%	95%
Proportion of pest-free plants	9991 out of 10,000 plants	9994.5 out of 10,000 plants	9996.4 out of 10,000 plants	9997.9 out of 10,000 plants	9999.2 out of 10,000 plants
Proportion of infected plants	0.8 out of 10,000 plants	2.1 out of 10,000 plants	3.6 out of 10,000 plants	5.5 out of 10,000 plants	9 out of 10,000 plants

Summary of the information used for the evaluation

Possibility that the pest could become associated with the commodity

There is one record of the presence of *P. siskiyouensis* in the UK, reported on *Alnus incana* plants in 2013 within a conifer and broadleaves plantation in the southwest of England (Perez-Sierra et al., 2015). The host range of *P. siskiyouensis* include also *A. cordata*, *A. rubra* and *A. glutinosa*. *P. siskiyouensis* does not have a broad host range. *P. siskiyouensis* has been also reported in myrtlewood (*Umbellularia californica*) and tanoak (*Lithocarpus densiflorus*) in south-western Oregon (USA) (for references see Appendix A.3). There is a possibility that the pathogen is present in areas where the export nurseries are located. Possible pathways of the pathogen are water and soil. The possible entry of *P. siskiyouensis* from the surrounding environment may occur through wind, water and infested soil propagules on feet of animals/humans entering the field (if any). The pathogen can also enter with new seedlings of *Alnus* spp. and new plants of other species used for plant production in the nurseries.

Pest control measures applied during production

Pest control measures applied during the production of *Alnus* plants include: (a) registration of production sites; (b) the use of certified plant material; (c) surveillance, monitoring and sampling; (d) hygiene measures; (e) irrigation water testing; (f) washing of the roots of the field grown plants (up to 9 years old); (g) application of pest control products; (h) inspection and management of plants before export.

Evaluation of control measures

In general, the measures taken by the growers are effective against this pathogen. The following critical points were identified:

- The exposure time of large specimen tree can be up to 25 years of trees.
- The commodity is grown for up to 9 years in the field, before being transplanted into pots with growing media.
- The washing of the roots removes (parts of) the soil and the pathogen present in the soil, but it does not remove the pathogen present in the roots.
- Early infections are difficult to be detected by visual inspections.

Main uncertainties

- The extent and the distribution of the pathogen in the UK.
- Whether symptoms on *Alnus* spp. are distinguishable from *Phytophthora alni*.
- The efficiency of the hygiene measures especially concerning the cleaning of the machinery and the possible movement of soil within the nursery.

5.2.4 | Outcome of Expert Knowledge Elicitation

Table 9 and **Figure 6** show the outcome of the EKE regarding pest freedom after the evaluation of the currently proposed risk mitigation measures for the selected pests. Specifically, **Figure 6** provides the descending distribution function that describes the likelihood of pest freedom after evaluating the proposed risk mitigation measures for large *A. cordata* and *A. glutinosa* specimen trees (from 7 to 25 years old) in pots, designated for export to the EU, with respect to *E. mammata*, *P. ramorum* and *P. siskiyouensis* (**Figure 7**).

TABLE 9 Assessment of the likelihood of pest freedom following evaluation of current risk mitigation measures against pests on *Alnus cordata* and *A. glutinosa* plants designated for export to the EU. In panel A, the median value for the assessed level of pest freedom for each pest is indicated by 'M', the 5% percentile is indicated by 'L' and the 95% percentile is indicated by 'U'. The percentiles together span the 90% uncertainty range regarding pest freedom. The pest freedom categories are defined in panel B of the table.

Pest species	Sometimes pest free	More often than not pest free	Frequently pest free	Very frequently pest free	Extremely frequently pest free	Pest free with some exceptional cases	Pest free with few exceptional cases	Almost always pest free
1 <i>Entoleuca mammata</i>	L							
2 <i>Phytophthora ramorum</i>	L							
3 <i>Phytophthora siskiyouensis</i>							L	

PANEL A

Pest freedom category	Pest free plants out of 10,000
Sometimes pest free	≤ 5000
More often than not pest free	5000–≤ 9000
Frequently pest free	9000–≤ 9500
Very frequently pest free	9500–≤ 9900
Extremely frequently pest free	9900–≤ 9950
Pest free with some exceptional cases	9950–≤ 9990
Pest free with few exceptional cases	9990–≤ 9995
Almost always pest free	995–≤ 10,000

PANEL B

Legend of pest freedom categories

- L** Pest freedom category includes the elicited lower bound of the 90% uncertainty range
- M** Pest freedom category includes the elicited median
- U** Pest freedom category includes the elicited upper bound of the 90% uncertainty range




FIGURE 6 The elicited certainty (y-axis) of the number of pest-free *Alnus cordata* and *A. glutinosa* plants (x-axis; log-scaled) out of 10,000 plants designated for export to the EU introduced from UK for all evaluated pests visualised as descending distribution function. Horizontal lines indicate the percentiles (starting from the bottom 5%, 25%, 50%, 75%, 95%).




FIGURE 7 The explanation of the descending distribution function describing the likelihood of pest freedom for *Alnus cordata* and *A. glutinosa* plants designated for export to the EU based on the example of *Entoleuca mammata*.

6 | CONCLUSIONS

There are three pests (*E. mammata*, *P. ramorum* (non-EU isolates) and *P. siskiyouensis*) identified to be present in UK and considered to be potentially associated with the *A. cordata* and *A. glutinosa* plants imported from the UK and relevant for the EU. The likelihood of pest freedom after the evaluation of the implemented risk mitigation measures for large specimen trees (from 7 to 25 years old) of *A. cordata* and *A. glutinosa* designated for export to the EU was estimated.

For *E. mammata*, the likelihood of pest freedom following evaluation of current risk mitigation measures was estimated as 'pest free with few exceptional cases' with the 90% uncertainty range reaching from 'extremely frequently pest free' to 'almost always pest free'. The EKE indicated, with 95% certainty, that between 9905 and 10,000 plants per 10,000 will be free from *E. mammata*.

For *P. ramorum*, the likelihood of pest freedom following evaluation of current risk mitigation measures was estimated as 'pest free with some exceptional cases' with the 90% uncertainty range reaching from 'extremely frequently pest free' to 'almost always pest free'. The EKE indicated, with 95% certainty, that between 9920 and 10,000 plants per 10,000 will be free from *P. ramorum*.

For *P. siskiyouensis*, the likelihood of pest freedom following evaluation of current risk mitigation measures was estimated as 'almost always pest free' with the 90% uncertainty range reaching from 'pest free with few exceptional cases' to 'almost always pest free'. The EKE indicated, with 95% certainty, that between 9991 and 10,000 plants per 10,000 will be free from *P. siskiyouensis*.

GLOSSARY

Control (of a pest)	Suppression, containment or eradication of a pest population (FAO, 2024a, 2024b).
Entry (of a pest)	Movement of a pest into an area where it is not yet present, or present but not widely distributed and being officially controlled (FAO, 2024b).
Establishment (of a pest)	Perpetuation, for the foreseeable future, of a pest within an area after entry (FAO, 2024b).
Impact (of a pest)	The impact of the pest on the crop output and quality and on the environment in the occupied spatial units.
Introduction (of a pest)	The entry of a pest resulting in its establishment (FAO, 2024b).
Measures	Control (of a pest) is defined in ISPM 5 (FAO, 2024b) as 'Suppression, containment or eradication of a pest population' (FAO, 2024a). Control measures are measures that have a direct effect on pest abundance. Supporting measures are organisational measures or procedures supporting the choice of appropriate risk mitigation measures that do not directly affect pest abundance.
Pathway	Any means that allows the entry or spread of a pest (FAO, 2024b).
Phytosanitary measures	Any legislation, regulation or official procedure having the purpose to prevent the introduction or spread of quarantine pests, or to limit the economic impact of regulated non-quarantine pests (FAO, 2024b).
Protected zone	A Protected zone is an area recognised at EU level to be free from a harmful organism, which is established in one or more other parts of the Union.
Quarantine pest	A pest of potential economic importance to the area endangered thereby and not yet present there, or present but not widely distributed and being officially controlled (FAO, 2024b).
Regulated non-quarantine pest	A non-quarantine pest whose presence in plants for planting affects the intended use of those plants with an economically unacceptable impact and which is therefore regulated within the territory of the importing contracting party (FAO, 2024b).
Risk mitigation measure	A measure acting on pest introduction and/or pest spread and/or the magnitude of the biological impact of the pest should the pest be present. A risk mitigation measure may become a phytosanitary measure, action or procedure according to the decision of the risk manager.
Spread (of a pest)	Expansion of the geographical distribution of a pest within an area (FAO, 2024b).

ABBREVIATIONS

APHA	Animal and Plant Health Agency
CABI	Centre for Agriculture and Bioscience International
DEFRA	Department for Environment Food and Rural Affairs
EKE	Expert Knowledge Elicitation
EPPO	European and Mediterranean Plant Protection Organization
FAO	Food and agriculture organization
ISPM	International Standards for Phytosanitary Measures
NPPO	National Plant Protection Organisation
PLH	Plant health
PRA	Pest risk assessment

PZQPs	Protected zone quarantine pests
RNQPs	Regulated non-quarantine pests

REQUESTOR

European Commission

QUESTION NUMBER

EFSA-Q-2025-00040

COPYRIGHT FOR NON-EFSA CONTENT

EFSA may include images or other content for which it does not hold copyright. In such cases, EFSA indicates the copyright holder and users should seek permission to reproduce the content from the original source.

PANEL MEMBERS

Antonio Vicent Civera, Paula Baptista, Anna Berlin, Elisavet Chatzivassiliou, Jaime Cubero, Nik Cunniffe, Eduardo de la Peña, Nicolas Desneux, Francesco Di Serio, Anna Filipiak, Paolo Gonthier, Beata Hasiów-Jaroszewska, Hervé Jactel, Blanca B. Landa, Lara Maistrello, David Makowski, Panagiotis Milonas, Nikos T. Papadopoulos, Roel Potting, Hanna Susi, and Dirk Jan van der Gaag.

MAP DISCLAIMER

The designations employed and the presentation of material on any maps included in this scientific output do not imply the expression of any opinion whatsoever on the part of the European Food Safety Authority concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

REFERENCES

Anisimov, N. S., & Bezborodov, V. G. (2021). Longicorn beetles (Coleoptera: Disteniidae, Cerambycidae) of the Russky Island (Primorsky Krai, Russia). *Ecologica Montenegrina*, 40, 46–58.

Anonymous. (1960). Index of plant diseases in the United States. Agriculture Handbook no 165, USDA-ARS (US) 531 pp.

Atkinson, T. H. (online). Bark and Ambrosia Beetles. <http://www.barkbeetles.info/index.php> [Accessed: September 2024].

Barringer, L., & Ciafré, C. M. (2020). Worldwide feeding host plants of spotted lanternfly, with significant additions from North America. *Environmental Entomology*, 49(5), 999–1011. <https://doi.org/10.1093/ee/nvaa093>

Batsanksalashvili, M., Kaydan, M. B., Kirkadze, G., & Japoshvili, G. (2017). Updated checklist of scale insects (Hemiptera: Coccoimorpha) in Sakartvelo (Georgia). *Annals of Agrarian Science*, 15(2), 252–268. <https://doi.org/10.1016/j.aasci.2017.05.002>

Brito, R., Specht, A., Gonçalves, G. L., Moreira, G. R. P., Carneiro, E., Santos, F. L., Roque-Specht, V. F., Mielke, O. H. H., & Casagrande, M. M. (2019). Spodoptera marima: A new synonym of Spodoptera ornithogalli (Lepidoptera: Noctuidae), with notes on adult morphology, host plant use and genetic variation along its geographic range. *Neotropical Entomology*, 48(3), 433–448. <https://doi.org/10.1007/s13744-018-0654-z>

CABI (Centre for Agriculture and Bioscience International). (online). CABI Crop Protection Compendium. Available online: <https://www.cabidigitallibrary.org/product/qc> [Accessed: 20 September 2024].

Callan, B. E. (1998). *Diseases of Populus in British Columbia: A diagnostic manual* (p. 157). Natural Resources Canada, Canadian Forest Service, Pacific Forestry Centre.

Ciesla, W. M., & Kruse, J. J. (2009). Large aspen tortrix [revised]. USDA Forest Service, Forest Insect & Disease Leaflet 139 (revised), 8 pp.

Dubey, A. K., & Ko, C.-C. (2012). Sexual dimorphism among species of *Aleurocanthus* Quaintance & Baker (Hemiptera: Aleyrodidae) in Taiwan, with one new species and an identification key. *Zootaxa*, 3177, 1–23.

EFSA PLH Panel (EFSA Panel on Plant Health). (2018). Guidance on quantitative pest risk assessment. *EFSA Journal*, 16(8), 5350. <https://doi.org/10.2903/j.efsa.2018.5350>

EFSA PLH Panel (EFSA Panel on Plant Health). (2019). Guidance on commodity risk assessment for the evaluation of high risk plants dossiers. *EFSA Journal*, 17(4), 5668. <https://doi.org/10.2903/j.efsa.2019.5668>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Gonthier, P., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Potting, R., Stefani, E., Van der Werf, W., Vicent Civera, A., Yuen, J., Zappalà, L., ... Thulke, H.-H. (2024a). Standard protocols for plant health scientific assessments. *EFSA Journal*, 22(9), 8891. <https://doi.org/10.2903/j.efsa.2024.8891>

EFSA PLH Panel (EFSA Panel on Plant Health), Civera, A. V., Baptista, P., Berlin, A., Chatzivassiliou, E., Cubero, J., Cunniffe, N., de la Peña, E., Desneux, N., Di Serio, F., Filipiak, A., Hasiów-Jaroszewska, B., Jactel, H., Landa, B. B., Maistrello, L., Makowski, D., Milonas, P., Papadopoulos, N. T., Potting, R., ... Gonthier, P. (2024b). Commodity risk assessment of *Betula pendula* and *Betula pubescens* plants from the UK. *EFSA Journal*, 22(11), 9051. <https://doi.org/10.2903/j.efsa.2024.9051>

EFSA PLH Panel (EFSA Panel on Plant Health), Civera, A. V., Baptista, P., Berlin, A., Chatzivassiliou, E., Cubero, J., Cunniffe, N., de la Peña, E., Desneux, N., Di Serio, F., Filipiak, A., Gonthier, P., Hasiów-Jaroszewska, B., Jactel, H., Landa, B. B., Maistrello, L., Makowski, D., Milonas, P., Papadopoulos, N. T., ... Potting, R. (2025). Commodity risk assessment of *Alnus cordata*, *Alnus glutinosa* and *Alnus incana* plants from the UK. *EFSA Journal*, 23(1), 9189. <https://doi.org/10.2903/j.efsa.2025.9189>

EFSA Scientific Committee. (2018). Scientific opinion on the principles and methods behind EFSA's guidance on uncertainty analysis in scientific assessment. *EFSA Journal*, 16(1), 5122. <https://doi.org/10.2903/j.efsa.2018.5122>

EPPO (European and Mediterranean Plant Protection Organization). (online). EPPO Global Database. Available online: <https://gd.eppo.int/> [Accessed: 20 September 2024].

Eskalen, A., Stouthamer, R., Lynch, S. C., Rugman-Jones, P. F., Twizeyimana, M., Gonzalez, A., & Thibault, T. (2013). Host range of Fusarium dieback and its ambrosia beetle (Coleoptera: Scolytinae) vector in southern California. *Plant Disease*, 97(7), 938–951. <https://doi.org/10.1094/PDIS-11-12-1026-RE>

EUROPHYT. (online). European Union Notification System for Plant Health Interceptions – EUROPHYT. Available online: <https://ec.europa.eu/food/plants/plant-health-and-biosecurity/European-union-notification-system-plant-health-interceptions> [Accessed: 21/1/2025].

FAO (Food and Agriculture Organization of the United Nations). (2019). *ISPM (international standards for phytosanitary measures) No. 36. Integrated measures for plants for planting*. FAO. <https://www.ippc.int/en/publications/636/>

FAO (Food and Agriculture Organization of the United Nations). (2024a). *ISPM (international standards for phytosanitary measures) No 4. Requirements for the establishment of pest free areas*. FAO. <https://www.ippc.int/en/publications/614/>

FAO (Food and Agriculture Organization of the United Nations). (2024b). *ISPM (international standards for phytosanitary measures) No. 5. Glossary of phytosanitary terms*. FAO. <https://www.ippc.int/en/publications/622/>

Farashiani, M. E., Sadeghi, S. E., & Abaii, M. (2001). Geographic distribution and hosts of sart longhorn beetle, *Aeolesthes sarta* Solsky (Col.: Cerambycidae) in Iran. *Journal of Entomological Society of Iran*, 20, 81–96.

Ferguson, D. C. (1975). Host records for Lepidoptera reared in eastern North America. *USDA Technical Bulletin*, 1521, 1–49.

Fleming, W. E. (1972). Biology of the Japanese beetle. Technical Bulletin, Agricultural Research Service, USDA no 1449, 129 pp.

Gardi, C., Kaczmarek, A., Streissl, F., Civitelli, C., Do Vale Correia, C., Mikulová, A., Yuen, J., & Stancanelli, G. (2024). EFSA standard protocol for commodity risk assessment. *Zenodo*. <https://doi.org/10.5281/zenodo.13149775>

Gilligan, T. M., & Epstein, M. E. (2014). TortAI. Tortricidae of Agricultural Importance. https://idtools.org/id/leps/tortai/Acleris_senescens.htm

Lim, J., Jung, S.-Y., Lim, J.-S., Jang, J., Kim, K.-M., Lee, Y.-M., & Lee, B.-W. (2014). A review of host plants of Cerambycidae (Coleoptera: Chrysomeloidea) with new host records for fourteen Cerambycids, including the Asian longhorn beetle (*Anoplophora glabripennis* Motschulsky), in Korea. *Korean Journal of Applied Entomology*, 53(2), 111–133. <https://doi.org/10.5656/KSAE.2013.11.1.061>

Malembic-Maher, S., Desqué, D., Khalil, D., Salar, P., Bergey, B., Danet, J.-L., Duret, S., & Foissac, X. (2020). When a Palearctic bacterium meets a Nearctic insect vector: Genetic and ecological insights into the emergence of the grapevine Flavescence dorée epidemics in Europe. *PLoS Pathogens*, 16(3), e1007967. <https://doi.org/10.1371/journal.ppat.1007967>

Mehle, N., Rupar, M., Seljak, G., Ravnikar, M., & Dermastia, M. (2011). Molecular diversity of 'flavescence doree' phytoplasma strains in Slovenia. *Bulletin of Insectology*, 64, S29–S30.

O'Hanlon, R., Choiseul, J., Corrigan, M., Catarame, T., & Destefanis, M. (2016). Diversity and detections of Phytophthora species from trade and non-trade environments in Ireland. *EPPO Bulletin*, 46(3), 594–602. <https://doi.org/10.1111/epp.12331>

Palmer, W. A. (1987). The phytophagous insect fauna associated with *Baccharis halimifolia* L. and *B. Neglecta* Britton in Texas, Louisiana, and northern Mexico. *Proceedings of the Entomological Society of Washington*, 89, 185–199.

Park, J. D., Kim, M.-Y., Lee, S.-G., Shin, S.-C., Kim, J., & Park, I.-K. (2009). Biological characteristics of *Lycorma delicatula* and the control effects of some insecticides. *Korean Journal of Applied Entomology*, 48(1), 53–57. <https://doi.org/10.5656/KSAE.2009.48.1.053>

Plant-SyNZ. (online). Landcare Research (NZ). Host plants of a herbivore -Oemona hirta. <https://plant-synt.landcareresearch.co.nz/index.asp> [Accessed: June 2021].

Prentice, R. M. (1966). *Forest Lepidoptera of Canada recorded by the Forest insect survey. Volume 4, Publication 1142* (pp. 543–840). Microlepidoptera, Department of Forestry.

Radonjic, S., Hrncic, S., Krstic, O., Cvrkovic, T., Mitrovic, M., Jovic, J., & Tosevski, I. (2013). First report of Alder yellows phytoplasma infecting common and grey alder (*Alnus glutinosa* and *A. Incana*) in Montenegro. *Plant Disease*, 97(5), 686–686. <https://doi.org/10.1094/PDIS-11-12-1087-PDN>

Regione Lombardia Servizio Fitosanitario. (online). Organismi nocivi, *Popillia* o Japanese beetle (*Popillia japonica*) - Ispezioni Visive- Principali piante ospiti. *Popillia o Japanese beetle (Popillia japonica)* [Accessed: 30 September, 2024].

Scalenet. (online). García Morales, M., Denno, B. D., Miller, D. R., Miller, G. L., Ben-Dov, Y., & Hardy, N. B. (online). *ScaleNet: A literature-based model of scale insect biology and systematics*. Database. [Accessed: 20 September 2024].

Shrewsbury, P. M., Harding, N. M., Rojas, M. S., & Gill, S. (2013). Japanese maple scale: Woody ornamental host plants. UMD Extension Publication EBR-18 2013. <https://extension.umd.edu/sites/extension.umd.edu/files/publications/Japanese%20Maple%20Scale%20%282%29.pdf>

Sjöman, H., Östberg, J., & Nilsson, J. (2014). Review of host trees for the wood-boring pests *Anoplophora glabripennis* and *Anoplophora chinensis*: an urban forest perspective. *Arboriculture & Urban Forestry*, 40(3), 143–164.

TRACES-NT. (online). Trade Control and Expert System. <https://webgate.ec.europa.eu/tracesnt> [Accessed: 21/1/2025].

Turchetti, T., Maresi, G., & Santagada, A. (1991). *Cryphonectria parasitica* blight on different hosts in the Cilento region, S. Italy.

USDA Fungal Databases (USDA). (online). U.S. Department of agriculture. <https://fungi.ars.usda.gov/> [Accessed: 20 September 2024]

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: EFSA PLH Panel (EFSA Panel on Plant Health), Civera, A. V., Baptista, P., Berlin, A., Chatzivassiliou, E., Cubero, J., Cunniffe, N., de la Peña, E., Desneux, N., Di Serio, F., Filipiak, A., Gonthier, P., Hasiów-Jaroszewska, B., Jactel, H., Landa, B. B., Maistrello, L., Makowski, D., Milonas, P., Papadopoulos, N. T., ... Potting, R. (2025). Commodity risk assessment of *Alnus cordata* and *Alnus glutinosa* specimen trees from the UK. *EFSA Journal*, 23(4), e9383. <https://doi.org/10.2903/j.efsa.2025.9383>

APPENDIX A

Datasheets of pests selected for further evaluation

A.1 | ENTOLEUCA MAMMATA

A.1.1 | Organism information

Taxonomic information	Current valid scientific name: <i>Entoleuca mammata</i> (Wahlenb.) J.D. Rogers & Y.M. Ju Synonyms: <i>Anthostoma blakei</i> , <i>Anthostoma morsei</i> , <i>Fuckelia morsei</i> , <i>Hypoxylon blakei</i> , <i>Hypoxylon holwayi</i> , <i>Hypoxylon mammatum</i> , <i>Hypoxylon morsei</i> , <i>Hypoxylon pauperatum</i> , <i>Hypoxylon pruinatum</i> , <i>Nemania mammata</i> , <i>Rosellinia pruinata</i> , <i>Sphaeria mammata</i> , <i>Sphaeria pruinata</i> (according to Index Fungorum) Name used in the EU legislation: <i>Entoleuca mammata</i> (Wahlenb.) Rogers and Ju Order: Xylariales Family: Xylariaceae Common name: hypoxylon canker of poplar, canker of aspen
Group	Fungi
EPPO code	HYPOMA
Regulated status	<i>Entoleuca mammata</i> is listed in Annex III of Commission Implementing Regulation (EU) 2019/2072 as protected zone quarantine pest for Ireland. The pathogen is quarantine pest in China and Israel. It is on the A1 list of Türkiye (EPPO, online_a).
Pest status in the UK	<i>Entoleuca mammata</i> is present in the UK, with few occurrences in England, Wales, Channel Islands and Scotland (CABI, online; EPPO, online_b).
Pest status in the EU	Not relevant, <i>E. mammata</i> is an EU-regulated pest.
Host status on Alnus spp.	<i>Entoleuca mammata</i> has been reported on various species of <i>Alnus</i> . These included <i>Alnus</i> spp. (Callan, 1998; French et al., 1969; Goos, 2010; Hawksworth, 1972; <i>A. crispa</i> var. <i>mollis</i> (Connors, 1967), <i>A. crispa</i> var. <i>sinuata</i> (Ginns, 1986), <i>A. incana</i> (Anonymous, 1960), <i>A. rugosa</i> (Connors, 1967; Ginns, 1986), <i>A. sinuata</i> (Callan, 1998; Connors, 1967) and <i>A. tenuifolia</i> (Anonymous, 1960). Given the fact that has been reported in several <i>Alnus</i> spp., the Panel assumes that <i>A. cordata</i> and <i>A. glutinosa</i> can be a host or <i>E. mammata</i> .
Risk assessment information	Pest Risk Assessments available: <ul style="list-style-type: none">– Scientific Opinion on the pest categorisation of <i>Entoleuca mammata</i> (EFSA PLH Panel, 2017).– UK Risk Register Details for <i>Entoleuca mammata</i> (DEFRA, online).– Express Pest Risk Analysis: <i>Entoleuca mammata</i> (Klejdysz et al., online).– Commodity risk assessment of <i>Acer campestris</i>, <i>A. platanoides</i> and <i>A. pseudoplatanus</i> plants from the UK (EFSA PLH Panel, 2023a, 2023b, 2023c).– Commodity risk assessment of <i>Sorbus aucuparia</i> plants from the UK (EFSA PLH Panel, 2024a, 2024b).– Commodity risk assessment of <i>Alnus cordata</i>, <i>A. glutinosa</i> and <i>A. incana</i> from the UK (EFSA PLH Panel, 2025).
Other relevant information for the assessment	
Biology	<i>Entoleuca mammata</i> is an ascomycete fungus known as an important agent of canker disease in <i>Populus</i> species, mostly <i>Populus tremuloides</i> and <i>P. tremula</i> ; other hardwood species like <i>Salix</i> spp. can also be infected (EFSA PLH Panel, 2017). The pathogen is native to North America and was introduced to Europe several centuries ago (Kasanen et al., 2004); the ascospores of <i>E. mammata</i> can infect the living wood of the hosts penetrating the periderm and invading tissues under healthy bark and through mechanical wounds, as well as through injuries caused by woodpeckers and insects, in particular the North American cerambycid beetles (mostly <i>Saperda inornata</i> and <i>Oberea</i> spp.) (Anderson et al., 1979a) and the cicada <i>Magicicada septendecim</i> (Ostry and Anderson, 1983) water stress can increase host susceptibility (EFSA PLH Panel, 2017). <i>Entoleuca mammata</i> is mostly found on trees 15–40 years old, but all ages can be infected (EFSA PLH Panel, 2017; EPPO online_c). Infection usually starts from branches and twigs and then can spread to the main stem. The cankers expand very rapidly (7–8 cm per month) in summer and more slowly during winter; branches and stems can be girdled causing drying and breakage. The pathogen mostly develops in the range from 8°C to 32°C, the optimum temperature is 28°C; toxins host-specific produced by the fungus are involved in pathogenesis (EFSA PLH Panel, 2017; EPPO, online_c; Stermer et al., 1984). <i>Entoleuca mammata</i> overwinters in host tissues as both mycelium and spores. Conidia are produced 5–14 months after infection, but their role in the disease transmission is considered not relevant (EFSA PLH Panel, 2017). The pathogen spreads over long distances via windborne ascospores, which are produced only 2–3 years after infection; cankers on felled trees on the ground can continue to produce ascospores for 23 months. Ascospores are dispersed at a temperature above –4°C and in wet weather; a minimum of 16°C is required for starting germination, which became rapid at 28–32°C (EFSA PLH Panel, 2017). Infected wood, mostly with bark, maybe a pathway for the passive spread of <i>E. mammata</i> in international trade; however, also young plants may carry ascospores or mycelium of the fungus, which can exist as a latent infection on living material inadvertently moved (EFSA PLH Panel, 2017; EPPO online_c).

(Continues)

(Continued)

Symptoms	Main type of symptoms	The symptoms are observed on <i>Populus</i> trees. Early symptoms of cankers on the bark appear as slightly sunken, yellowish-orange areas with an irregular border. Young cankers can be easily identified by removing the bark to expose the white mycelium in the cambial zone. The outer bark in older cankers is then lifted into blister-like patches and breaks away, exposing blackened areas prominently visible on green branches and trunks. Callus formation only occasionally develops because cankers spread very quickly (Anderson et al., 1979b; EPPO, online_c). Wilting of leaves may be observed when the trees are girdled, as well as sprouting of new shoots on stems and branches. Infected trees can be secondarily colonised by other fungi, accelerating the host decline (EPPO, online_c). There is no information on the symptoms caused to <i>Alnus</i> plants.
	Presence of asymptomatic plants	Infections by <i>E. mammata</i> have an incubation period, with symptoms typically appearing on average 2 years after ascospore infection; therefore, asymptomatic plants can be found (Ostry and Anderson, 2009).
	Confusion with other pests	Some <i>Hypoxyylon</i> species present in Europe on deciduous trees (<i>H. confluens</i> and <i>H. udum</i>) show symptoms similar to those caused by <i>E. mammata</i> but can be easily distinguished in the laboratory by the ascospore characteristics (EFSA PLH Panel, 2017).
Host plant range		In North America, <i>E. mammata</i> mainly infects quaking aspen (<i>Populus tremuloides</i>); minor damage is recorded on <i>P. grandidentata</i> , <i>P. balsamifera</i> and various <i>Populus</i> hybrids. Other reported hosts in North America are <i>Acer</i> , <i>Alnus</i> , <i>Betula</i> , <i>Carpinus</i> , <i>Fagus</i> , <i>Picea</i> , <i>Pyrus</i> , <i>Salix</i> , <i>Sorbus</i> and <i>Ulmus</i> (Manion and Griffin, 1986). In Europe, the main host is <i>Populus tremula</i> ; other hosts are <i>Populus alba</i> , <i>P. nigra</i> , <i>P. trichocarpa</i> and the hybrid <i>P. tremula</i> x <i>P. tremuloides</i> (Ostry, 2013). The fungus is reported in <i>Salix</i> sp. (Eriksson, 2014; Vasilyeva and Scheuer, 1996) and <i>S. aucuparia</i> (Eriksson, 2014; Vasilyeva and Scheuer, 1996). There are no reports on <i>Alnus</i> spp. in Europe.
Reported evidence of impact		The fungus is an EU regulated pest.
Evidence that the commodity is a pathway		Plants for planting of <i>Alnus</i> species may carry ascospores and mycelium of <i>E. mammata</i> also as asymptomatic plants (EFSA PLH Panel; EPPO online_c). Mechanical wounds including pruning may facilitate infection courts. Plants in pots with or without leaves can be a pathway, because the ascospore or mycelium can be found in the branches.
Surveillance information		<i>Entoleuca mammata</i> is not a regulated pest for UK and it is not under official control and surveillance (Dossier Section 1).

A.1.2 | Possibility of pest presence in the nursery

A.1.2.1 | Possibility of entry from the surrounding environment

Entoleuca mammata is present in the UK in England, Wales, Channel Islands and Scotland (CABI, online; EPPO, online_b). The pathogen can naturally spread with ascospores dispersed by air currents also over long distance.

Exporting nurseries are predominately situated in the rural areas. The surrounding land would tend to be arable farmland with some pasture for animals and small areas of woodland. Hedges are often used to define field boundaries and grown along roadsides. Woodlands tend to be a standard UK mixed woodland with a range of UK native trees, that include host plants for the fungus such as oak (*Quercus robur*), poplar (*Populus* spp.). Hedges are made up of a range of species, including *Acer* and *Alnus* (Dossier Section 1).

Uncertainties:

- The presence of the pathogen on host plants in the surrounding area.

Taking into consideration the above evidence and uncertainties, the Panel considers that it is possible for *E. mammata* to enter the nurseries from the surrounding environment via ascospores transported by wind and air currents.

A.1.2.2 | Possibility of entry with new plants/seeds

Plants are produced by seeds and grafting. Seeds purchased in the UK are certified under The Forest Reproductive Material (Great Britain) Regulations 2002 ([legislation.gov.uk](https://www.legislation.gov.uk)); seedlings sourced in the UK are certified with UK Plant Passports; a small percentage of plants may be obtained from EU (Netherlands); seedlings from the EU countries are certified with phytosanitary certificates.

The pathogen is not known to be seedborne or seed transmitted, therefore not expected to enter the nursery via the seed pathway. Grafted plants (scion and buds in the case of grafting) originate from the nursery itself.

The seedling pathways are unlikely because the conditions of their production (seedbed in the greenhouse, pest free growing media) are expected to prevent the infection of seedlings.

The nurseries use virgin peat or peat-free compost (a mixture of coir, tree bark, wood fibre, etc.) as a growing media (Dossier Section 1).

The growing media is heat-treated by commercial suppliers during production to eliminate pests and diseases (Dossier Section 1). There is no evidence that soil or growing media may be a pathway for *E. mammata*.

Uncertainties

- None.

Taking into consideration the above evidence and uncertainties, the Panel considers that it is unlikely for the pathogen to enter the nurseries via seeds or seedlings of *A. cordata* and *A. glutinosa*.

A.1.2.3 | Possibility of spread within the nursery

Host plants of *E. mammata* are grown outdoors, in the open air, or field, in the nurseries producing *Alnus* spp. plants (Dossier Section 1). Therefore, is it possible that *E. mammata* can spread within the nursery from infested host plants of other species to the plot with *Alnus* spp. plants. *Entoleuca mammata* could spread from these other host plant species to *Alnus* plants via ascospores by air currents.

Uncertainties

- Efficiency in detecting and removing *E. mammata* infected plants.

Taking into consideration the above evidence and uncertainties, the Panel considers that the spread of the pathogen within the nurseries is possible by air currents.

A.1.3 | Information from interceptions

In the EUROPHYT/TRACES-NT database, there are no records of notification of *Alnus* plants for planting neither from the UK nor from other countries due to the presence of *E. mammata* between the years 1995 and January 2025 (EUROPHYT/TRACES-NT, online).

A.1.4 | Evaluation of the risk mitigation measures

In the table below, all risk mitigation measures currently applied in the UK are listed and an indication of their effectiveness on *E. mammata* is provided. The description of the implemented risk mitigation measures is provided in [Table 8](#).

No.	Risk mitigation measure	Effect on the pest	Evaluation and uncertainties
1	Registration of production sites	Yes	<p>All nurseries are registered as professional operator with the UK NPPO, by the Animal Plant Health Agency (APHA) for England and Wales, or with SASA for Scotland, and are authorised to issue UK plant passports (Dossier Section 1).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – The risk mitigation measure is expected to be effective in reducing the likelihood of presence of the pathogen on the commodity. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – Whether early symptoms on <i>Alnus</i> spp. are easily recognisable.
2	Certification of propagation material	Yes	<p><i>Alnus cordata</i> and <i>A. glutinosa</i> seeds purchased in the UK are certified under The Forest Reproductive Material (Great Britain) Regulations 2002 (legislation.gov.uk); seedlings sourced in the UK are certified with UK Plant Passports; a small percentage of plants may be obtained from EU (Netherlands); seedlings from the EU countries are certified with phytosanitary certificates. (Dossier Section 1).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – The risk mitigation measure is expected to be effective in reducing the likelihood of presence of the pathogen on the commodity. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – None.
3	Origin and treatment of growing media	No	<p>Rooted plants in pots: In the production or procurement of these plants, the use of growing media is assessed for the potential to harbour and transmit plant pests. Growers most commonly use virgin peat or peat-free compost, which is a mixture of coir, tree bark, wood fibre, etc. The compost is heat-treated by commercial suppliers during production to eliminate pests and diseases. It is supplied in sealed bulk bags or shrink-wrapped bales and stored off the ground on pallets, these are completely hygienic and free from contamination. Where delivered in bulk, compost is kept in a dedicated bunker, either indoors, or covered by tarpaulin outdoors, and with no risk of contamination with soil or other material (Dossier Section 1).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – Not relevant because the fungus is not reported as a soil-borne pathogen.

(Continues)

(Continued)

No.	Risk mitigation measure	Effect on the pest	Evaluation and uncertainties
4	Surveillance, monitoring and sampling	Yes	<p>Inspection is carried out at least once a year as part of the Quarantine Surveillance programme (Great Britain uses the same framework for its surveillance programme as the EU). Surveillance is based on visual inspection with samples taken from symptomatic material, and where appropriate, samples are also taken from asymptomatic material (e.g. plants, tubers, soil, watercourses) (Dossier Section 1).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – If infected plants are detected and removed from the nursery, this measure could have some effect. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – Whether symptoms caused by the pathogen on <i>Alnus</i> spp. are recognisable. – Whether <i>Alnus</i> spp. plants are subjected to annual surveys.
5	Hygiene measures	Yes	<p>According to the Dossier Section 1, all the nurseries have plant hygiene and housekeeping rules and practices in place, which are communicated to all relevant employees. These measures include:</p> <ul style="list-style-type: none"> – Cleaning and sterilisation of tools. – Waste treatment and disposal. <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – It is highly unlikely that the fungus spreads by the pruning tools. – The correct disposal of infected plant material prevents the spread of the fungus. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – The efficiency of hygiene measures performed in the nurseries.
6	Irrigation water quality and/or treatments	No	<p>Growers are required to assess water sources, irrigation and drainage systems used in the plant production for the potential to harbour and transmit plant pests. Water is routinely sampled and sent for analysis. No quarantine pests have been found (Dossier Section 1).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – The fungus is not spread by irrigation water. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – None.
7	Application of pest control products	Yes	<p>Crop protection is achieved using a combination of measures including approved plant protection products, biological control or physical measures. Plant protection products are only used when necessary and records of all plant protection treatments are kept (Dossier Section 1).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – Although <i>E. mammata</i> is generally not a target of the pesticide treatments in the nurseries, some protection products could reduce the likelihood of the infection by the pathogen. – The efficacy of plant protection products on large trees (up to 25 years old) could be reduced. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – No specific information on the efficacy of the plant protection products used.
8	Washing of the roots	No	<p>Specimen trees (up to 25 years old) are grown in soil for up to 9 years, root washed and placed in pots (based on the request letter of DEFRA to the EU Commission on September 25th, 2023).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – Not relevant because the fungus is not reported as a soil-borne pathogen.
9	Inspections and management of plants before export	Yes	<p>The UK NPPO carries out inspections and testing where required by the country of destination's plant health legislation, to ensure all requirements are fulfilled and a valid phytosanitary certificate with the correct additional declarations is issued. Separate to any official inspection, plant material is checked by growers for plant health issues before dispatch.</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – The inspections and management of plants before export can detect the pathogen. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – Whether early symptoms caused by the pathogen on <i>Alnus</i> species are identified by visual inspections.

A.1.5 | Overall likelihood of pest freedom for the exported commodity (comparative)

A.1.5.1 | Comparison with other relevant commodity Risk Assessments involving *Entoleuca mammata*

Entoleuca mammata was already assessed as actionable pest for graftwood, bare-root plants, cell-grown plants and rooted plants (of up to 7 years old) in pots of *A. cordata*, *A. glutinosa* and *A. incana* (EFSA PLH Panel, 2025). *Alnus* commodity types evaluated originate from nurseries sharing the same production system. Therefore, for the estimation of the pest freedom level of *E. mammata* in exported specimen trees (from 7 to 25 years old) of *Alnus* plants, the Panel decided to base the estimated values on the ones elicited for the other commodities of *A. cordata*, *A. glutinosa* and *A. incana* (EFSA PLH Panel,

2025). In addition, because of the similarity of the commodities, the production systems, the locations of the nurseries, the Panel also considered the EKE values from the previous Scientific opinion on *Betula pendula* and *B. pubescens* from the UK (EFSA PLH Panel, 2024a, 2024b).

A.1.6 | Overall likelihood of pest freedom for specimen trees (from 7 to 25 years old)

A.1.6.1 | Reasoning for a scenario which would lead to a reasonably low number of infected specimen trees (from 7 to 25 years old)

The scenario assumes a low pressure of the pathogen in the nurseries and in the surroundings. There are no host records for *A. cordata* and *A. glutinosa*. The scenario assumes *A. cordata* and *A. glutinosa* to be minor hosts for the pathogen. The scenario also assumes that symptoms of the disease are visible and promptly detected during inspections, resulting in removal of infected trees.

A.1.6.2 | Reasoning for a scenario which would lead to a reasonably high number of infected specimen trees (from 7 to 25 years old)

The scenario assumes a high pressure of the pathogen in the surrounding environment of the nurseries because suitable hosts are present. The scenario assumes that the pathogen can infect leaves, which may still be present on the plants at the time of export. The scenario also assumes that the pathogen is not detected during the inspections because of presence of asymptomatic plants or difficulties in recognising early symptoms. Grafting can increase the incidence of the pathogen (via infected buds or by woundings).

In comparison with the other commodities (graftwood, bare-root plants, cell-grown plants and rooted plants in pots), the canopy size of large specimen trees (up to 25 years old) is larger, increasing the difficulties in detecting the pathogen. Furthermore, the exposure time is longer, increasing the probability for the plants to be infected.

A.1.6.3 | Reasoning for a central scenario equally likely to over- or underestimate the number of infected specimen trees (from 7 to 25 years old) (Median)

In comparison to potted plants of up to 7 years old, the specimen trees are generally and older up to 25 years old and larger, increasing the time of exposure to the pathogen. The scenario assumes a limited presence of the pathogen in the nurseries and in the surroundings, and a limited reported susceptibility of *A. cordata* and *A. glutinosa*.

A.1.6.4 | Reasoning for the precision of the judgement describing the remaining uncertainties (1st and 3rd quartile/interquartile range)

The limited information on the occurrence of the pathogen in the nurseries and the surroundings and the susceptibility of *A. cordata* and *A. glutinosa*. Results in high level of uncertainties.

A.1.7 | Elicitation outcomes of the assessment of the pest freedom for *Entoleuca mammata* specimen trees (from 7 to 25 years old)

The elicited and fitted values for *E. mammata* for pest infestation and pest freedom agreed by the Panel are shown in Tables A.1 and A.2 and in Figure A.1.

TABLE A.1 Elicited and fitted values of the uncertainty distribution of pest infestation by *Entoleuca mammata* per 10,000 plants of specimen trees (from 7 to 25 years old).

Percentile	1%	2.5%	5%	10%	17%	25%	33%	50%	67%	75%	83%	90%	95%	97.5%	99%
Elicited values	0					14		25		50					140
EKE	0.961	1.94	3.34	5.85	9.04	13.0	17.2	26.8	39.5	48.2	60.2	75.0	94.8	114	139

Note: The EKE results is the *BetaGeneral* (1.3319, 379.19, 0, 10,000) distribution fitted with @Risk version 7.6.

Based on the numbers of estimated infected plants the pest freedom was calculated (i.e. = 10,000 – number of infected plants per 10,000). The fitted values of the uncertainty distribution of the pest freedom are shown in Table A.2.

TABLE A.2 The uncertainty distribution of plants free of *Entoleuca mammata* per 10,000 plants of specimen trees (from 7 to 25 years old) calculated by Table A.1.

Percentile	1%	2.5%	5%	10%	17%	25%	33%	50%	67%	75%	83%	90%	95%	97.5%	99%
Values	9860					9950		9975		9986					10,000
EKE results	9861	9886	9905	9925	9940	9952	9960	9973	9983	9987	9991	9994	9997	9998	9999

Note: The EKE results are the fitted values.




FIGURE A.1 (Continued)

(B)




FIGURE A.1 (Continued)




FIGURE A.1 (A) Elicited uncertainty of pest infection per 10,000 plants of specimen trees (histogram in blue – vertical blue line indicates the elicited percentile in the following order: 1%, 25%, 50%, 75%, 99%) and distributional fit (red line); (B) uncertainty of the proportion of pest-free plants per 10,000 (i.e. = 1 – pest infection proportion expressed as percentage); (C) descending uncertainty distribution function of pest infection per 10,000 plant.

A.1.8 | Reference list

Anderson N. A., Ostry, M. E., & Anderson, G. W. (1979a). Insect wounds as infection sites for Hypoxylon mammatum on trembling aspen. *Phytopathology*, 69, 476–479. <https://doi.org/10.1094/phyto-69-476>

Anderson R. L., Anderson, G. W., & Schipper Jr, A. L. (1979b). Hypoxylon canker of aspen. *USDA Forest Insect and Disease Leaflet*, 6, 6.

Anonymous. (1960). Index of Plant Diseases in the United States. *USDA Agriculture Handbook*, 165, 1–531.

CABI (Centre for Agriculture and Bioscience International). (online). Hypoxylon mammatum (poplar canker). Available online: <https://www.cabi.org/cpc/datasheet/28323> [Accessed: 12 October 2024].

Callan, B. E. (1988). Diseases of *Populus* in British Columbia: A Diagnostic Manual. Natural Resources Canada, Canadian Forest Service, Pacific Forestry Centre, Victoria, BC. 157.

Connors I. L. (1967). An annotated index of plant diseases in Canada and fungi recorded on plants in Alaska, Canada and Greenland. *Research Branch Canada Department of Agriculture*, 1251, 1–381

DEFRA (Department for Environment, Food and Rural Affairs). (online). UK risk register details for Entoleuca mammata. Available online: <https://plant-healthportal.defra.gov.uk/pests-and-diseases/uk-plant-health-risk-register/viewPestRisks.cfm?csref=11840> [Accessed: 26 February 2024].

EFSA PLH Panel (EFSA Panel on Plant Health), Jeger, M., Bragard, C., Caffier, D., Candresse, T., Chatzivassiliou, E., Dehnen-Schmutz, K., Gilioli, G., Gregoire, J.-C., Jaques Miret, J.A., MacLeod, A., Navajas Navarro, M., Niere, B., Parnell, S., Potting, R., Rafoss, T., Rossi, V., Urek, G., Van Bruggen, A., ... Pautasso, M. (2017). Scientific Opinion on the pest categorisation of Entoleuca mammata. *EFSA Journal*, 15(7), 4925. <https://doi.org/10.2903/j.efsa.2017.4925>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Potting, R., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent-Civera, A., Yuen, J., Zappalà, L., ... Gonthier, P. (2023a). Scientific Opinion on the commodity risk assessment of Acer campestre plants from the UK. *EFSA Journal*, 21(7), 8071. <https://doi.org/10.2903/j.efsa.2023.8071>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Potting, R., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent-Civera, A., Yuen, J., Zappalà, L., ... Gonthier, P. (2023b). Scientific Opinion on the commodity risk assessment of Acer platanoides plants from the UK. *EFSA Journal*, 21(7), 8073. <https://doi.org/10.2903/j.efsa.2023.8073>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Potting, R., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent-Civera, A., Yuen, J., Zappalà, L., ... Gonthier, P. (2023c). Scientific Opinion on the commodity risk assessment of Acer pseudoplatanus plants from the UK. *EFSA Journal*, 21(7), 8074. <https://doi.org/10.2903/j.efsa.2023.8074>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Potting, R., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent-Civera, A., Yuen, J., Zappalà, L., ... Gonthier, P. (2024a). Commodity risk assessment of Betula pendula and Betula pubescens plants from the UK. *EFSA Journal*, 22(11), e9051. <https://doi.org/10.2903/j.efsa.2024.9051>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Potting, R., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent-Civera, A., Yuen, J., Zappalà, L., ... Gonthier, P. (2024b). Commodity risk assessment of Sorbus aucuparia plants from the UK. *EFSA Journal*, 22(6), e8837. <https://doi.org/10.2903/j.efsa.2024.8837>

EFSA PLH Panel (EFSA Panel on Plant Health), Civera, A. V., Baptista, P., Berlin, A., Chatzivassiliou, E., Cubero, J., Cunniffe, N., de la Peña, E., Desneux, N., Di Serio, F., Filipiak, A., Gonthier, P., Hasiów-Jarzewska, B., Jactel H. Landa, B. B., Maistrello, L., Makowski, D., Milonas, P., Papadopoulos, N. T., ... Potting, R. (2025). Commodity risk assessment of *Alnus cordata*, *Alnus glutinosa* and *Alnus incana* plants from the UK. *EFSA Journal*, 23(1), e9189. <https://doi.org/10.2903/j.efsa.2025.9189>

EPPO (European and Mediterranean Plant Protection Organization). (online_a). Entoleuca mammata (HYPOMA), Categorization. Available online: <https://gd.eppo.int/taxon/HYPOMA/categorization> [Accessed: 26 October 2024].

EPPO (European and Mediterranean Plant Protection Organization) (online_b). Entoleuca mammata (HYPOMA), Distribution. Available online: <https://gd.eppo.int/taxon/HYPOMA/distribution> [Accessed: 26 October 2024].

EPPO (European and Mediterranean Plant Protection Organization). (online_c). Datasheets on Quarantine Pests: Hypoxylon mammatum (HYPOMA), Documents. Available online: <https://gd.eppo.int/taxon/HYPOMA/documents> [Accessed: 26 October 2024].

Eriksson, O. E. (2014). Checklist of the non-lichenized ascomycetes of Sweden. *Acta Universitatis Upsaliensis Symbolae Botanicae Upsalienses*, 36, 499.

EUROPHYT. (online). European Union Notification System for Plant Health Interceptions – EUROPHYT. Available online: http://ec.europa.eu/food/plant/plant_health_biosecurity/europhyt/index_en.htm [Accessed: 21/1/2025].

French, D. W., Hodges Jr, C. S., & Froyd, J. D. (1969). Pathogenicity and taxonomy of Hypoxylon mammatum. *Canadian Journal of Botany*, 47, 223–226. <https://doi.org/10.1139/b69-029>

Ginns, J. H. (1986). Compendium of plant disease and decay fungi in Canada 1960–1980. *Research Branch Canada Department of Agriculture Publication*, 1813, 416.

Goos, R. D. (2010). The mycota of Rhode Island: A checklist of the fungi recorded in Rhode Island (including lichens and myxomycetes) Vol. 4. *Rhode Island Natural History Survey* 222.

Hawksworth, D. L. (1972). Hypoxylon mammatum. CMI Descriptions of Pathogenic Fungi and Bacteria No. 356. CABI, Wallingford, UK.

Kasanen, R., Hantula, J., Ostry, M. E., Pinon, J., & Kurnkela, T. (2004). North American populations of Entoleuca mammata are genetically more variable than populations in Europe. *Mycological Research*, 108, 766–774. <https://doi.org/10.1017/s0953756204000334>

Klejdys, T., Kubasik, W., Strażyński, P., Gawlik, M., Pruciak, A., Rzepecka, D., Kałuski, T. (online). Express pest risk analysis for Hypoxylon mammatum. Available online: https://www.plantquarantine.pl/user_storage/36/pliki/pra_133/Entoleuca%20mammata.pdf [Accessed: 2 October 2024].

Manion P. D., & Griffin, D. H. (1986). Sixty-five years of research on Hypoxylon canker of aspen. *Plant Disease*, 70(8), 803–808. <https://doi.org/10.1094/pd-70-803>

Ostry, M. E. (2013). Hypoxylon canker Gonthier P Nicolotti G. *Infectious Forest Diseases Cabi International Wallingford*, 407, 419

Ostry, M. E., & Anderson, N. A. (1983). Infection of trembling aspen by Hypoxylon mammatum through cicada oviposition wounds. *Phytopathology*, 73, 1092–1096. <https://doi.org/10.1094/phyto-73-1092>

Ostry, M. E., & Anderson, N. A. (2009). Genetics and ecology of the Entoleuca mammata–Populus pathosystem: implications for aspen improvement and management. *Forest Ecology and Management*, 257, 390–400. <https://doi.org/10.1016/j.foreco.2008.09.053>

Stermer, B. A., Scheffer, R. P., & Hart, J. H. (1984). Isolation of toxins from Hypoxylon mammatum and demonstration of some toxin effects on selected clones of *Populus tremuloides*. *Phytopathology*, 74, 654–658. <https://doi.org/10.1094/phyto-74-654>

TRACES-NT. (online). TRAde Control and Expert System. Available online: <https://webgate.ec.europa.eu/tracesnt> [Accessed: 21/1/2025].

Vasilyeva, L. N., & Scheuer, C. (1996). Recent collections of stromatic pyrenomycetes from Austria, especially from Styria Mitt. Naturwiss. Vereins Steiermark, 126, 61–82.

A.2 | PHYTOPHTHORA RAMORUM

A.2.1 | Organism information

Taxonomic information	<p>Current valid scientific name: <i>Phytophthora ramorum</i> Werres, De Cock & Man in 't Veld</p> <p>Synonyms: –</p> <p>Name used in the EU legislation: <i>Phytophthora ramorum</i> (non-EU isolates) Werres, De Cock & Man in 't Veld [PHYTRA]</p> <p>Order: Peronosporales</p> <p>Family: Peronosporaceae</p> <p>Common name: Sudden Oak Death (SOD), ramorum bleeding canker, ramorum blight, ramorum leaf blight, twig and leaf blight</p> <p>Name used in the Dossier: <i>Phytophthora ramorum</i></p>
Group	Oomycetes
EPPO code	PHYTRA
Regulated status	<p>The pathogen is listed in Annex II of Commission Implementing Regulation (EU) 2019/2072 as <i>Phytophthora ramorum</i> (non-EU isolates) Werres, De Cock & Man in 't Veld [PHYTRA]. The EU isolates of <i>P. ramorum</i> are listed as regulated non quarantine pest (RNQP).</p> <p>The pathogen is included in the EPPO A2 list (EPPO, online_a).</p> <p><i>Phytophthora ramorum</i> is listed as a quarantine pest in the UK (EPPO, online_b).</p>
Pest status in the UK	<p><i>Phytophthora ramorum</i> is present in the UK (Brown and Brasier, 2007; Dossier Section 2; CABI, online; EPPO, online_c).</p> <p>According to the Dossier Section 2, non-EU isolates of <i>P. ramorum</i> are present in the UK: not widely distributed and under official control. It has been found in most regions of the UK, but it is more often reported in wetter, western regions.</p>
Pest status in the EU	<i>Phytophthora ramorum</i> is a regulated pest in the EU.
Host status on Alnus spp.	<p><i>Alnus cordata</i> is reported as a host (EPPO online, O'Hanlon et al., 2016).</p> <p>According to Sansford et al. (2009), <i>A. glutinosa</i> range from low susceptibility to resistant host.</p> <p><i>Alnus glutinosa</i> shows symptoms after inoculation in the lab (Denman et al., 2005; Matsiakh and Menkis, 2023).</p>
Risk Assessment information	<p>Pest Risk Assessments available:</p> <ul style="list-style-type: none"> – Risk analysis for <i>Phytophthora ramorum</i> Werres, de Cock & Man in't Veld, causal agent of sudden oak death, ramorum leaf blight and ramorum dieback (Cave et al., 2008). Risk analysis of <i>Phytophthora ramorum</i>, a newly recognised pathogen threat to Europe and the cause of sudden oak death in the USA (Sansford et al., 2009). – Scientific opinion on the pest risk analysis on <i>Phytophthora ramorum</i> prepared by the FP6 project RAPRA (EFSA Panel on Plant Health, 2011); – Pest risk management for <i>Phytophthora kernoviae</i> and <i>Phytophthora ramorum</i> (EPPO, 2013). – UK Risk Register Details for <i>Phytophthora ramorum</i> (DEFRA, online). – Commodity risk assessment of <i>Acer campestre</i> from the UK (EFSA PLH Panel, 2023a). – Commodity risk assessment of <i>Acer platanoides</i> from the UK (EFSA PLH Panel, 2023b). – Commodity risk assessment of <i>Acer pseudoplatanus</i> from the UK (EFSA PLH Panel, 2023c). – Commodity risk assessment of <i>Cornus</i> spp. from the UK (EFSA PLH Panel, 2024a). – Commodity risk assessment of <i>Sorbus aucuparia</i> from the UK (EFSA PLH Panel, 2024b). – Commodity risk assessment of <i>Betula</i> spp. from the UK (EFSA PLH Panel, 2024c). – Commodity risk assessment of <i>Alnus cordata</i>, <i>A. glutinosa</i> and <i>A. incana</i> from the UK (EFSA PLH Panel, 2025).
Other relevant information for the assessment	
Biology	<p><i>Phytophthora ramorum</i> is most probably native to East Asia (Poimala and Lilja, 2013; Jung et al., 2021). The pathogen is present in Asia (Japan, Vietnam), Europe (Belgium, Croatia, Denmark, Finland, France, Germany, Guernsey, Ireland, Luxembourg, the Netherlands, Norway, Poland, Portugal, Slovenia, the UK), North America (Canada, US) and South America (Argentina) (EPPO, online_c).</p> <p><i>Phytophthora ramorum</i> is heterothallic oomycete species belonging to clade 8c (Blair et al., 2008) with two mating types: A1 and A2 (Boutet et al., 2010).</p> <p><i>Phytophthora</i> species generally reproduce through (a) dormant (resting) spores which can be either sexual (oospores) or asexual (chlamydospores); and (b) fruiting structures (sporangia) which contain zoospores (Erwin and Ribeiro, 1996).</p> <p><i>Phytophthora ramorum</i> produces sporangia on the surfaces of infected leaves and twigs of host plants. These sporangia can be splash-dispersed or carried by wind and rain to longer distances. The sporangia germinate in free water to produce zoospores that penetrate and initiate an infection on new hosts. In infected plant material, the chlamydospores are produced and can serve as resting structures (Davidson et al., 2005; Grünwald et al., 2008). Trunk cankers (e.g. on <i>Quercus</i>, <i>Fagus</i>) are not known to support sporulation (DEFRA, 2008). The pathogen is also able to survive in soil (Shishkoff, 2007). In the west of Scotland, it persisted in soil for at least 2 years after its hosts were removed (Elliot et al., 2013). Oospores were only observed in pairing tests under controlled laboratory conditions (Brasier and Kirk, 2004). Optimal temperatures under laboratory conditions were 16–26°C for growth, 14–26°C for chlamydospore production and 16–22°C for sporangia production (Englander et al., 2006).</p> <p><i>P. ramorum</i> is mainly a foliar pathogen; however, it was also reported to infect shoots, stems and occasionally roots of various host plants (Grünwald et al., 2008; Parke and Lewis, 2007). According to Brown and Brasier (2007), <i>P. ramorum</i> commonly occupies xylem beneath phloem lesions and may spread within xylem and possibly recolonise the phloem from the xylem. <i>P. ramorum</i> can remain viable within xylem for two or more years after the overlying phloem has been excised.</p>

(Continues)

(Continued)

	<p><i>Phytophthora ramorum</i> can disperse by aerial dissemination, water, movement of infected plant material and soil containing propagules on footwear, tires of trucks and mountain bikes or the feet of animals (Davidson et al., 2002; Brasier, 2008).</p> <p>Infected foliar hosts can be a major source of inoculum, which can lead to secondary infections on nearby host plants. Important foliar hosts in Europe are <i>Rhododendron</i> spp. and <i>Larix kaempferi</i> (Brasier and Webber, 2010, Grünwald et al., 2008).</p> <p>Possible pathways of entry for <i>P. ramorum</i> are plants for planting (excluding seed and fruit) of known susceptible hosts; plants for planting (excluding seed and fruit) of non-host plant species accompanied by contaminated attached growing media; soil/growing medium (with organic matter) as a commodity; soil as a contaminant; foliage or cut branches; susceptible (isolated) bark and susceptible wood (EFSA PLH Panel, 2011).</p>
Symptoms	<p>Main type of symptoms</p> <p>There is no information on the symptoms caused by <i>P. ramorum</i> to <i>Alnus</i> spp. plants. <i>P. ramorum</i> causes different types of symptoms depending on the host species and the plant tissue infected.</p> <p>According to DEFRA (2008), <i>P. ramorum</i> causes three different types of disease:</p> <ol style="list-style-type: none"> 'Ramorum bleeding canker' – cankers on trunks of trees, which emit a dark ooze. As they increase in size they can lead to tree death. 'Ramorum leaf blight' – infection of the foliage, leading to discoloured lesions on the leaves. 'Ramorum dieback' – shoot and bud infections which result in wilting, discolouration and dying back of affected parts. <p>Presence of asymptomatic plants</p> <p>If roots are infected by <i>P. ramorum</i>, the plants can be without aboveground symptoms for months until developmental or environmental factors trigger disease expression (Roubtsova and Bostock, 2009; Thompson et al., 2021).</p> <p>Application of some fungicides may reduce symptoms and therefore mask infection, making it more difficult to determine whether the plant is pathogen-free (DEFRA, 2008).</p> <p>Confusion with other pests</p> <p>Various symptoms caused by <i>P. ramorum</i> can be confused with other pathogens, such as: canker and foliar symptoms caused by other <i>Phytophthora</i> species (<i>P. cinnamomi</i>, <i>P. cambivora</i>, <i>P. citricola</i> and <i>P. cactorum</i>); leaf lesions caused by rust in early stages; leafspots caused by sunburn; dieback of twigs and leaves caused by <i>Botryosphaeria dothidea</i> (Davidson et al., 2003).</p> <p><i>Phytophthora ramorum</i> can be easily distinguished from other <i>Phytophthora</i> species based on morphology and molecular tests EPPO (2006).</p>
Host plant range	<p><i>Phytophthora ramorum</i> has a very wide host range, which is expanding.</p> <p>Main host plants include <i>Camellia</i> spp., <i>Larix decidua</i>, <i>L. kaempferi</i>, <i>Pieris</i> spp., <i>Rhododendron</i> spp., <i>Syringa vulgaris</i>, <i>Viburnum</i> spp. and the North American trees species, <i>Lithocarpus densiflorus</i> and <i>Quercus agrifolia</i> (EPPO online_d).</p> <p>Further proven hosts confirmed by Koch's postulates are <i>Abies grandis</i>, <i>A. magnifica</i>, <i>Acer circinatum</i>, <i>A. macrophyllum</i>, <i>A. pseudoplatanus</i>, <i>Adiantum aleuticum</i>, <i>A. jordanii</i>, <i>Aesculus californica</i>, <i>A. hippocastanum</i>, <i>Arbutus menziesii</i>, <i>Arbutus unedo</i>, <i>Arctostaphylos columbiana</i>, <i>Agrostis glauca</i>, <i>A. hooveri</i>, <i>A. manzanita</i>, <i>A. montereyensis</i>, <i>A. morroensis</i>, <i>A. pilosula</i>, <i>A. pumila</i>, <i>A. silvicola</i>, <i>A. viridissima</i>, <i>Calluna vulgaris</i>, <i>Castanea sativa</i>, <i>Ceanothus thyrsiflorus</i>, <i>Chamaecyparis lawsoniana</i>, <i>Chryssolepis chrysophylla</i>, <i>Cinnamomum camphora</i>, <i>Cornus kousa</i>, <i>Cornus hybrids</i>, <i>Corylus cornuta</i>, <i>Fagus sylvatica</i>, <i>Frangula californica</i>, <i>Frangula purshiana</i>, <i>Fraxinus excelsior</i>, <i>Gaultheria procumbens</i>, <i>G. shallon</i>, <i>Griselinia littoralis</i>, <i>Hamamelis virginiana</i>, <i>Heteromeles arbutifolia</i>, <i>Kalmia</i> spp., <i>Larix x eurolepis</i>, <i>Laurus nobilis</i>, <i>Lonicera hispida</i>, <i>Lophostemon confertus</i>, <i>Loropetalum chinense</i>, <i>Magnolia x loebneri</i>, <i>M. oltsopa</i>, <i>M. stellata</i>, <i>Mahonia aquifolium</i>, <i>Maianthemum canadense</i>, <i>Parrotia persica</i>, <i>Photinia fraseri</i>, <i>Phoradendron serotinum</i> subsp. <i>macrophyllum</i>, <i>Photinia x fraseri</i>, <i>Prunus laurocerasus</i>, <i>Pseudotsuga menziesii</i> var. <i>menziesii</i>, <i>Quercus cerris</i>, <i>Q. chryssolepis</i>, <i>Q. falcata</i>, <i>Q. ilex</i>, <i>Q. kelloggii</i>, <i>Q. parvula</i> var. <i>shrevei</i>, <i>Rosa gymnocarpa</i>, <i>Salix caprea</i>, <i>Sequoia sempervirens</i>, <i>Taxus baccata</i>, <i>Trientalis latifolia</i>, <i>Umbellularia californica</i>, <i>Vaccinium myrtillus</i>, <i>V. ovatum</i>, <i>V. parvifolium</i> and <i>Vinca minor</i> (APHIS USDA, 2022; Cave et al., 2008; EPPO, online d; Jung et al., 2016).</p>
Reported evidence of impact	Not relevant, <i>P. ramorum</i> is an EU-regulated quarantine pest.
Evidence that the commodity is a pathway	Life stages of <i>P. ramorum</i> can be present on leaves, stems, branches or roots of whips, bare-root plants and potted plants. <i>P. ramorum</i> can be present in soil; however, potted plants contain only new growing media. <i>P. ramorum</i> is regularly intercepted in the EU on different plant species intended for planting (EUROPHYT/TRACE-NT, online). Therefore, plants for planting of <i>Alnus</i> spp. are possible pathway for <i>P. ramorum</i> .
Surveillance information	<p>The UK carries out surveys for <i>P. ramorum</i> (Dossier Section 1). At growing sites, <i>P. ramorum</i>-infested plants are destroyed, and potentially infested plants are 'held' (prohibited from moving). The UK has a containment policy in the wider environment with official action taken to remove infected trees (Dossier Section 1).</p> <p>As part of an annual survey at ornamental retail and production sites (frequency of visits determined by a decision matrix), <i>P. ramorum</i> is inspected on common host plants. An additional inspection, during the growing period, is carried out at plant passport production sites. Inspections are carried out at a survey to 300 non-woodland wider environment sites annually (Dossier Sections 1).</p>

A.2.2 | Possibility of pest presence in the nursery

A.2.2.1 | Possibility of entry from the surrounding environment

Phytophthora ramorum is present in the UK, it has been found in most regions of the UK, but it is more often reported in wetter, western regions (Dossier Section 1). The possible entry of *P. ramorum* from surrounding environment to the nurseries may occur through aerial dissemination, water and animals (Davidson et al., 2002).

The surrounding land would tend to be arable farmland with some pasture for animals and small areas of woodland. Hedges are often used to define field boundaries and grown along roadsides. These hedges contain host plants such as *Acer* and *Alnus*.

Uncertainties

- The dispersal range of *P. ramorum* sporangia.
- There is no information available on the distance of the nurseries to sources of pathogen in the surrounding environment.

A.2.2.2 | Possibility of entry with new plants/seeds

Plants are produced by seeds and grafting. Seeds purchased in the UK are certified under The Forest Reproductive Material (Great Britain) Regulations 2002 ([legislation.gov.uk](https://www.legislation.gov.uk)); seedlings sourced in the UK are certified with UK Plant Passports; a small percentage of plants may be obtained from EU (Netherlands); seedlings from the EU countries are certified with phytosanitary certificates.

The pathogen is not known to be seedborne or seed transmitted, therefore not expected to enter the nursery via the seed pathway. Grafted plants (scion and buds in the case of grafting) originate from the nursery itself.

The seedling pathways is unlikely because the conditions of their production (seedbed in the greenhouse, pest-free growing media) are expected to prevent the infection of seedlings.

The nurseries use virgin peat or peat-free compost (a mixture of coir, tree bark, wood fibre, etc.) as a growing media (Dossier Section 1).

The growing media is heat-treated by commercial suppliers during production to eliminate pests and diseases (Dossier Section 1).

Uncertainties

- None.

Taking into consideration the above evidence and uncertainties, the Panel considers that it is unlikely for the pathogen to enter the nurseries via seeds or seedlings of *Alnus*.

A.2.2.3 | Possibility of spread within the nursery

Host plants of *P. ramorum* are grown outdoors, in the open air, or field, in the nurseries producing *Alnus* spp. plants (Dossier Section 1). Therefore, is it possible that *P. ramorum* can spread within the nursery from infested host plants of other species to the plot with *Alnus* spp. plants.

Phytophthora ramorum can spread within the nurseries by aerial dissemination/water splash: via soil, water, movement of infested plant material (e.g. infested leaves) and animals/humans (Davidson et al., 2002). However, due to the quarantine status of *P. ramorum*, the presence of symptomatic plants is expected to be limited in the nurseries.

Uncertainties

- If asymptomatic plants are detected and efficiently removed.

Taking into consideration the above evidence and uncertainties, the Panel considers that the spread of the pathogen within the nurseries is possible by air currents.

A.2.3 | Information from interceptions

Phytophthora ramorum is regularly intercepted in the EU on different plant species intended for planting (EUROPHYT/TRADES-NT, online). In the EUROPHYT/TRADES-NT database, there are no records of notification of *Alnus* plants for planting neither from the UK nor from other countries due to the presence of *P. ramorum* between the years 1995 and January 2025 (EUROPHYT/TRADES-NT, online).

A.2.4 | Evaluation of the risk mitigation measures

In the table below, all risk mitigation measures currently applied in the UK are listed and an indication of their effectiveness on *P. ramorum* is provided. The description of the implemented risk mitigation measures is provided in [Table 8](#).

No.	Risk mitigation measure	Effect on the pest	Evaluation and uncertainties
1	Registration of production sites	Yes	<p>All nurseries are registered as professional operator with the UK NPPO, by the Animal Plant Health Agency (APHA) for England and Wales, or with SASA for Scotland, and are authorised to issue UK plant passports (Dossier Section 1).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – The risk mitigation measure is expected to be effective in reducing the likelihood of presence of the pathogen on the commodity. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – Whether early symptoms on <i>Alnus</i> spp. are easily recognisable.
2	Certification of plant material	Yes	<p><i>Alnus cordata</i> and <i>A. glutinosa</i> seeds purchased in the UK are certified under The Forest Reproductive Material (Great Britain) Regulations 2002 (legislation.gov.uk); seedlings sourced in the UK are certified with UK Plant Passports; a small percentage of plants may be obtained from EU (Netherlands); seedlings from the EU countries are certified with phytosanitary certificates. (Dossier Section 1).</p> <p>The starting material of <i>Alnus</i> production consists of seed and seedlings. Seeds are certified. Seedlings for production sourced in the UK are certified with UK Plant Passports; seedlings from the EU countries are certified with phytosanitary certificates.</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – <i>Phytophthora ramorum</i> is a quarantine pest and it is highly unlikely that the pathogen is present on the certified starting material. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – None.
3	Origin and treatment of growing media	Yes	<p>In the production or procurement of these plants, the use of growing media is assessed for the potential to harbour and transmit plant pests. Growers most commonly use virgin peat or peat-free compost, which is a mixture of coir, tree bark, wood fibre, etc. The compost is heat-treated by commercial suppliers during production to eliminate pests and diseases. It is supplied in sealed bulk bags or shrink-wrapped bales and stored off the ground on pallets, these are completely hygienic and free from contamination. Where delivered in bulk, compost is kept in a dedicated bunker, either indoors, or covered by tarpaulin outdoors, and with no risk of contamination with soil or other material (Dossier Section 1).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – The measure is efficient in preventing the entry of the pathogen via the substrate into the nursery. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – None.
4	Surveillance, monitoring and sampling	Yes	<p>Inspection is carried out at least once a year as part of the Quarantine Surveillance programme (Great Britain uses the same framework for its surveillance programme as the EU). Surveillance is based on visual inspection with samples taken from symptomatic material, and where appropriate, samples are also taken from asymptomatic material (e.g. plants, tubers, soil, watercourses) (Dossier Section 1).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – The surveillance, monitoring and sampling can detect the pathogen. No results are reported. – Due to the canopy size of older trees (up to 25 years old), the pathogen could remain undetected. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – The efficiency of the surveillance, monitoring and sampling.
5	Hygiene measures	Yes	<p>According to the Dossier Section 1, all the nurseries have plant hygiene and housekeeping rules and practices in place, which are communicated to all relevant employees. These measures include:</p> <ul style="list-style-type: none"> – Cleaning and sterilisation of tools. – Waste treatment and disposal. <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – It is unlikely that the fungus spreads by the pruning tools. – The correct disposal of infected plant material prevents the spread of the fungus. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – The efficiency of hygiene measures performed in the nurseries.

(Continued)

No.	Risk mitigation measure	Effect on the pest	Evaluation and uncertainties
6	Irrigation water quality and/or treatments	Yes	<p>Growers are required to assess water sources, irrigation and drainage systems used in the plant production for the potential to harbour and transmit plant pests. Water is routinely sampled and sent for analysis. No quarantine pests have been found (Dossier Section 1).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – There is no disinfestation treatment applied to the irrigation water. However, irrigation water is routinely sampled and tested for quarantine pests. This procedure can reduce the risk. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – The frequency of sampling and the method used for the detection of the pathogen.
7	Application of pest control products	Yes	<p>Crop protection is achieved using a combination of measures including approved plant protection products, biological control or physical measures. Plant protection products are only used when necessary and records of all plant protection treatments are kept (Dossier Section 1).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – Some plant protection products are applied and could reduce the likelihood of the infection by the pathogen, but detailed information is lacking in the Dossier. – The efficacy of plant protection products on large trees (up to 25 years old) could be reduced. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – No specific information on the efficacy of the plant protection products used.
8	Washing of the roots	Yes	<p>Specimen trees (up to 25 years old) are grown in soil for up to 9 years, root washed and placed in pots (based on the request letter of DEFRA to the EU Commission on September 25th, 2023).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – The washing of the roots removes (parts of) the soil and the pathogen present in the soil. However, the root plant system of plants of 9 years may be difficult to clean completely from soil. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – The effectiveness of the washing to remove all soil with the pathogen.
9	Inspections and management of plants before export	Yes	<p>The UK NPPO carries out inspections and testing where required by the country of destination's plant health legislation, to ensure all requirements are fulfilled and a valid phytosanitary certificate with the correct additional declarations is issued.</p> <p>Separate to any official inspection, plant material is checked by growers for plant health issues before dispatch.</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> – The inspections and management of plants before export can detect the pathogen. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – Whether early symptoms caused by the pathogen on <i>Alnus</i> species are identified by visual inspections.

A.2.5 | Overall likelihood of pest freedom for the exported commodity (comparative)

A.2.5.1 | Comparison with other relevant commodity Risk Assessments involving *Phytophthora ramorum*

Phytophthora ramorum was already assessed as actionable pest for graftwood, bare-root plants, cell-grown plants and rooted plants in pots (of up to 7 years old) of *A. cordata*, *A. glutinosa* and *A. incana* (EFSA PLH Panel, 2025). *Alnus* commodity types evaluated originate from nurseries sharing the similar production system. Therefore, for the estimation of the pest freedom level of *P. ramorum* in exported specimen trees (from 7 to 25 years old) of *Alnus* plants, the Panel decided to base the estimated values on the ones elicited for the other commodities of *A. cordata*, *A. glutinosa* and *A. incana* (EFSA PLH Panel, 2025). In addition, because of the similarity of the commodities, the production systems, the locations of the nurseries, the Panel also considered the EKE values from the previous Scientific opinion on *Betula pendula* and *B. pubescens* from the UK (EFSA PLH Panel, 2024).

A.2.6 | Overall likelihood of pest freedom for specimen trees (from 7 to 25 years old)

A.2.6.1 | Reasoning for a scenario which would lead to a reasonably low number of infected specimen trees (from 7 to 25 years old)

The scenario assumes a low pressure of the pathogen in the nurseries and in the surroundings. Plants are exported as dormant plants without leaves. The scenario assumes *Alnus* spp. to be minor hosts for the pathogen. The scenario also assumes that symptoms of the disease are visible and promptly detected during inspections. Furthermore, due to the extended period during which specimen trees are kept in pots in the nursery (up to 25 years old) soil-borne infections originating from the bare soil cultivation are more likely to express symptoms of the disease, therefore be detected during repeated inspections.

A.2.6.2 | Reasoning for a scenario which would lead to a reasonably high number of infected specimen trees (from 7 to 25 years old)

The scenario assumes a high pressure of the pathogen in the surrounding environment of the nurseries because suitable hosts are present. The scenario assumes that the pathogen can infect leaves, which may still be present on the plants at the time of export. The scenario also assumes that the pathogen is not detected during the inspections because of the presence of asymptomatic plants or difficulties in recognising early symptoms. Grafting can increase the incidence of the pathogen (via infected buds or by woundings).

In comparison with the other commodities (graftwood, bare-root plants, cell-grown plants and rooted plants of up to 7 years old in pots), the canopy size of large specimen trees (up to 25 years old) is larger, increasing the difficulties in detecting the pathogen, particularly an early infection. Furthermore, the exposure time is longer, increasing the probability for the plants to be infected.

A.2.6.3 | Reasoning for a central scenario equally likely to over- or underestimate the number of infected specimen trees (from 7 to 25 years old) (Median)

In comparison to potted plants up to 7 years old, the specimen trees are generally larger and older up to 25 years old, increasing the time of exposure to the pathogen. The scenario assumes a limited presence of the pathogen in the nurseries and in the surroundings, and a limited reported susceptibility of *Alnus* spp. The pathogen is a regulated quarantine pest in the UK and under official control.

A.2.6.4 | Reasoning for the precision of the judgement describing the remaining uncertainties (1st and 3rd quartile/interquartile range)

The limited information on the occurrence of the pathogen in the nurseries and the surroundings and the susceptibility of *Alnus* spp. results in high level of uncertainties.

A.2.7 | Elicitation outcomes of the assessment of the pest freedom for *Phytophthora ramorum* specimen trees (from 7 to 25 years old)

The elicited and fitted values for *P. ramorum* for pest infestation and pest freedom agreed by the Panel are shown in Tables A.3 and A.4 and in Figure A.2.

TABLE A.3 Elicited and fitted values of the uncertainty distribution of pest infestation by *Phytophthora ramorum* per 10,000 plants of specimen trees (from 7 to 25 years old).

Percentile	1%	2.5%	5%	10%	17%	25%	33%	50%	67%	75%	83%	90%	95%	97.5%	99%
Elicited values	0					12		25		45					105
EKE	0.639	1.43	2.64	4.95	8.00	11.9	16.0	25.3	37.1	44.7	54.7	66.1	79.8	91.6	105

Note: The EKE results is the *BetaGeneral* (1.1527, 5.328, 0, 175) distribution fitted with @Risk version 7.6.

Based on the numbers of estimated infected plants, the pest freedom was calculated (i.e. = 10,000 – number of infected plants per 10,000). The fitted values of the uncertainty distribution of the pest freedom are shown in Table A.4.

TABLE A.4 The uncertainty distribution of plants free of *Phytophthora ramorum* per 10,000 plants of specimen trees (from 7 to 25 years old) calculated by Table A.3.

Percentile	1%	2.5%	5%	10%	17%	25%	33%	50%	67%	75%	83%	90%	95%	97.5%	99%
Values	9895					9955		9975		9988					10,000
EKE results	9895	9908	9920	9934	9945	9955	9963	9975	9984	9988	9992	9995	9997	9998.6	9999.4

Note: The EKE results are the fitted values.

(A)




FIGURE A.2 (Continued)

(B)




FIGURE A.2 (Continued)




FIGURE A.2 (A) Elicited uncertainty of pest infection per 10,000 plants of specimen trees (histogram in blue—vertical blue line indicates the elicited percentile in the following order: 1%, 25%, 50%, 75%, 99%) and distributional fit (red line); (B) uncertainty of the proportion of pest-free plants per 10,000 (i.e. = 1 – pest infection proportion expressed as percentage); (C) descending uncertainty distribution function of pest infection per 10,000 plants.

A.2.8 | Reference list

APHIS USDA (Animal and Plant Health Inspection Service U.S. Department of Agriculture). (2022). APHIS Lists of Proven Hosts of and Plants Associated with Phytophthora ramorum. September 2022. 12 pp. Available online: https://www.aphis.usda.gov/plant_health/plant_pest_info/pram/downloads/pdf_files/usdaprlist.pdf [Accessed: 27 September 2024].

Blair, J. E., Coffey, M. D., Park, S. Y., Geiser, D. M., & Kang, S. (2008). A multi-locus phylogeny for Phytophthora utilizing markers derived from complete genome sequences. *Fungal Genetics and Biology*, 45(3), 266–277. <https://doi.org/10.1016/j.fgb.2007.10.010>

Boutet, X., Vercauteren, A., Heungens, C., & Kurt, A. (2010). Mating of Phytophthora ramorum: functionality and consequences. In: Frankel, S. J., Kliejunas, J. T., and Palmieri, K. M. (eds.). Proceedings of the Sudden Oak Death Fourth Science Symposium. Albany, CA: US Department of Agriculture, Forest Service, Pacific Southwest Research Station, 229, 97–100.

Brasier, C. (2008). Phytophthora ramorum + P. kernoviae = international biosecurity failure. In: Frankel SJ, Kliejunas JT, Palmieri KM (eds). Proceedings of the sudden oak death third science symposium. USDA Forest Service, Pacific Southwest Research Station, Albany, CA: US Department of Agriculture, 214, 133–139.

Brasier, C., & Kirk, S. (2004). Production of gametangia by Phytophthora ramorum in vitro. *Mycological Research*, 108(7), 823–827. <https://doi.org/10.1017/s0953756204000565>

Brasier, C., & Webber, J. (2010). Sudden larch death. *Nature*, 466, 824–825. <https://doi.org/10.1038/466824a>

Brown, A. V., & Brasier, C. M. (2007). Colonization of tree xylem by Phytophthora ramorum, P. kernoviae and other Phytophthora species. *Plant Pathology*, 56(2), 227–241. <https://doi.org/10.1111/j.1365-3059.2006.01511.x>

CABI (Centre for Agriculture and Bioscience International). (online). Phytophthora ramorum (Sudden Oak Death (SOD)). Available online: <https://www.cabi.org/cpc/datasheet/40991> [Accessed: 27 September 2024].

Cave, G. L., Randall-Schadel, B., & Redlin, S. C. (2008). Risk analysis for Phytophthora ramorum Werres, de Cock & Man in't Veld, causal agent of sudden oak death, ramorum leaf blight, and ramorum dieback. US Department of Agriculture, Animal and Plant Health Inspection Service, Raleigh, NC. 88 pp.

Davidson, J. M., Rizzo, D. M., Garbelotto, M., Tjosvold, S., & Slaughter, G. W. (2002). Phytophthora ramorum and sudden oak death in California: II. Transmission and survival. In: Standiford, R. B., McCreary, D., & Purcell, K. L. (eds.). Proceedings of the fifth symposium on oak woodlands: Oaks in California's challenging landscape. San Diego, California, US Department of Agriculture, Forest Service, Pacific Southwest Research Station: 184, 741–749.

Davidson, J. M., Werres, S., Garbelotto, M., Hansen, E. M., & Rizzo, D. M. (2003). Sudden oak death and associated diseases caused by Phytophthora ramorum. *Plant Health Progress*, 4(1), 12. <https://doi.org/10.1094/php-2003-0707-01-dg>

Davidson, J. M., Wickland, A. C., Patterson, H. A., Falk, K. R., & Rizzo, D. M. (2005). Transmission of Phytophthora ramorum in mixed-evergreen forest in California. *Phytopathology*, 95, 587–596. <https://doi.org/10.1094/phyto-95-0587>

DEFRA (Department for Environment, Food and Rural Affairs). (online). UK Risk Register Details for Phytophthora ramorum. Available online: <https://planthealthportal.defra.gov.uk/pests-and-diseases/uk-plant-health-risk-register/viewPestRisks.cfm?csref=23022> [Accessed: 27 September 2024].

DEFRA (Department for Environment, Food and Rural Affairs). (online). Fera list of natural hosts for Phytophthora ramorum with symptom and location. 2015. <https://planthealthportal.defra.gov.uk/assets/uploads/P-ramorum-host-list-finalupdate-NOV-20-15.pdf>. [Accessed: 15 September 2024].

Denman, S., Kirk, S. A., Brasier, C. M., & Webber, J. F. (2005). In vitro leaf inoculation studies as an indication of tree foliage susceptibility to Phytophthora ramorum in the UK. *Plant Pathology*, 54(4), 512–521. <https://doi.org/10.1111/j.1365-3059.2005.01243.x>

DiLeo, M. V., Bienapfl, J. C., & Rizzo, D. M. (2008). Phytophthora ramorum infects hazelnut, vine maple, blue blossom, and manzanita species in California. *Plant Health Progress*, <https://doi.org/10.1094/php-2008-0118-02-br>

EFSA PLH Panel (EFSA Panel on Plant Health). (2011). Scientific Opinion on the Pest Risk Analysis on Phytophthora ramorum prepared by the FP6 project RAPRA. *EFSA Journal*, 9(6), 2186. <https://doi.org/10.2903/j.efsa.2011.2186>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Potting, R., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent Civera, A., Yuen, J., Zappalà, L., Battisti, A., Mas, H., Rigling, D., Faccoli, M., Gardi, C., Iacopetti, G., Mikulová, A., Mosbach-Schulz, O., Stergulc, F., Streissl, F., & Gonthier, P. (2023a). Scientific Opinion on the commodity risk assessment of Acer campestre plants from the UK. *EFSA Journal*, 21(7), 8071. <https://doi.org/10.2903/j.efsa.2023.8071>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Potting, R., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent Civera, A., Yuen, J., Zappalà, L., Battisti, A., Mas, H., Rigling, D., Faccoli, M., Gardi, C., Iacopetti, G., Mikulová, A., Mosbach-Schulz, O., Stergulc, F., Streissl, F., & Gonthier, P. (2023b). Scientific Opinion on the commodity risk assessment of Acer platanoides plants from the UK. *EFSA Journal*, 21(7), 8073. <https://doi.org/10.2903/j.efsa.2023.8073>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Potting, R., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent Civera, A., Yuen, J., ... Gonthier, P. (2023c). Commodity risk assessment of Acer pseudoplatanus plants from the UK. *EFSA Journal*, 21(7), 8074. <https://doi.org/10.2903/j.efsa.2023.8074>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Potting, R., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent Civera, A., Yuen, J., ... Gonthier, P. (2023d). Commodity risk assessment of Fagus sylvatica plants from the UK. *EFSA Journal*, 21(7), 8118. <https://doi.org/10.2903/j.efsa.2023.8118>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Potting, R., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent Civera, A., Yuen, J., ... Potting, R. (2023e). Commodity risk assessment of Quercus robur plants from the UK. *EFSA Journal*, 21(10), 8314. <https://doi.org/10.2903/j.efsa.2023.8314>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Gonthier, P., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent Civera, A., Yuen, J., ... Potting, R. (2024a). Commodity risk assessment of Betula pendula and Betula pubescens plants from the UK. *EFSA Journal*, 22(11), 9051. <https://doi.org/10.2903/j.efsa.2024.9051>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Gonthier, P., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent Civera, A., Yuen, J., ... Potting, R. (2024b). Commodity risk assessment of Cornus alba and Cornus sanguinea plants from the UK. *EFSA Journal*, 22(3), 8657. <https://doi.org/10.2903/j.efsa.2024.8657>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Gonthier, P., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent Civera, A., Yuen, J., ... Potting, R. (2024c). Commodity risk assessment of Sorbus aucuparia plants from the UK. *EFSA Journal*, 22(6), 8837. <https://doi.org/10.2903/j.efsa.2024.8837>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Gonthier, P., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent Civera, A., Yuen, J., ... Potting, R. 2024d Commodity risk assessment of *Corylus avellana* plants from the UK. *EFSA Journal*, 22(1), 8495. <https://doi.org/10.2903/j.efsa.2024.8495>

EFSA PLH Panel (EFSA Panel on Plant Health), Bragard, C., Baptista, P., Chatzivassiliou, E., Di Serio, F., Gonthier, P., Jaques Miret, J. A., Justesen, A. F., MacLeod, A., Magnusson, C. S., Milonas, P., Navas-Cortes, J. A., Parnell, S., Reignault, P. L., Stefani, E., Thulke, H.-H., Van der Werf, W., Vicent Civera, A., Yuen, J., ... Potting, R. (2025). Commodity risk assessment of *Alnus cordata*, *Alnus glutinosa* and *Alnus incana* plants from the UK. *EFSA Journal*, 23(1), 9189. <https://doi.org/10.2903/j.efsa.2025.9189>

Elliott, M., Meagher, T. R., Harris, C., Searle, K., Purse, B. V., & Schlenzig, A. (2013). The epidemiology of *Phytophthora ramorum* and *P. kernoviae* at two historic gardens in Scotland. In Frankel, S. J., Kliejunas, J. T., Palmieri, K. M., and Alexander, J. M. (eds.). Sudden oak death fifth science symposium. Albany, CA, USA: US Department of Agriculture, Forest Service, Pacific Southwest Research Station, 23–32.

Englander, L., Browning, M., and Tooley, P. W. (2006). Growth and sporulation of *Phytophthora ramorum* in vitro in response to temperature and light. *Mycologia*, 98(3), 365–373. <https://doi.org/10.3852/mycologia.98.3.365>

EPPO (European and Mediterranean Plant Protection Organization). (2013). Pest risk management for *Phytophthora kernoviae* and *Phytophthora ramorum*. EPPO, Paris. Available online: http://www.eppo.int/QUARANTINE/Pest_Risk_Analysis/PRA_intro.htm [Accessed: 27 September 2023]

EPPO (European and Mediterranean Plant Protection Organization). (online_a). EPPO A2 List of pests recommended for regulation as quarantine pests, version 2021-09. Available online: https://www.eppo.int/ACTIVITIES/plant_quarantine/A2_list [Accessed: 27 September 2023].

EPPO (European and Mediterranean Plant Protection Organization). (online_b). *Phytophthora ramorum* (PHYTRA), Categorization. Available online: <https://gd.eppo.int/taxon/PHYTRA/categorization> [Accessed: 27 September 2024]

EPPO (European and Mediterranean Plant Protection Organization). (online_c). *Phytophthora ramorum* (PHYTRA), Distribution. Available online: <https://gd.eppo.int/taxon/PHYTRA/distribution> [Accessed: 27 September 2024].

EPPO (European and Mediterranean Plant Protection Organization). (online_d). *Phytophthora ramorum* (PHYTRA), Host plants. Available online: <https://gd.eppo.int/taxon/PHYTRA/hosts> [Accessed: 27 September 2023]. EPPO (European and Mediterranean Plant Protection Organization). (online_e). *Phytophthora ramorum* (PHYTRA), Photos. Available online: <https://gd.eppo.int/taxon/PHYTRA/photos> [Accessed: 27 September 2024].

EPPO (European and Mediterranean Plant Protection Organization). (2006). EPPO Standards PM 7/66 Diagnostics. *Phytophthora ramorum*. *EPPO Bulletin*, 36, 145–155.

Erwin, D. C., & Ribeiro, O. K. (1996). *Phytophthora* diseases worldwide. St. Paul, Minnesota: APS Press, American Phytopathological Society, 562 pp.

EUROPHYT. (online). European Union Notification System for Plant Health Interceptions - EUROPHYT Available online: http://ec.europa.eu/food/plant/plant_health_biosecurity/europhyt/index_en.htm [Accessed: 21/1/2025].

Jung, T., Jung, M. H., Webber, J. F., Kageyama, K., Hieno, A., Masuya, H., Uematsu, S., Pérez-Sierra, A., Harris, A. R., Forster, J., Rees, H., Scanu, B., Patra, S., Kudláček, T., Janoušek, J., Corcobado, T., Milenović, I., Nagy, Z., Csorba, I., Bakonyi, J., & Brasier, C. M. (2021). The destructive tree pathogen *Phytophthora ramorum* originates from the laurosilva forests of East Asia. *Journal of Fungi*, 7(3), 226. <https://doi.org/10.3390/jof7030226>

Jung, T., Orlikowski, L., Henricot, B., Abad-Campos, P., Aday, A. G., Agúin Casal, O., ... Peréz-Sierra, A. (2016). Widespread *Phytophthora* infestations in European nurseries put forest, semi-natural and horticultural ecosystems at high risk of *Phytophthora* diseases. *Forest Pathology*, 46(2), 134–163.

Matsiakh, I., & Menkis, A. (2023). An overview of *Phytophthora* species on woody plants in Sweden and other Nordic Countries. *Microorganisms*, 11(5), 1309.

O'Hanlon, R., Choiseul, J., Corrigan, M., Catarame, T., & Destefanis, M. (2016). Diversity and detections of *Phytophthora* species from trade and non-trade environments in Ireland. *EPPO Bulletin*, 46(3), 594–602. <https://doi.org/10.1111/epb.12331>

Parke, J. L., & Lewis, C. (2007). Root and stem infection of *Rhododendron* from potting medium infested with *Phytophthora ramorum*. *Plant Disease*, 91, 1265–1270. <https://doi.org/10.1094/pdis-91-10-1265>

Poimala, A., & Lilja, A. (2013). NOBANIS – Invasive Alien Species Fact Sheet – *Phytophthora ramorum*. From: Online Database of the European Network on Invasive Alien Species. 14 pp. Available online: <https://www.nobanis.org/globalassets/speciesinfo/p/phytophthora-ramorum/phytophthora-ramorum.pdf> [Accessed: 27 September 2023].

Roubtsova, T. V., & Bostock, R. M. (2009). Episodic abiotic stress as a potential contributing factor to onset and severity of disease caused by *Phytophthora ramorum* in *Rhododendron* and *Viburnum*. *Plant Disease*, 93(9), 912–918. <https://doi.org/10.1094/pdis-93-9-0912>

Sanford, C. E., Inman, A. J., Baker, R., Brasier, C., Frankel, S., de Gruyter, J., Husson, C., Kehlenbeck, H., Kessel, G., Moralejo, E., Steeghs, M., Webber, J., & Werres, S. (2009). Report on the risk of entry, establishment, spread and socio-economic loss and environmental impact and the appropriate level of management for *Phytophthora ramorum* for the EU. Deliverable Report 28. EU Sixth Framework Project RAPRA. 310 pp.

Shishkoff, N. (2007). Persistence of *Phytophthora ramorum* in soil mix and roots of nursery ornamentals. *Plant Disease*, 91(10), 1245–1249. <https://doi.org/10.1094/pdis-91-10-1245>

TRACES-NT. (online). TRAde Control and Expert System. Available online: <https://webgate.ec.europa.eu/tracesnt> [Accessed: 21/1/2025].

A.3 | PHYTOPHTHORA SISKIYOUENSIS

A.3.1 | Organism information

Taxonomic information	Current valid scientific name: <i>Phytophthora siskiyouensis</i> Reeser & E.M. Hansen Synonyms: - Name used in the EU legislation: - Order: Peronosporales Family: Peronosporaceae Common name: No common name in EPPO GD Name used in the Dossier: <i>Phytophthora siskiyouensis</i>
Group	Oomycetes
EPPO code	PHYTSK
Regulated status	The pest is not regulated in the EU territory and not listed in Commission Implementing Regulation (EU) 2019/2072.
Pest status in UK	Present [unknown distribution, (CABI; USDA, online) one reported outbreak in Dorset (Dossier Section 5.2)].
Pest status in the EU	The pathogen is not Present in EU (CABI; USDA, online).

(Continued)

Host status on <i>Alnus</i> spp.	<p><i>Phytophthora siskiyouensis</i> has been reported in <i>Alnus incana</i> (grey alder) (Perez-Sierra et al., 2015), <i>A. glutinosa</i> (European alder) (Smith et al., 2006), <i>A. cordata</i> (Italian alder) and <i>A. rhombifolia</i> (white alder) (Rooney-Latham et al., 2009), <i>A. rubra</i> (red alder) (Sims et al., 2015a; 2015b; Navarro et al., 2015; CABI; USDA, online). Pathogenicity tests in the UK showed <i>A. cordata</i> to be the most susceptible host species followed by <i>A. glutinosa</i> and <i>A. incana</i> (Perez-Sierra et al., 2014).</p>
Risk Assessment information	<ul style="list-style-type: none"> - Pest Risk Assessments are not available. - Commodity risk assessment of <i>Alnus cordata</i>, <i>A. glutinosa</i> and <i>A. incana</i> from the UK (EFSA PLH Panel, 2025).
Other relevant information for the assessment	
Biology	<p><i>Phytophthora siskiyouensis</i> is a relatively new species (Reeser et al., 2007). Little is known about its biology/life cycle, but is mainly found in soil and water streams. <i>Phytophthora siskiyouensis</i> (Reeser et al., and 2007) was discovered first in streams and soil in Curry County, Oregon in areas dominated by native forest. Later the pathogen was reported to cause the death of alders planted in cities in California (Rooney-Latham et al. 2009) and Australia (Smith et al. 2006).</p> <p><i>Phytophthora siskiyouensis</i> produces several types of structures specialised for survival, dispersal, or infection (Forest Phytophtoras of the World b, online).</p> <p><i>Phytophthora siskiyouensis</i> can be distinguished from other <i>Phytophthora</i> species by the shape of the terminal and intercalary sporangia, and the shape of the antheridia. With its semi-papillate sporangia and predominately paragynous antheridia, <i>P. Phytophthora siskiyouensis</i> would be placed in Waterhouse's (1963) Group III.</p> <p><u>Asexual phase:</u></p> <p>Sporangia are semi-papillate, some with two and more rarely with three papilla; persistent or caducous with variable pedicel length (2–66 µm L); ovoid, ellipsoid, reniform (or kidney), distorted shapes (26–95 L x 23–50 W µm); hyphal swellings small, often associated with sporangiophore; sporangia originated terminal but often subterminal and occasionally intercalary in unbranched or simple sympodial sporangiophores. Hyphal swellings and chlamydospores are absent.</p> <p><u>Sexual phase:</u></p> <p>Homothallic. Oogonia are globose to subglobose (23–42 µm diam), occasionally much elongated or with a funnel shape tapering toward the stalk, frequently sessile and occasionally laterally intercalary; antheridia predominately paragynous, spherical, ellipsoid or ovoid, club-shaped (10–18 L x 8–13 W µm), terminal, occasionally intercalary and usually clivous, attached anywhere on the oogonium; oospores are globose to subglobose (19–33 µm diam) and usually aplerotic. The above combination of characteristics clearly separates <i>P. siskiyouensis</i> from other known <i>Phytophthora</i> species. (Abad e al., 2023a; 2023b; ITP online; Forest Phytophtoras of the World b, online).</p> <p><u>Genetics:</u></p> <p>ITS sequences showed greatest similarity to <i>P. tropicalis</i> and <i>P. capsici</i> in phylogenetic clade 2, but consistent differences were found between <i>P. siskiyouensis</i> and these related species. <i>P. siskiyouensis</i> is distinguished morphologically from these relatives by its homothallic condition, predominately paragynous antheridia and semipapillate sporangia on unbranched sporangiophores (Hansen et al., 2011; Reeser et al., 2007).</p>
Symptoms	<p>Main type of symptoms</p> <p>The predominant symptoms on the diseased <i>Alnus</i> trees are sparse foliage, dieback in the canopy and bleeding cankers on the trunks. Cankers occur primarily at the bases of the trunks near the soil line and extend upward. When the outer bark is removed from the cankers, a cinnamon-brown margin is observed separating cream-coloured healthy tissue from dark orange-brown diseased tissue. The diseased tissue extends through the bark to the vascular cambium and sapwood interface, characteristic of <i>Phytophthora</i> diseases (Rooney-Latham et al., 2009). Symptoms include bleeding stem lesions (CABI, online). The symptoms caused by <i>P. siskiyouensis</i> on various hosts include canker, root rot, blight, dieback, leaf necrosis, twig blight (Forest Phytophtoras of the World, a).</p> <p>Presence of asymptomatic plants</p> <p>No information is available concerning the presence of <i>P. siskiyouensis</i> in symptomless host plants.</p> <p>Confusion with other pests</p> <p><i>Phytophthora siskiyouensis</i> symptoms include bleeding stem lesions similar to those caused by <i>Phytophthora alni</i> (CABI, online). It should be noted that <i>P. alni</i> is widespread both in UK and EU affecting all species of alder (Forest Research, online). <i>Phytophthora siskiyouensis</i> is most readily recognised from other <i>Phytophthora</i> species by its uniquely and irregularly shaped sporangia formed in water and by the frequent occurrence of sessile and intercalary oogonia in agar. Sporangia may also be formed on some agar media with a mostly elongated ovoid shape and cymose sporangiophore reminiscent of <i>P. hibernalis</i>. Sporangia in water are characteristically reniform with off-centre semipapillae and sporangiophore attachment, the irregular sporangia are reminiscent of <i>P. citricola</i>. <i>P. siskiyouensis</i> is distinguished readily from the latter species, however, by having deciduous sporangia with variable length pedicels, variable orientation of semipapilla and sporangiophore attachment, intercalary and sessile attachment of some oogonia and mostly aplerotic oospores. <i>P. siskiyouensis</i> sporangia resemble those depicted for <i>Phytophthora quercina</i>, except that they are slightly larger, semipapillate, weakly deciduous, and are formed singly on simple, unbranched sporangiophores. Sexual structures also resemble <i>P. quercina</i>, except that oogonia may be sessile or intercalary, and antheridia may be paragynous or amphigynous. The oogonial stalk and arrangement of paragynous antheridia are similar to those described for <i>P. hedraiantha</i> (Forest Phytophtoras of the World, online b).</p>

(Continues)

(Continued)

Host plant range	<i>Phytophthora siskiyouensis</i> does not appear to be highly specific to alder species and is known to cause occasional symptoms on a variety of associated plants (Hansen et al., 2011). Apart from the <i>Alnus</i> species the host range of the oomycete includes <i>Notholithocarpus densiflorus</i> (Tanoak) and <i>Umbellularia californica</i> (Reeser et al., 2008). <i>P. siskiyouensis</i> is considered a potential weak pathogen on seedlings of other tree species, including <i>Citrus</i> , <i>Acacia</i> , and <i>Eucalyptus</i> (Smith et al., 2004).
Reported evidence of impact	<i>Phytophthora siskiyouensis</i> is of great concern as a pathogen of Italian alder (<i>A. cordata</i>) and the native white alder (<i>A. rhombifolia</i>) in California (Rooney-Latham et al., 2009), and European alder (<i>A. glutinosa</i> (L.) Gaertn) in Australia (Smith et al., 2006) (Forest Phytophtoras of the World a, online). In the UK Plant Pest Risk Register <i>P. siskiyouensis</i> has an impact rating of 4 out of 5 (UK Plant Health Portal, online).
Evidence that the commodity is a pathway	<i>Phytophthora siskiyouensis</i> is apparently present in the nursery industry as a pathogen on horticultural alder planting stock (Forest Phytophtoras of the World b, online). <i>Alnus</i> spp. are hosts of <i>P. siskiyouensis</i> . Life stages of <i>P. siskiyouensis</i> can be present on leaves, stems, branches or roots of whips, bare root plants and potted plants. <i>P. siskiyouensis</i> can be present in soil, however potted plants contain only new growing media. Therefore, plants for planting of <i>Alnus</i> spp. are possible pathway for <i>P. siskiyouensis</i> .
Surveillance information	No information.

A.3.2 | Possibility of pest presence in the nursery

A.3.2.1 | Possibility of entry from the surrounding environment

There is a high uncertainty on the prevalence and distribution of this pathogen in the UK (unknown distribution reported in UK Pest Risk Register). There is only one report of the presence of the pathogen in the southwest of England (Perez-Sierra et al., 2015). There is a possibility that the pathogen is present in areas where the export nurseries are located. Possible pathways of the pathogen are water and soil.

Uncertainties

- The exact pathways are still uncertain.
- There is a high uncertainty on the prevalence and distribution of this pathogen in the UK in areas where the nurseries are located.

A.3.2.2 | Possibility of entry with new plants/seeds

Plants are produced by seeds and grafting. Seeds purchased in the UK are certified under The Forest Reproductive Material (Great Britain) Regulations 2002 ([legislation.gov.uk](https://www.legislation.gov.uk)); seedlings sourced in the UK are certified with UK Plant Passports; A small percentage of plants may be obtained from EU (Netherlands); seedlings from the EU countries are certified with phytosanitary certificates.

The pathogen is not known to be seedborne or seed transmitted, therefore not expected to enter the nursery via the seed pathway. Grafted plants (scion and buds in the case of grafting) originate from the nursery itself.

The seedling pathways is unlikely because the conditions of their production (seedbed in the greenhouse, pest free growing media) are expected to prevent the infection of seedlings.

The nurseries use virgin peat or peat-free compost (a mixture of coir, tree bark, wood fibre, etc.) as a growing media (Dossier Section 1).

The growing media is heat-treated by commercial suppliers during production to eliminate pests and diseases (Dossier Section 1).

Uncertainties

- None.

Taking into consideration the above evidence and uncertainties, the Panel considers that it is unlikely for the pathogen to enter the nurseries via seeds or seedlings of *Alnus*.

A.3.2.3 | Possibility of spread within the nursery

Phytophthora siskiyouensis has a very narrow host range and it is unlikely that other host plants than *Alnus* spp. plants, such as *Lithocarpus densiflorus* and *Umbellularia californica* are present in the nurseries.

Uncertainties

- There are no uncertainties.

Taking into consideration the above evidence and uncertainties, the Panel considers that it is unlikely for the pathogen to spread from other host plants within the nurseries.

A.3.3 | Information from interceptions

In the EUROPHYT/TRACES-NT database, there are no records of notification of *Alnus* plants for planting neither from the UK nor from other countries due to the presence of *P. siskiyouensis* between the years 1995 and January 2025 (EUROPHYT/TRACES-NT, online).

A.3.4 | Evaluation of the risk mitigation measures

In the table below, all risk mitigation measures currently applied in the UK are listed and an indication of their effectiveness on *P. siskiyouensis* is provided. The description of the implemented risk mitigation measures is provided in [Table 8](#).

No.	Risk mitigation measure	Effect on the pest	Evaluation and uncertainties
1	Registration of production sites	Yes	<p>All nurseries are registered as professional operators with the UK NPPO, by the Animal and Plant Health Agency (APHA) and is authorised to issue UK plant passports (Dossier Section 1).</p> <p><u>Evaluation</u></p> <ul style="list-style-type: none"> – Every nursery exporting to the EU is under supervision of the NPPO. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – None.
2	Certification of plant material	Yes	<p><i>Alnus cordata</i> and <i>A. glutinosa</i> seeds purchased in the UK are certified under The Forest Reproductive Material (Great Britain) Regulations 2002 (legislation.gov.uk); seedlings sourced in the UK are certified with UK Plant Passports; a small percentage of plants may be obtained from EU (Netherlands); seedlings from the EU countries are certified with phytosanitary certificates. (Dossier Section 1).</p> <p>The starting material of <i>Alnus</i> production consists of seed and seedlings. Seeds are certified. Seedlings for production sourced in the UK are certified with UK Plant Passports; seedlings from the EU countries are certified with phytosanitary certificates.</p> <p><u>Evaluation</u>:</p> <ul style="list-style-type: none"> – Despite <i>P. siskiyouensis</i> is not a quarantine pest, it has been assessed by the UK Plant Health Risk Register and attention to its detection is adopted. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – None.
3	Origin and treatment of growing media	Yes	<p>Rooted plants in pots: In the production or procurement of these plants, the use of growing media is assessed for the potential to harbour and transmit plant pests. Growers most commonly use virgin peat or peat-free compost, which is a mixture of coir, tree bark, wood fibre, etc. The compost is heat-treated by commercial suppliers during production to eliminate pests and diseases. It is supplied in sealed bulk bags or shrink-wrapped bales and stored off the ground on pallets, these are completely hygienic and free from contamination. Where delivered in bulk, compost is kept in a dedicated bunker, either indoors, or covered by tarpaulin outdoors, and with no risk of contamination with soil or other material (Dossier Section 1).</p> <p><u>Evaluation</u>:</p> <ul style="list-style-type: none"> – The measure is efficient in preventing the entry of the pathogen via the substrate into the nursery. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – None.
4	Surveillance, monitoring and sampling	Yes	<p>Inspection is carried out at least once a year as part of the Quarantine Surveillance programme (Great Britain uses the same framework for its surveillance programme as the EU). Surveillance is based on visual inspection with samples taken from symptomatic material, and where appropriate, samples are also taken from asymptomatic material (e.g., plants, tubers, soil, watercourses) (Dossier Section 1).</p> <p><u>Evaluation</u>:</p> <ul style="list-style-type: none"> – The surveillance, monitoring and sampling can detect the pathogen. No results are reported. – Due to the canopy size of older trees (up to 25 years old), the pathogen could remain undetected. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> – The efficiency of the surveillance, monitoring and sampling.

(Continues)

(Continued)

No.	Risk mitigation measure	Effect on the pest	Evaluation and uncertainties
5	Hygiene measures	Yes	<p>According to the Dossier Section 1, all the nurseries have plant hygiene and housekeeping rules and practices in place, which are communicated to all relevant employees.</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> These measures could be effective in reducing the risk of introduction and/or spread of the pathogen. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> The efficiency of the hygiene measures performed in the nurseries.
6	Irrigation water quality and/or treatments	Yes	<p>Growers are required to assess water sources, irrigation and drainage systems used in the plant production for the potential to harbour and transmit plant pests. Water is routinely sampled and sent for analysis. No quarantine pests have been found (Dossier Section 1).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> There is no disinfection treatment applied to the irrigation water. However, irrigation water is routinely sampled and tested for quarantine pests. This procedure can reduce the risk. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> The frequency of sampling and the method used for the detection of the pathogen.
7	Application of pest control products	Yes	<p>Crop protection is achieved using a combination of measures including approved plant protection products, biological control or physical measures. Plant protection products are only used when necessary and records of all plant protection treatments are kept (Dossier Section 1).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> Some plant protection products are applied and could reduce the likelihood of the infection by the pathogen, but detailed information is lacking in the Dossier. The efficacy of plant protection products on large trees (up to 25 years old) could be reduced. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> The details about the products applied and the application scheme are unknown, and the efficiency is unclear.
8	Washing of the roots	Yes	<p>Specimen trees (up to 25 years old) are grown in soil for up to 9 years, root washed and placed in pots (based on the request letter of DEFRA to the EU Commission on September 25th, 2023).</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> The washing of the roots removes (parts of) the soil and the pathogen present in the soil. However, the root plant system of plants of 9 years may be difficult to clean completely from soil. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> The effectiveness of the washing to remove all soil with the pathogen.
9	Inspections and management of plants before export	Yes	<p>The UK NPPO carries out inspections and testing where required by the country of destination's plant health legislation, to ensure all requirements are fulfilled and a valid phytosanitary certificate with the correct additional declarations is issued. Separate to any official inspection, plant material is checked by growers for plant health issues before dispatch.</p> <p><u>Evaluation:</u></p> <ul style="list-style-type: none"> The inspections and management of plants before export can detect the pathogen. <p><u>Uncertainties:</u></p> <ul style="list-style-type: none"> Whether early symptoms caused by the pathogen on <i>Alnus</i> species are identified by visual inspections.

A.3.5 | Overall likelihood of pest freedom specimen tree (from 7 to 25 years old)

A.3.5.1 | Comparison with other relevant commodity Risk Assessments involving *Phytophthora siskiyouensis*

Phytophthora siskiyouensis was already assessed as actionable pest for graftwood, bare-root plants, cell-grown plants and rooted plants in pots (of up to 7 years old) of *A. cordata*, *A. glutinosa* and *A. incana* (EFSA PLH Panel, 2025). *Alnus* commodity types evaluated originate from the same nurseries, sharing the same production system. Therefore, for the estimation of the pest freedom level of *P. siskiyouensis* in exported specimen trees (from 7 to 25 years old) of *Alnus* plants, the Panel decided to base the estimated values the ones elicited for the other commodities of *A. cordata*, *A. glutinosa* and *A. incana* (EFSA PLH Panel, 2025).

A.3.5.2 | Reasoning for a scenario which would lead to a reasonably low number of infected potted plants and specimen trees (from 7 to 25 years old)

Phytophthora siskiyouensis has a very restricted distribution in the UK. The scenario assumes a low pressure of the pathogen in the nurseries and in the surroundings. Plants are exported as dormant plants without leaves. The scenario assumes *Alnus* spp. to be minor hosts for the pathogen. The scenario also assumes that symptoms of the disease are visible and promptly detected during inspections.

Furthermore, the extended period during which specimen trees are kept in pots in the nursery (up to 25 years old) would allow soil-borne infections originating from the bare soil cultivation to express symptoms of the disease, therefore be detected during repeated inspections.

A.3.5.3 | Reasoning for a scenario which would lead to a reasonably high number of infected potted plants and specimen trees (from 7 to 25 years old)

The scenario assumes a high pressure of the pathogen in the surrounding environment of the nurseries because suitable hosts are present. The scenario assumes that the pathogen can infect leaves, which may still be present on the plants at the time of export. The scenario also assumes that the pathogen is not detected during the inspections because of the presence of asymptomatic plants or difficulties in recognising early symptoms. Grafting can increase the incidence of the pathogen (via infected buds or by woundings).

In comparison with the other commodities (graftwood, bare-root plants, cell-grown plants and rooted plants in pots), the canopy size of large specimen trees (up to 25 years old) is larger, increasing the difficulties in detecting the pathogen. Furthermore, the exposure time is longer, increasing the probability for the plants to be infected.

A.3.5.4 | Reasoning for a central scenario equally likely to over- or underestimate the number of specimen trees (from 7 to 25 years old) (Median)

Phytophthora siskiyouensis has a very restricted distribution in the UK. In comparison to potted plants up to 7 years old, the specimen trees are generally larger and older up to 25 years old, increasing the time of exposure to the pathogen. The scenario assumes a limited presence of the pathogen in the nurseries and in the surroundings, and a limited reported susceptibility of *Alnus* spp.

A.3.5.5 | Reasoning for the precision of the judgement describing the remaining uncertainties (1st and 3rd quartile/interquartile range)

The limited information on the occurrence of the pathogen in the nurseries and the surroundings and the susceptibility of *Alnus* spp. results in high level of uncertainties.

A.3.6 | Elicitation outcomes of the assessment of the pest freedom for *Phytophthora siskiyouensis* specimen trees (up to 25 years old)

The elicited and fitted values for *P. siskiyouensis* for pest infestation and pest freedom agreed by the Panel are shown in Tables A.5 and A.6 and in Figure A.3.

TABLE A.5 Elicited and fitted values of the uncertainty distribution of pest infestation by *Phytophthora siskiyouensis* per 10,000 specimen trees (from 7 to 25 years old).

Percentile	1%	2.5%	5%	10%	17%	25%	33%	50%	67%	75%	83%	90%	95%	97.5%	99%
Elicited values	0					2		4		5					12
EKE	0.372	0.588	0.842	1.23	1.66	2.13	2.60	3.57	4.75	5.50	6.47	7.62	9.05	10.4	12.0

Note: The EKE results are the *BetaGeneral* (2.1433, 19.658, 0, 41.5) distribution fitted with @Risk version 7.6.

Based on the numbers of estimated infected plants, the pest freedom was calculated (i.e. = 10,000 – number of infected plants per 10,000). The fitted values of the uncertainty distribution of the pest freedom are shown in Table A.6.

TABLE A.6 The uncertainty distribution of plants free of *Phytophthora siskiyouensis* m per 10,000 specimen trees (from 7 to 25 years old) calculated by Table A.5.

Percentile	1%	2.5%	5%	10%	17%	25%	33%	50%	67%	75%	83%	90%	95%	97.5%	99%
Values	9988					9995		9996		9998					10,000
EKE results	9988	9990	9991	9992	9993.5	9994.5	9995.3	9996.4	9997.4	9997.9	9998.3	9998.8	9999.2	9999.4	9999.6

Note: The EKE results are the fitted values.




FIGURE A.3 (Continued)




FIGURE A.3 (Continued)




FIGURE A.3 (A) Elicited uncertainty of pest infection per 10,000 plants of specimen trees (histogram in blue—vertical blue line indicates the elicited percentile in the following order: 1%, 25%, 50%, 75%, 99%) and distributional fit (red line); (B) uncertainty of the proportion of pest-free bundles per 10,000 (i.e. = 1 – pest infection proportion expressed as percentage); (C) descending uncertainty distribution function of pest infection per 10,000 plants.

A.3.7 | Reference list

Abad, Z. G., Burgess, T. I., Redford, A. J., Bienapfl, J. C., Srivastava, S., Mathew, R., & Jennings, K. (2023). IDphy: An international online resource for molecular and morphological identification of Phytophthora. *Plant Disease*, 107(4), 987–998. <https://doi.org/10.1094/PDIS-02-22-0448-FE>

Abad, Z. G., Burgess, T. I., Bourret, T., Bensch, K., Cacciola, S. O., Scanu, B., Mathew, R., Kasiborski, B., Srivastava, S., Kageyama, K., Bienapfl, J. C., Verkleij, G., Broders, K., Schena, L., & Redford, A. J. (2023b). Phytophthora: Taxonomic and phylogenetic revision of the genus. *Studies in Mycology*, 106, 259–349. <https://doi.org/10.3114/sim.2023.106.05>

CABI (Centre for Agriculture & Bioscience International). Phytophthora siskiyouensis. Available online: <https://www.cabidigitallibrary.org/doi/10.1079/cabicompendium.110465#REF-DDB-199728> [accessed 29/10/2024].

CABI (Centre for Agriculture & Bioscience International). Phytophthora siskiyouensis. Available online: <https://www.cabidigitallibrary.org/doi/full/10.5555/20153264575> [accessed 29/10/2024].

EFSA PLH Panel (EFSA Panel on Plant Health) Civera, A. V., Baptista, P., Berlin, A., Chatzivassiliou, E., Cubero, J., Cunniffe, N., de la Peña, E., Desneux, N., Di Serio, F., Filipiak, A., Gonthier, P., Hasiów-Jaroszewska, B., Jactel, H., Landa, B. B., Maistrello, L., Makowski, D., Milonas, P., Papadopoulos, N. T., Susi, H., van der Gaag, D. J., ... Potting, R. (2025). Commodity risk assessment of *Alnus cordata*, *Alnus glutinosa* and *Alnus incana* plants from the UK. *EFSA Journal*, 23(1), e9189. <https://doi.org/10.2903/j.efsa.2025.9189>

EPPO (European and Mediterranean Plant Protection Organization). (online). Phytophthora siskiyouensis (PHYTSK). Available online: <https://gd.eppo.int/taxon/PHYTSK>. [Accessed:29/10/2024].

EUROPHYT, (online). European Union Notification System for Plant Health Interceptions - EUROPHYT. Available online: https://ec.europa.eu/food/plant/plant_health_biosecurity/europhyt/index_en.htm [Accessed: 21/1/2025].

Forest Phytophtoras of the World, a. Siskiyouensis disease. Forest Phytophtoras of the World. Available online: <https://forestphytophtoras.org/species/siskiyouensis/disease> [Accessed: 30/10/2024].

Forest Phytophtoras of the World, b. Phytophthora basics. Forest Phytophtoras of the World. Available online: <https://forestphytophtoras.org/phytophthora-basics> [Accessed: 30/10/2024].

Forest Research. Phytophthora disease of alder (Phytophthora alni). Available online: <https://www.forestryresearch.gov.uk/?s=phytophthora-disease-of-alder%2520phytophthora-almi> [Accessed: 30/10/2024].

Hansen, E. M., Reeser, P., & Rooney-Latham, S. (2011). Phytophthora siskiyouensis. *Forest Phytophtoras*, 1(1). <https://doi.org/10.5399/osu/fp.1.1.1826>

ITP (Identification Technology Program). Available online: <https://idtools.org/phytophthora/index.cfm?packageID=1131&entityID=5176> [Accessed: 30/10/2024].

Navarro, S., Sims, L., Hansen, E., & Vannini, Á. (2015). Pathogenicity to alder of Phytophthora species from riparian ecosystems in western Oregon. *Forest Pathology*, 45(5), 358–366. <https://doi.org/10.1111/fp.121752>

Perez-Sierra, A., Kalantarzadeh, M., Sancisi-Frey, S., & Brasier, C. M. (2015). Phytophthora siskiyouensis causing stem lesions and cankers on *Alnus incana*. *New Disease Reports*, 31, 172. <https://doi.org/10.5197/j.2044-0588.2015.031.017>

Perez-Sierra, A., Sancisi-Frey, S., Kalantarzadeh, M., & Brasier, C. (2014). Unexpected discovery of Phytophthora siskiyouensis in the UK. 7th Meeting IUFRO Working Party 7.02.09 Phytophtora in Forest and Natural Ecosystems, 10–14 November 2014, Esquel, Chubut, Patagonia, Argentina.

Reeser, P. W., Hansen, E. M., & Sutton, W. (2007) Phytophthora siskiyouensis, a new species from soil, water, myrtlewood (*Umbellularia californica*) and tanoak (*Lithocarpus densiflorus*) in southwestern Oregon. *Mycologia*, 99(5), 639–6432.

Reser, P. W., Sutton, W., & Hansen, E. M. (2008). Phytophthora species causing tanoak stem cankers in Southwestern Oregon. *Plant Diseases*, 92, 1252. <https://doi.org/10.1094/PDIS-92-8-1252B>

Rooney-Latham, S., Blomquist, C. L., Pastalka, T., & Costello, L. (2009). Collar rot on Italian alder trees in California caused by Phytophthora siskiyouensis. *PL. Health Programs*, 10(1), 20. <https://doi.org/10.1094/PHP-2009-0413-01-RS>

Sims, L., Goheen, E., Kanaskie, A., & Hansen, E. (2015a). Alder canopy dieback and damage in western Oregon riparian ecosystems. *Northwest Science*, 89(1), 34–46. <https://doi.org/10.3955/046.089.0103>

Sims, L. L., Sutton, W., Reeser, P., & Hansen, E. M. (2015b). The Phytophthora species assemblage and diversity in riparian alder ecosystems of western Oregon, USA. *Mycologia*, 107, 889–902.

Smith, I. W., Cunningham, J., & Pascoe, I. (2006). Another new? Species of Phytophthora on alder 'down under'. (Australia). Poster 30, Progress in Research on Phytophthora Diseases of Forest Trees (C. Brasier, T. Jung, and W. Oßwald, eds.). Proceedings of the Third Meeting of the International Union of Forest Research Organizations (IUFRO) Working Party S07.02.09: Phytophtoras in forests and natural ecosystems. Forest Research, Farnham, Surrey, UK.

Smith, I. A., Cunningham, J., & Pascoe, I. (2004). Another new? Species of Phytophthora on alder down under (Australia). P. of the IUFRO Res. Work. Gr. Conf. on Phytophtora Forests and Natural Ecosystems. 11–17 September 2004. Freising, Germany.

TRACES-NT. (online). TRAde Control and Expert System. Available online: <https://webgate.ec.europa.eu/tracesnt> [Accessed: 21/1/2025].

UK Plant Health Portal, DEFRA. (2024). Available online: <https://planthealthportal.defra.gov.uk/pests-and-diseases/uk-plant-health-riskregister/viewPestRisks.cfm?csref=27280> [Accessed: 31/10/2024].

Waterhouse, G. M. (1963). Key to the species of Phytophthora de Bary. *Mycol. Pap.* 92. 22 pp. Commonwealth Mycological Institute, Kew, UK.

APPENDIX B

Web of Science All Databases Search String 30/01/2024

In the table below, the search string used in Web of Science is reported. In total, 1248 papers were retrieved. Titles and abstracts were screened, and 78 pests were added to the list of pests (Appendix D).

Web of Science All databases	TOPIC: "Alnus incana" OR "Alnus glutinosa" OR "Alnus cordata" OR "Alnus sp." OR "Alnus spp." OR "grey alder" OR "European alder" OR "Italian alder" AND TOPIC: "pathogen*" OR "pathogenic bacteria" OR "fung*" OR "oomycet*" OR "myce*" OR "bacteri*" OR "virus*" OR "viroid*" OR "insect\$" OR "mite\$" OR "phytoplasm*" OR "arthropod*" OR "nematod*" OR "disease\$" OR "infecti*" OR "damag*" OR "symptom*" OR "pest\$" OR "vector" OR "hostplant\$" OR "host plant\$" OR "host" OR "root lesions" OR "decline\$" OR "infestation\$" OR "damages\$" OR "symptom\$" OR "dieback" OR "die back" OR "malaise" OR "aphid\$" OR "curculio" OR "thrip\$" OR "cicad\$" OR "miner\$" OR "borer\$" OR "weevil\$" OR "plant bug\$" OR "spittlebug" OR "moth\$" OR "mealybug\$" OR "cutworm\$" OR "pillbug\$" OR "root feeder\$" OR "caterpillar\$" OR "foliar feeder\$" OR "virosis" OR "viruses" OR "blight\$" OR "wilt\$" OR "wilted" OR "canker" OR "scab" OR "rot" OR "rots" OR "rotten" OR "damping off" OR "damping-off" OR "blister" OR "smut" OR "mould" OR "mould" OR "damping syndrome" OR "mildew" OR "scald\$" OR "root knot" OR "root-knot" OR "rootkit" OR "cyst\$" OR "dagger" OR "plant parasitic" OR "parasitic plant" OR "plant\$parasitic" OR "root feeding" OR "root\$feeding" OR "acari" OR "host\$" OR "gall" OR "gall\$" OR "whitefly" OR "whitefl*" OR "aleyrodidae" OR "thysanoptera" OR "moths" OR "scale" OR "scale\$" OR "thripidae" OR "leafhoppers" OR "leafhopper\$" OR "plant pathogens" OR "fungal" OR "aphididae" OR "Scolytinae" OR "bark beetle" NOT "heavy metal\$" OR "pollut*" OR "weather" OR "propert*" OR "probes" OR "spectr*" OR "antioxidant\$" OR "transformation" OR "Secondary plant metabolite" OR "metabolite" OR "Postharvest" OR "Pollin*" OR "Ethylene" OR "Thinning" OR "fertil*" OR "Mulching" OR "Nutrient\$" OR "human virus" OR "animal disease\$" OR "plant extracts" OR "immunological" OR "purified fraction" OR "traditional medicine" OR "medicine" OR "mammal\$" OR "bird\$" OR "human disease\$" OR "cancer" OR "therapeutic" OR "psoriasis" OR "blood" OR "medicinal ethnobotany" OR "Nitrogen-fixing" OR "patients" OR "Probiotic drugs" OR "Antioxidant" OR "Anti-Inflammatory" OR "plasma levels" OR "ethnomedicinal" OR "traditional uses of medicinal plants" OR "Antitumor" OR "Neuroprotective" OR "Hypoglycemic" OR "ozone sensitivity" OR "cardiotonic" NOT TOPIC: 'Abagrotis variata' OR 'Acalitus brevitarsus' OR 'Acalitus phyllereus' OR 'Acantharia sinensis' OR 'Acanthococcus spiraeae' OR 'Acanthococcus tokaeae' OR 'Acanthosoma haemorrhoidale' OR 'Acanthostigma alni' OR 'Acaphylla trinotus' OR 'Acaricalus trinotus' OR 'Aceria alniborealis' OR 'Aceria alniviridis' OR 'Aceria bistriata' OR 'Aceria longirostris' OR 'Aceria nalepaei' OR 'Acerra normalis' OR 'Achatia distincta' OR 'Achatodes zea' OR 'Achlya flavicornis' OR 'Acleistia alniella' OR 'Acleris alnivora' OR 'Acleris braunana' OR 'Acleris caliginosana' OR 'Acleris cervinana' OR 'Acleris cornana' OR 'Acleris emargana' OR 'Acleris fuscana' OR 'Acleris hudsoniana' OR 'Acleris inana' OR 'Acleris logiana placidana' OR 'Acleris maccana' OR 'Acleris notana' OR 'Acleris semiannula' OR 'Acleris senescens' OR 'Acleris umbrana' OR 'Acmaeodera revelierei' OR 'Acrobasis betulella' OR 'Acrobasis rubrifasciella' OR 'Acronicta alni' OR 'Acronicta americana' OR 'Acronicta auricoma' OR 'Acronicta barnesi' OR 'Acronicta cuspis' OR 'Acronicta dactylinia' OR 'Acronicta distans' OR 'Acronicta euphorbiae' OR 'Acronicta fragilis' OR 'Acronicta funeralis' OR 'Acronicta grisea' OR 'Acronicta hastulifera' OR 'Acronicta hesperida' OR 'Acronicta impleta' OR 'Acronicta impressa' OR 'Acronicta innotata' OR 'Acronicta lanceolaria' OR 'Acronicta leporina' OR 'Acronicta lepusculina' OR 'Acronicta longa' OR 'Acronicta megacephala' OR 'Acronicta oblinita' OR 'Acronicta psi' OR 'Acronicta rumicis' OR 'Acronicta sperata' OR 'Acronicta tridens' OR 'Acrospermum compressum' OR 'Acrostalagmus luteoalbus' OR 'Actebia fennica' OR 'Actias artemis' OR 'Actias gnomae' OR 'Actias luna' OR 'Actias selene' OR 'Actias selene ningpoana' OR 'Aculus epiphyllus' OR 'Adoxophyes orana' OR 'Aegomorphus clavipes' OR 'Aegosoma scabricorne' OR 'Aethalura ignobilis' OR 'Aethalura intertexta' OR 'Aethalura punctulata' OR 'Aeglastica alni' OR 'Aglia tau' OR 'Agnocoris reclairei' OR 'Agonimia repleta' OR 'Agonopterix argillacea' OR 'Agrilus angustulus' OR 'Agrilus graminis' OR 'Agrilus hastulifer' OR 'Agrilus roscidus' OR 'Agrilus viridis' OR 'Agriopsis aurantiaria' OR 'Agriopsis marginaria' OR 'Agroathelia rofsii' OR 'Agrochola litura' OR 'Agrochola lota' OR 'Agrochola pulchella' OR 'Agrochola purpurea' OR 'Agromyza alnivora' OR 'Alatospora acuminata' OR 'Alicis repandata' OR 'Alder yellows phytoplasm' OR 'Alebra albostriella' OR 'Alebra wahlbergi' OR 'Aleuris distinctata' OR 'Aleochara spiniferus' OR 'Aleochara aurantius' OR 'Allygus mixtus' OR 'Allygus modestus' OR 'Alnecium auctum' OR 'Alnetoidea alneti' OR 'Alosterna tabacicolor' OR 'Alsophila aescularia' OR 'Alsophila japonensis' OR 'Alternaria alternata' OR 'Alternaria botrytis' OR 'Alternaria chartarum' OR 'Altica bimarginata' OR 'Altica tamaricis' OR 'Alysium resinace' OR 'Amaurodon mustialaensis' OR 'Ammiculicola parva' OR 'Amorbia humerosana' OR 'Amorpha juglandis' OR 'Ampedus cinnabarinus' OR 'Ampedus elongantulus' OR 'Ampedus nigerrimus' OR 'Ampedus nigrinus' OR 'Ampedus pomonae' OR 'Ampedus pomorum' OR 'Ampedus rufipennis' OR 'Ampedus sanguinolentus' OR 'Amphinema byssoides' OR 'Amphipyra perflua' OR 'Amphipyra pyramidoides' OR 'Amphirosellinia evansii' OR 'Amphisphaeria umbrina' OR 'Anacampsis innocuella' OR 'Anacampsis niveopulvella' OR 'Anaesthetis testacea' OR 'Anagoga occiduaria' OR 'Anaplectoides pressus' OR 'Anavitriella pampinaria' OR 'Andropolia aedon' OR 'Andropolia contacta' OR 'Angerona prunaria' OR 'Anguillosporella vermiformis' OR 'Anguillosporella vermiformis' OR 'Angustimassarina alni' OR 'Anisandrus maiche' OR 'Anomoloma myceliosum' OR 'Anoplodera sexguttata' OR 'Anoplophora chinensis' OR 'Anoplophora glabripennis' OR 'Anoplus plantaris' OR 'Anoplus roboris' OR 'Anoplus setulosus' OR 'Antepione thisoaria' OR 'Antheraea polyphemus polyphemus' OR 'Anthonomus undulatus' OR 'Anthostoma ellisi' OR 'Anthostoma gastrinum' OR 'Anthostoma melanotes' OR 'Antrodia albida' OR 'Antrodia heteromorpha' OR 'Antrodia sinuosa' OR 'Antrodiella genistae' OR 'Antrodiella semisupina' OR 'Apamea auranticolor' OR 'Apamea castanea' OR 'Apateodes torrefacta' OR 'Apatura ilia' OR 'Apatura iris' OR 'Aphelenchoides composticola' OR 'Aphelenchus avenae' OR 'Aphis spiraecola' OR 'Aphrophora alni' OR 'Apionomonia alniella' OR 'Aploporthella bavarica' OR 'Apiospora arundinis' OR 'Apiosporopsis carpinea' OR 'Aplosporella alnicala' OR 'Apocheima hispidaria' OR 'Apoda
---------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

(Continues)

(Continued)

limacodes' OR 'Apodachlyla brachynema' OR 'Apoderus coryli' OR 'Apophlymus pectoralis' OR 'Apotomis bifida' OR 'Apotomis funerea' OR 'Apterona paludella' OR 'Aptinothrips elegans' OR 'Aracima muscosa' OR 'Aradus betulae' OR 'Arboridia erecta' OR 'Arboridia parvula' OR 'Arboridia ribauti' OR 'Archiearis infans oregonensis' OR 'Archips argyrospila' OR 'Archips breviplicanus' OR 'Archips cerasivorana' OR 'Archips crataegana' OR 'Archips fuscocupreanus' OR 'Archips mortuana' OR 'Archips negundana' OR 'Archips podana' OR 'Archips rosana' OR 'Archips viola' OR 'Archips xylosteana' OR 'Arctia caja' OR 'Arctia caja americana' OR 'Arctia caja waroi' OR 'Arctorthezia cataphracta' OR 'Arcyria stipata' OR 'Argyresthia brockeella' OR 'Argyresthia calliphanes' OR 'Argyresthia goedartella' OR 'Argyrotaenia ljungiana' OR 'Argyrotaenia mariana' OR 'Argyrotaenia velutinana' OR 'Armillaria altimontana' OR 'Armillaria cepistipes' OR 'Armillaria gallica' OR 'Armillaria mellea' OR 'Armillaria nabsnona' OR 'Armillaria ostoyae' OR 'Armillaria puiggarii' OR 'Armillaria sinapina' OR 'Armillaria singular' OR 'Arocatus longiceps' OR 'Arocatus melanocephalus' OR 'Arocatus roeselii' OR 'Aromia moschata' OR 'Arthopyrenia didymelloides' OR 'Arthopyrenia grisea' OR 'Artomyces pyxidatus' OR 'Ascocoryne cylindrium' OR 'Ascocoryne sarcoides' OR 'Ascospirella lutea' OR 'Ascotremella faginea' OR 'Aseptis binotata' OR 'Aspergillus nidulans' OR 'Aspergillus niger' OR 'Aspidiotus nerii' OR 'Asterococcus oblates' OR 'Asteroma alneum' OR 'Asteroma alniellum' OR 'Asteroma alnigena' OR 'Asteromassaria alni' OR 'Asteromassaria macrospora' OR 'Asteromella alnicola' OR 'Asteromella ulmi' OR 'Astetholida lucida' OR 'Astrosphaeriella applanata' OR 'Asymmetrasca decedens' OR 'Atemelia torquatella' OR 'Athelia arachnoidea' OR 'Athelia decipiens' OR 'Athelia epiphylla' OR 'Athelia nivea' OR 'Atheliachaete sanguinea' OR 'Atopospora betulina' OR 'Attelabus nitens' OR 'Aureobasidium pullulans' OR 'Auricularia auricula-judae' OR 'Autographa ampla' OR 'Autographa corusca' OR 'Bactrodesmium biformatum' OR 'Bactrodesmium obovatum' OR 'Baeopelma foersteri' OR 'Baileya doubledayi' OR 'Baileya ophthalmica' OR 'Baltazarria galactina' OR 'Barbatosphaeria barbirostris' OR 'Barrmaelia oxyacanthae' OR 'Basidiobolus cinereum' OR 'Basilarchia arthemis' OR 'Basilarchia arthemis arthemis' OR 'Basilarchia arthemis rubrofasciata' OR 'Batia lambdella' OR 'Bertia moriformis' OR 'Besma quercivoria' OR 'Betacallis alnicolens' OR 'Betulaphis quadriflora' OR 'Bibarrambla allenella' OR 'Bipolaris oryzae' OR 'Biscogniauxia bartholomaei' OR 'Biscogniauxia mediterranea' OR 'Biscogniauxia nummularia' OR 'Biscogniauxia repanda' OR 'Bispora effusa' OR 'Biston betularia' OR 'Biston betularia cognataria' OR 'Biston strataria' OR 'Bjerkanderia adusta' OR 'Blastobasis decolorella' OR 'Blepharidopterus angulatus' OR 'Blepharidopterus dubius' OR 'Blepharita adusta' OR 'Boeremia exigua' OR 'Boernerina alni' OR 'Boernerina alni ssp. Insularia' OR 'Boernerina depressa' OR 'Boernerina occidentalis' OR 'Boernerina variabilis' OR 'Bohemannia quadrimaculella' OR 'Bombardia bombarda' OR 'Botryobasidium candidans' OR 'Botryobasidium pruinatum' OR 'Botryobasidium ramosissimum' OR 'Botryobasidium subcoronatum' OR 'Botryobasidium vagum' OR 'Botryobasidium vagum' OR 'Botryodiplodia alni' OR 'Botrytis cinerea' OR 'Bourdottedigloea concisa' OR 'Bourdottedigloea dura' OR 'Bourdottedigloea grisea' OR 'Bourdottedigloea vestita' OR 'Brachysporiella dennisii' OR 'Brachysporiella pulchra' OR 'Brachysporium bloxamii' OR 'Brachysporium britannicum' OR 'Brachysporium fusiforme' OR 'Brachysporium polyseptatum' OR 'Brenneria alni' OR 'Brenneria salicis' OR 'Brevipalpus californicus' OR 'Brevipalpus lewisi' OR 'Brevipalpus obovatus' OR 'Brunnipes calyculiformis' OR 'Bryobia praetiosa' OR 'Bucculatrix cidarella' OR 'Bucculatrix locuples' OR 'Bucculatrix thoracella' OR 'Bulbillomyces farinosus' OR 'Byctiscus betulae' OR 'Byssomerulius corium' OR 'Byssosphaeria alnea' OR 'Caber exanthemata' OR 'Caber purus' OR 'Caber pusaria' OR 'Cactodera betulae' OR 'Cacumisporium capitulatum' OR 'Cadophora melini' OR 'Calaphis alni' OR 'Callidium violaceum' OR 'Callimorpha dominula' OR 'Callistosporium luteo-olivaceum' OR 'Calliteara pudibunda' OR 'Calocera cornea' OR 'Caloptilia alni' OR 'Caloptilia alnicolella' OR 'Caloptilia elongella' OR 'Caloptilia falconipennella' OR 'Caloptilia glutinella' OR 'Caloptilia invariabilis' OR 'Caloptilia issikii' OR 'Caloptilia pulchella' OR 'Caloptilia pulvrea' OR 'Calosphaeria cryptospora' OR 'Calycellina lachnobrachya' OR 'Calycellina leucella' OR 'Calycina alniella' OR 'Calycina alniella' OR 'Calycina citrina' OR 'Calyptella capula' OR 'Camaropella lutea' OR 'Camaropella microspora' OR 'Camarops polysperma' OR 'Campaea margaritaria' OR 'Campaea perlata' OR 'Camposporium cambrense' OR 'Campylospora parvula' OR 'Canephora hirsuta' OR 'Capitotricha bicolor' OR 'Capitotricha scabrovillosa' OR 'Capnodium citri' OR 'Capronia pilosella' OR 'Capua vulgana' OR 'Carpatolechia proximella' OR 'Catocala elocata' OR 'Caviariella aquatica' OR 'Cellypha subgelatinosa' OR 'Celypha rivulana' OR 'Cenangium graddonii' OR 'Cenopalpus lanceolatiseae' OR 'Centrotus cornutus' OR 'Ceraceomyces tessulatus' OR 'Cerambyx scopolii' OR 'Cerastis rubricosa' OR 'Ceratocystiopsis synnemata' OR 'Ceratocystis piceae' OR 'Ceratostomella rostrata' OR 'Ceriopor leptocephalus' OR 'Ceriopor scutellatus' OR 'Ceriopor squamosus' OR 'Ceriopor varius' OR 'Ceriosporopsis cambrensis' OR 'Cerioporia purpurea' OR 'Cerioporia reticulata' OR 'Cerioporia rhodella' OR 'Cerioporia spissa' OR 'Cerioporia viridans' OR 'Ceriporiopsis mucida' OR 'Cerrena unicolor' OR 'Chaenothecopsis savonica' OR 'Chaetomium cochlioides' OR 'Chaetospermum chaetosporum' OR 'Chaetosphaerella phaeostroma' OR 'Chaetosphaeria myriocarpa' OR 'Chaetosphaeria preussii' OR 'Chaetosphaeria pygmaea' OR 'Chaetothyrium setosum' OR 'Chalara alnicola' OR 'Chalara aurea' OR 'Chalara cylindrica' OR 'Chalara inflatipes' OR 'Cheimophila salicella' OR 'Cheiromycella folicola' OR 'Cheirospora alni' OR 'Chilecomadia valdiviana' OR 'Chionaspis alnus' OR 'Chionaspis lintneri' OR 'Chionaspis ortholobis' OR 'Chionaspis salicis' OR 'Chionaspis vistariae' OR 'Chlorencoelia versiformis' OR 'Chloridium caesium' OR 'Chloridium cylindrosporum' OR 'Chlorociboria aeruginascens' OR 'Chlorociboria aeruginosa' OR 'Chloroclysta miata' OR 'Chloroclysta siterata' OR 'Chloroclysta truncata' OR 'Chlorophanus pollinosus' OR 'Chlorophorus glabromaculatus' OR 'Chlorophorus varius' OR 'Chlorosea nevadaria' OR 'Chondrostereum purpureum' OR 'Choreutis betuliperda' OR 'Choreutis diana' OR 'Choristoneura conflictana' OR 'Choristoneura diversana' OR 'Choristoneura rosaceana' OR 'Choristoneura zapulata' OR 'Chromaphis hirsutistibus' OR 'Chrysobothris affinis' OR 'Chrysobothris femorata' OR 'Chrysobothris mali' OR 'Chrysomela aenea' OR 'Chrysomela lapponica' OR 'Chytrium confervae' OR 'Ciboria acerina' OR 'Ciboria alni' OR 'Ciboria amentacea' OR 'Ciboria caucus' OR 'Ciboria semincola' OR 'Ciboria tenuistipes' OR 'Ciboria viridifusca' OR 'Cicadella viridis' OR 'Cimbex connatus' OR 'Cimbex femoratus' OR 'Cimbex luteus' OR 'Cinereomyces lindbladii' OR 'Cingilia catenaria' OR 'Cistella xylita' OR 'Cixius cunicularius' OR 'Cixius nervosus' OR 'Cixius similis' OR 'Cixius stigmaticus' OR 'Cladara atroliturata' OR 'Cladonia caespiticia' OR 'Cladosporium allicinum' OR 'Cladosporium alneum' OR 'Cladosporium alnicola' OR 'Cladosporium cladosporioides' OR 'Cladosporium epiphyllum' OR 'Cladosporium herbarum' OR 'Cladosporium inversicolor' OR 'Cladosporium laxicapitulatum' OR 'Cladosporium lignicola' OR 'Cladosporium macrocarpum' OR 'Cladosporium oxysporum' OR 'Cladosporium sphaerospermum' OR 'Clathrosphaerina zalewskii' OR 'Clavatospora longibrachiata' OR 'Clepsis melaleucana' OR 'Clepsis persicana' OR 'Clethrobius comes' OR 'Clitopilus ardosiacus' OR 'Clostera albosigma' OR 'Clostera inclusa' OR 'Closterotomus fulvomaculatus' OR 'Clytus arietis' OR 'Cochylis nana' OR 'Coeliodes rubicundus' OR 'Coleophora ahenella' OR 'Coleophora alniella' OR 'Coleophora alnifoliae' OR 'Coleophora anatipenella' OR 'Coleophora bindarella' OR 'Coleophora comptoniella' OR 'Coleophora fuscedinella' OR 'Coleophora limosipennella' OR 'Coleophora milvipennis' OR 'Coleophora

(Continued)

orbitella' OR 'Coleophora persimplexella' OR 'Coleophora pruniella' OR 'Coleophora serratella' OR 'Coleophora siccifolia' OR 'Coleophora violacea' OR 'Coleotechnites alnifructella' OR 'Colletotrichum gloeosporioides' OR 'Colocasia coryli' OR 'Colocasia flavidicorns' OR 'Colocasia flavidicorns electa' OR 'Comstockaspis perniciosa' OR 'Conferticium ochraceum' OR 'Coniochaeta dakotensis' OR 'Coniochaeta hoffmannii' OR 'Coniochaeta lignaria' OR 'Coniochaeta lignaria' OR 'Coniochaeta pulveracea' OR 'Coniochaeta subcorticalis' OR 'Coniochaeta velutina' OR 'Coniomela rimicola' OR 'Coniophora arida var. arida' OR 'Coniophora olivacea' OR 'Coniophora puteana' OR 'Coniortodes salicellus' OR 'Coniothyrium fuckelii' OR 'Conistra vaccinii' OR 'Coprinellus micaceus' OR 'Coprinopsis alnivora' OR 'Cordana boothii' OR 'Cordana pauciseptata' OR 'Corinectria fuckeliana' OR 'Corniculariella urceola' OR 'Coronicum alboglaucum' OR 'Coronophora angustata' OR 'Coronophora annexa' OR 'Coronophora gregaria' OR 'Coronophora ovipara' OR 'Corticium boreoreoseum' OR 'Corticium confine' OR 'Corticium roseocarneum' OR 'Corticium roseum' OR 'Corynesporopsis quercicola' OR 'Coryneum sydowianum' OR 'Cosmospora cymosa' OR 'Cosmospora diminuta' OR 'Cosmospora viridescens' OR 'Cossus cossus' OR 'Craniophora ligustris' OR 'Crepidotus alnicola' OR 'Crepidotus cesatii' OR 'Crepidotus fulvotomentosus' OR 'Crepidotus haerens' OR 'Crepidotus mollis' OR 'Crepidotus occidentalis' OR 'Crepidotus ochraceus' OR 'Crepidotus pseudoflammus' OR 'Crepidotus submollis' OR 'Crepidotus subverrucisporus' OR 'Criconema annuliferum' OR 'Criconemoides parvus' OR 'Cristinia coprophila' OR 'Crocallis elinguaria' OR 'Croesus septentrionalis' OR 'Croesus varus' OR 'Crustodontia chrysocreas' OR 'Crustomyces subabruptus' OR 'Cryptonectria parasitica' OR 'Cryptonectria radicalis' OR 'Cryptoblabes bistriga' OR 'Cryptocephalus bipunctatus' OR 'Cryptocephalus coryli' OR 'Cryptocephalus decemmaculatus' OR 'Cryptocephalus pusillus' OR 'Cryptoecoryneum condensatum' OR 'Cryptodiaporthe pyrrhocystis' OR 'Cryptodiaporthe tiliacea' OR 'Cryptorhynchus lapathi' OR 'Cryptosporella alnicola' OR 'Cryptosporella alni-cordata' OR 'Cryptosporella alni-rubrae' OR 'Cryptosporella alni-sinuatae' OR 'Cryptosporella alni-tenuifoliae' OR 'Cryptosporella amistadensis' OR 'Cryptosporella betulae' OR 'Cryptosporella femoralis' OR 'Cryptosporella jaklitschii' OR 'Cryptosporella marylandica' OR 'Cryptosporella multicontinentalis' OR 'Cryptosporella pacifica' OR 'Cryptosporella suffusa' OR 'Cryptosporium neesii' OR 'Cryptosympodula appendiculata' OR 'Cryptovalaria americana' OR 'Cryptovalaria rossica' OR 'Crypturaphis grissii' OR 'Crystallitius serpens' OR 'Cubamyces lactineus' OR 'Cucullia intermedia' OR 'Cucullia lucifuga' OR 'cucumber mosaic virus' OR 'Cucurbitaria alni' OR 'Cucurbitaria fraxini' OR 'Curculio betulae' OR 'Curculio rubidus' OR 'Curvularia lunata' OR 'Cyanosporus caesius' OR 'Cyanostolus aeneus' OR 'Cyathicula amenti' OR 'Cyathicula microspora' OR 'Cyclophora albipunctata' OR 'Cyclophora pendularia' OR 'Cyclophora pendulinaria' OR 'Cydia leguminana' OR 'Cylindrium elongatum' OR 'Cylindrobasidium evolvens' OR 'Cylindrobasidium laeve' OR 'Cylindrotrichum clavatum' OR 'Cyphella fasciculata' OR 'Cyphella fulva' OR 'Cyrtoclytus capra' OR 'Cystostereum pini-canadensis' OR 'Cytidia salicina' OR 'Cytidiella albida' OR 'Cytospora ceratosperma' OR 'Cytospora coenobitica' OR 'Cytospora diatrypa' OR 'Cytospora diatrypoides' OR 'Cytospora leucosperma' OR 'Cytospora leucostoma' OR 'Cytospora melanodiscus' OR 'Cytospora nastroma' OR 'Cytospora populina' OR 'Cytospora pulcherrima' OR 'Cytospora stenospora' OR 'Cytospora truncata' OR 'Cytospora umbrina' OR 'Cytosporella antarctica' OR 'Cytosporina ludibunda' OR 'Dacrymyces capitatus' OR 'Dacrymyces chrysospermus' OR 'Dacrymyces stillatus' OR 'Dactylaria candidula' OR 'Dactylaria obtriangularia' OR 'Daedaleopsis confragosa' OR 'Daedaleopsis nipponica' OR 'Daedaleopsis tricolour' OR 'Daldinia andina' OR 'Daldinia barkalovii' OR 'Daldinia childiae' OR 'Daldinia concentrica' OR 'Daldinia decipiens' OR 'Daldinia lloydii' OR 'Daldinia loculata' OR 'Daldinia occidentalis' OR 'Daldinia petriniae' OR 'Daldinia vernicosa' OR 'Dasineura tortilis' OR 'Dasychira vagans grisea' OR 'Dasyscyphella dryina' OR 'Dematiocypha catenate' OR 'Dematiocypha dematiicola' OR 'Dematiocypha olivacea' OR 'Dematiocypha richonis' OR 'Dematophora necatrix' OR 'Dendrocorticium violaceum' OR 'Dendrophoma merizophila' OR 'Dendrophora erumpens' OR 'Dendroplella multisepata' OR 'Dendrocyter marmaroides' OR 'Dendrothrips degeeri' OR 'Dendrothrips ornatus' OR 'Dendrothrips saltatrix' OR 'Denticollis linearis' OR 'Dentocorticium portoricense' OR 'Deporaus betulae' OR 'Deroplia genei' OR 'Diacrisia sannio' OR 'Dialonectria episphaeria' OR 'Diaporthe alnea' OR 'Diaporthe eres' OR 'Diaporthe nivosa' OR 'Diaporthe padi var. padi' OR 'Diaporthe rufidis' OR 'Diaporthe valsiformis' OR 'Diaporthe verrucella' OR 'Diasria esurialis' OR 'Diaspidiotus aesculi' OR 'Diaspidiotus alni' OR 'Diaspidiotus gigas' OR 'Diaspidiotus ostreaeformis' OR 'Diaspidiotus wuenni' OR 'Diatrype bullata' OR 'Diatrype disciformis' OR 'Diatrype macounii' OR 'Diatrype megastoma' OR 'Diatrype stigma' OR 'Diatrypella betulina' OR 'Diatrypella decorata' OR 'Diatrypella discoidea' OR 'Diatrypella favacea' OR 'Diatrypella placenta' OR 'Diatrypella rimosa' OR 'Dibeloniella citrinella' OR 'Dicerca aenea' OR 'Dicerca alni' OR 'Dicerca berolinensis' OR 'Dichostereum effuscatum' OR 'Dichostereum pallescens' OR 'Dichotomopilus funicola' OR 'Diderma radiatum' OR 'Didymosphaeria dochmia' OR 'Didymosphaeria nana' OR 'Didymosphaeria oblitescens' OR 'Didymosphaeria oregonensis' OR 'Diplococcum lawrencei' OR 'Diplococcum spicatum' OR 'Diplodia alni' OR 'Diplodia cavanillesiana' OR 'Diplodia seriata' OR 'Diptacus dipterochelus' OR 'Diptacus sacramentae' OR 'Ditiola peziziformis' OR 'Ditopella aseptatospora' OR 'Ditopella biseptata' OR 'Ditopella cryptosphaeria' OR 'Ditopella ditopa' OR 'Ditopellosis alni' OR 'Ditylenchus intermedius' OR 'Ditylenchus myceliophagus' OR 'Diurnea fagella' OR 'Diurnea lipsiella' OR 'Dolba hyloeus' OR 'Donkioporia expansa' OR 'Dothidea collecta' OR 'Dothiora europaea' OR 'Dothiorella guttulata' OR 'Dothiorella sarmentorum' OR 'Drepana arcuata' OR 'Drepana bilineata' OR 'Drepana curvata' OR 'Drepana falcata' OR 'Drymochares truquii' OR 'Drymus brunneus' OR 'Dryocoetinus alni' OR 'Dryocoetinus villosus' OR 'Durandiella alni' OR 'Durella melanochlora' OR 'Dysstroma citrata' OR 'Dysstroma truncata' OR 'Dysstroma walkerae' OR 'Eacles imperialis' OR 'Eacles imperialis imperialis' OR 'Echinosphaeria strigosa' OR 'Ectoedemia minimella' OR 'Ectopsocus petersi' OR 'Ectropis crepuscularia' OR 'Ectropis excellens' OR 'Edwardsiana alnicola' OR 'Edwardsiana bergmani' OR 'Edwardsiana candidula' OR 'Edwardsiana crataegi' OR 'Edwardsiana geometrica' OR 'Edwardsiana gratiosa' OR 'Edwardsiana helva' OR 'Edwardsiana hippocastani' OR 'Edwardsiana lanternae' OR 'Edwardsiana lethierryi' OR 'Edwardsiana menzbieri' OR 'Edwardsiana plebeja' OR 'Edwardsiana plurispinosa' OR 'Edwardsiana rosea' OR 'Edwardsiana sardoa' OR 'Edwardsiana soror' OR 'Edwardsiana spinigera' OR 'Efibula avellanea' OR 'Egira crucialis' OR 'Egira hiemalis' OR 'Egira rubrica' OR 'Egira simplex' OR 'Eilema griseola' OR 'Elasmostethus interstinctus' OR 'Elasmostethus minor' OR 'Elasmucha antennata' OR 'Elasmucha fiebri' OR 'Elasmucha grisea' OR 'Electrophaes corylata' OR 'Ellisembia coronata' OR 'Elmerina caryae' OR 'Elodes marginata' OR 'Elpiste lorquinaria' OR 'Empoasca decedens' OR 'Enargia decolor' OR 'Encoelia furfuracea' OR 'Endocliata auratus' OR 'Endocliata undulifer' OR 'Endophragmiella angustispora' OR 'Endophragmiella collapsa' OR 'Endophragmiella ovoidea' OR 'Endophragmiella pallescens' OR 'Endromis versicolora' OR 'Ennomos alniaria' OR 'Ennomos autumnaria' OR 'Ennomos erosaria' OR 'Ennomos fuscantaria' OR 'Ennomos magnaria' OR 'Ennomos quercinaria' OR 'Entoleuca mammata' OR 'Eopyrenula leucoplaca' OR 'Eotetranychus carpini' OR 'Eotetranychus kankitus' OR 'Eotetranychus pallidus' OR 'Eotetranychus tiliarium' OR 'Eotetranychus uncatus' OR 'Epicoccum nigrum' OR 'Epinotia albanguiana' OR 'Epinotia corylana' OR 'Epinotia cruciana' OR 'Epinotia demarniana' OR 'Epinotia immundana' OR

(Continues)

(Continued)

'Epinotia rectiplicana' OR 'Epinotia rubicana' OR 'Epinotia solandriana' OR 'Epinotia sordidana' OR 'Epinotia subculana' OR 'Epinotia tenerana' OR 'Epinotia tetraquetra' OR 'Epinotodonta fumosa' OR 'Epione paralellaria' OR 'Epione repandaria' OR 'Epiphegia microcarpa' OR 'Epiphyas postvittana' OR 'Epirrita autumnata' OR 'Epirrita christyi' OR 'Epirrita dilutata' OR 'Episimus argutana' OR 'Epitimerus longitarsus' OR 'Erannis defoliaria' OR 'Erannis golda' OR 'Erannis tiliaria' OR 'Erannis tiliaria vancouverensis' OR 'Erastia aurantiaca' OR 'Ergates faber' OR 'Eriocampa ovata' OR 'Eriocampa umbratica' OR 'Eriocrania alpinella' OR 'Eriogaster arbusculae' OR 'Eriogaster lanestris' OR 'Eriophyes alniicanae' OR 'Eriophyes axillaris' OR 'Eriophyes euryporus' OR 'Eriophyes inangulis' OR 'Eriophyes laevis' OR 'Erysiphe aggregata' OR 'Erysiphe amanoii' OR 'Erysiphe aquilegiae' OR 'Erysiphe betulina' OR 'Erysiphe miyabei' OR 'Erysiphe miyabei' OR 'Erysiphe penicillata' OR 'Erysiphe pisii' OR 'Erysiphe salicis' OR 'Erysiphe vernalis' OR 'Erythricium laetum' OR 'Erythroneura angusta' OR 'Estigmene acrea' OR 'Euantennaria alaskensis' OR 'Euceraphis betulae' OR 'Euceraphis betulijaponicae' OR 'Euceraphis caerulescens' OR 'Euceraphis gillettei' OR 'Euceraphis ontakensis' OR 'Euchlaena marginaria' OR 'Euchlaena marginaria albertanensis' OR 'Eucephaea nebulata' OR 'Euclea delphinii' OR 'Eucolaspis brunnea' OR 'Eufidonia discospilata' OR 'Eulecanium alnicola' OR 'Eulecanium ciliatum' OR 'Eulecanium douglasi' OR 'Eulecanium tiliae' OR 'Eulia ministrana' OR 'Eulithis destinata' OR 'Eulithis testata' OR 'Eulithis xyli' OR 'Eulithis xyli speciosa' OR 'Eupithecia anticaria' OR 'Eupithecia columbiata' OR 'Eupithecia columbiata columbiata' OR 'Eupithecia columbiata holbergata' OR 'Eupithecia exiguata' OR 'Eupithecia fletcherata' OR 'Eupithecia gelidata' OR 'Eupithecia harrisonata' OR 'Eupithecia lachrymosa' OR 'Eupithecia maestosa' OR 'Eupithecia misturata' OR 'Eupithecia multistrigata' OR 'Eupithecia perfusca' OR 'Eupithecia pseudotsugata' OR 'Eupithecia ravocostaliata' OR 'Eupithecia satyra dodata' OR 'Eupithecia sheppardata' OR 'Eupithecia strattonata' OR 'Eupithecia subfuscata' OR 'Euplexia benesimilis' OR 'Euplexia lucipara' OR 'Euproctis similis' OR 'Eupsilia tristigmata' OR 'Eupterycyba jucunda' OR 'Eurhadina concinna' OR 'Eurhadina pulchella' OR 'Eurhadina ribauti' OR 'Eurois astrcta' OR 'Eurois occulta' OR 'Eutrichosiphum alnicola' OR 'Eutrichosiphum alnifoliae' OR 'Eutrichosiphum alnisuctum' OR 'Eutrichosiphum nepalensis' OR 'Eutrichosiphum raychaudhuri' OR 'Eutrichosiphum tattakanum' OR 'Eutypa flavovirens' OR 'Eutypa lata' OR 'Eutypa stenopora' OR 'Eutypella acericola' OR 'Eutypella alnifraga' OR 'Eutypella alpina' OR 'Eutypella cerviculata' OR 'Eutypella glandulosa' OR 'Eutypella leprosa' OR 'Eutypella persica' OR 'Eutypella stellulata' OR 'Eutypella tetraploa' OR 'Euura glutinosae' OR 'Euura pavid' OR 'Euura umbrata' OR 'Euura viridis' OR 'Euwallacea fornicate sensu lato' OR 'Euwallacea fornicate sensu stricto' OR 'Euwallacea kuroshio' OR 'Evora hemidesma' OR 'Exaeretia ciniflonella' OR 'Excipularia fusispora' OR 'Exidia cartilaginea' OR 'Exidia crenata' OR 'Exidia glandulosa' OR 'Exidia repanda' OR 'Exidia saccharina' OR 'Exidiopsis effusa' OR 'Exidiopsis molybdea' OR 'Exocentrus adspersus' OR 'Exocentrus punctipennis' OR 'Exochalara longissima' OR 'Fagocyba alnisuga' OR 'Fagocyba cruenta' OR 'Falcaria lacertinaria' OR 'Favolus alveolaris' OR 'Fellhanera gyrophorica' OR 'Fenestella fenestrata' OR 'Fenestella leucostoma' OR 'Fenestella minor' OR 'Fenestella princeps' OR 'Fenestella subvestita' OR 'Feniseca tarquinii novascotiae' OR 'Fenus dohrnii' OR 'Fenus pumila' OR 'Fenusella nana' OR 'Fibricium rude' OR 'Fibroporia destructor' OR 'Fibroporia vaillantii' OR 'Filosporella versimorpha' OR 'Fishia evelina' OR 'Flagelloscypha minutissima' OR 'Flagellospora curvula' OR 'Flammula alnicola' OR 'Flammulaster carpophilus' OR 'Flammulina velutipes' OR 'Fomes fomentarius' OR 'Fomitiporia punctata' OR 'Fomitopsis pinicola' OR 'Fonscolombia rotunda' OR 'Furcula bicuspis' OR 'Fusarium arthrosporioides' OR 'Fuscoporia contigua' OR 'Fuscoporia ferrea' OR 'Fuscoporia ferruginosa' OR 'Fuscoporia gilva' OR 'Galerina pallidispora' OR 'Galerucella lineola' OR 'Galerucella solarii' OR 'Galerucella tenella' OR 'Gamsomycetes longisporus' OR 'Ganoderma applanatum' OR 'Ganoderma australe' OR 'Gastrosarbus nigricollis' OR 'Gazalina chrysolopha' OR 'Geometra papilionaria' OR 'Globisporangium intermedium' OR 'Gloeocystidiellum luridum' OR 'Gloeocystidiellum porosellum' OR 'Gloeocystidiellum porosum' OR 'Gloeodontia columbiensis' OR 'Gloeophyllum odoratum' OR 'Gloeophyllum sepiarium' OR 'Gloeoporus ambiguus' OR 'Gloeoporus pannocinctus' OR 'Gloeosporium alnicola' OR 'Gloeosporium cylindrospermum' OR 'Gloiothele citrina' OR 'Gloiothele lactescens' OR 'Glyphisia septentrionalis' OR 'Glyphina betulae' OR 'Glyphina jacutensis' OR 'Glyphium corrugatum' OR 'Glyphium elatum' OR 'Gnomonia alnea' OR 'Gnomonia gnomon' OR 'Gnomonia nervisequa' OR 'Gnomonia pversa' OR 'Gnomoniella alnobetulae' OR 'Gnomoniella gnomon' OR 'Gnomoniella tubaeformis' OR 'Godronia cassandrae' OR 'Godronia fuliginosa' OR 'Gomphinaria amoena' OR 'Gonioctena flavicornis' OR 'Gonioctena interposita' OR 'Gonioctena pallida' OR 'Gonioctena viminalis' OR 'Gonocerus acuteangulatus' OR 'Gossyparia spuria' OR 'Grammoptera ruficornis' OR 'Grammoptera ustulata' OR 'Grapevine flavescent dorée phytoplasma' OR 'Graphium penicilliooides' OR 'Greenidea manii' OR 'Greenidea myriaca' OR 'Gretchenia dulciana' OR 'Gretchenia semialba' OR 'Gretchenia watchungana' OR 'Grynobius planus' OR 'Guepiniopsis alpina' OR 'Guignardia alnigena' OR 'Gymnopus androsaceus' OR 'Gyrodon lividus' OR 'Gyrophanopsis polonensis' OR 'Halysidota tessellaris' OR 'Hannabura alnicola' OR 'Hannabura alnosa' OR 'Hapalopilus rutilans' OR 'Hapalopilus rutilans' OR 'Haploa confusa' OR 'Haploporus odorus' OR 'Hedya dimidioalba' OR 'Hedya nubiferana' OR 'Helicobasidium mompa' OR 'Helicogloea pellucida' OR 'Helicogloea pinicola' OR 'Helicogloea septifera' OR 'Helicoma microscopicum' OR 'Helicoma muelleri' OR 'Helicosporium vegetum' OR 'Helicotylenchus anhelicus' OR 'Helicotylenchus digonicus' OR 'Helicotylenchus dihystera' OR 'Helicotylenchus erythrinae' OR 'Heliococcus boemicus' OR 'Heliozela resplendella' OR 'Helminthosphaeria odontiae' OR 'Helminthosporium italicum' OR 'Helminthosporium lusitanicum' OR 'Helminthosporium velutinum' OR 'Hemicroa australis' OR 'Hemicroa crocea' OR 'Hemicronemoides californianus' OR 'Hemicronemoides mangiferae' OR 'Hemicyclophora typica' OR 'Hemileuca nevadensis-complex' OR 'Hemimycena crispa' OR 'Hemithea aestivaria' OR 'Hendersoniopsis thelebola' OR 'Hericium coralloides' OR 'Hericium erinaceus' OR 'Herminia grisealis' OR 'Hesium domino' OR 'Hesperumia sulphuraria' OR 'Heterarthrus vagans' OR 'Heterobasidion annosum' OR 'Heterobasidion parviporum' OR 'Heterocampa biundata' OR 'Heterocordylus tumidicornis' OR 'Heteroradulum spinulosum' OR 'Hilberina breviseta' OR 'Hilberina rufa' OR 'Hohenbuehelia atrocoerulea' OR 'Hohenbuehelia constans var. alni' OR 'Hohenbuehelia petaloïdes' OR 'Hohenbuehelia spathulata' OR 'Homoglaea hircina' OR 'Homonopsis rubens' OR 'Homophrone naucoria' OR 'Homorthodes communis' OR 'Homostegia obscura' OR 'Hoplotylus femina' OR 'Hoplotylus silvaticus' OR 'Humicola fuscoatra' OR 'Hyalodendriella betulae' OR 'Hyalopeziza alni' OR 'Hyalopeziza digitipila' OR 'Hyalopeziza millepunctata' OR 'Hyalophora cecropia' OR 'Hyalophora columbia' OR 'Hyalophora columbia gloveri' OR 'Hyalophora euryalus' OR 'Hyaloscypha albohyalina' OR 'Hyaloscypha carpinacea' OR 'Hyaloscypha fuckelii' OR 'Hyaloscypha hyalina' OR 'Hyaloscypha spinulosa' OR 'Hydnomerulius pinastri' OR 'Hydnoporia corrugata' OR 'Hydnoporia fuscescens' OR 'Hydnoporia olivacea' OR 'Hydnoporia tabacina' OR 'Hydrelia flammeolaria' OR 'Hydrelia sylvata' OR 'Hydria undulata' OR 'Hydriomena furcata' OR 'Hydriomena impluviata' OR 'Hydriomena irata' OR 'Hydriomena nubilofasciata' OR 'Hydriomena pluviata' OR 'Hydriomena renunciata' OR 'Hydriomena ruberata' OR 'Hydrocina chaetocladia' OR 'Hygrophoropsis aurantiaca' OR 'Hylecoetus dermestoides' OR 'Hylotrupes bajulus' OR 'Hymenochaete cinnamomea' OR 'Hymenochaete pinnatifida' OR 'Hymenochaete spreta' OR 'Hymenochaetopsis intricata' OR 'Hymenoscyphus albopunctus' OR

(Continued)

'Hymenoscyphus calyculus' OR 'Hymenoscyphus caudatus' OR 'Hymenoscyphus epiphyllus' OR 'Hymenoscyphus fastidiosus' OR 'Hymenoscyphus flavofuscens' OR 'Hymenoscyphus imberbis' OR 'Hymenoscyphus rufescens' OR 'Hymenoscyphus tetracadius' OR 'Hymenoscyphus vernus' OR 'Hypagyrtis unipunctata' OR 'Hypatima rhomboidella' OR 'Hypena sordidula' OR 'Hyphantria cunea' OR 'Hyphoderma amoenum' OR 'Hyphoderma cristulatum' OR 'Hyphoderma litschaueri' OR 'Hyphoderma obtusiforme' OR 'Hyphoderma pilosum' OR 'Hyphoderma setigerum' OR 'Hyphoderma tenue' OR 'Hyphodermella corrugata' OR 'Hyphodiscosia mirabilis' OR 'Hyphodiscus hymeniophilus' OR 'Hyphodontia alutaria' OR 'Hyphodontia arguta' OR 'Hyphodontia barba-jovis' OR 'Hyphodontia pallidula' OR 'Hyphodontia quericina' OR 'Hyphodontia spathulata' OR 'Hyphodontia stipata' OR 'Hypochnicum bombycinum' OR 'Hypochnicum geogenium' OR 'Hypochnicum punctulatum' OR 'Hypochnicum sphaerosporum' OR 'Hypocrea aureo-viridis' OR 'Hypocrea gelatinosa' OR 'Hypomecis punctinalis' OR 'Hypomecis roboraria' OR 'Hypomyces gamsii' OR 'Hypospila californica' OR 'Hypothenemus eruditus' OR 'Hypoxylon fragiforme' OR 'Hypoxylon fuscum' OR 'Hypoxylon howeanum' OR 'Hypoxylon julianii' OR 'Hypoxylon macrosporum' OR 'Hypoxylon peckianum' OR 'Hypoxylon rubiginosum' OR 'Hypoxylon vogesiacum' OR 'Hyppa brunneicrista' OR 'Hyppa xyloides' OR 'Hypsizygus elongatipes' OR 'Hysterium angustum' OR 'Hysterium pulicare' OR 'Hysterographium fraxini' OR 'Idiocerus stigmatis' OR 'Idiocerus vitreus' OR 'Idiodonus cruentatus' OR 'Illinoia alni' OR 'Illinoia wilhelminae' OR 'Immersiella caudata' OR 'Incurvaria pectinea' OR 'Inonotus hispidus' OR 'Inonotus luteoumbrinus' OR 'Inonotus mikadoi' OR 'Inonotus obliquus' OR 'Ipimorpha retusa' OR 'Irantylenchus vicinus' OR 'Iridopsis emasculata' OR 'Iridopsis larvaria' OR 'Irpea crenicolor' OR 'Irpea lacteum' OR 'Irpea owensis' OR 'Ischnoderma resinosum' OR 'Isotomus speciosus' OR 'Itame anataria' OR 'Itame bitactata' OR 'Itame exauspicata' OR 'Itame loricaria julia' OR 'Jaapiella clethrophila' OR 'Jackrogersella cohaerens' OR 'Jackrogersella multifloris' OR 'Jattaea taediosa' OR 'Jodis lactearia' OR 'Jodis urosticta' OR 'Jumillera hypophlaea' OR 'Junewangia globulosa' OR 'Junghuhnia nitida' OR 'Junghuhnia subfimbriata' OR 'Karstenia alnicola' OR 'Kirschsteiniothelia atra' OR 'Kirschsteiniothelia recessa' OR 'Kleidocerys privignus' OR 'Kleidocerys resedae' OR 'Kneiffia subalutacea' OR 'Kretzschmaria deusta' OR 'Kretzschmaria zonata' OR 'Kuehneromyces mutabilis' OR 'Kybos mucronatus' OR 'Kybos smaragdula' OR 'Kybos strobli' OR 'Lacanobia contigua' OR 'Lacanobia radix' OR 'Lacanobia suasa' OR 'Lacanobia subjuncta' OR 'Lacanobia thalassina' OR 'Lachnum alneum' OR 'Lachnum hyalopus' OR 'Lachnum roridum' OR 'Lachnum virgineum' OR 'Lacinipolia cuneata' OR 'Lactarius lilacinus' OR 'Laetiporus sulphureus' OR 'Lambdina fiscellaria' OR 'Lambdina fiscellaria lugubrosa' OR 'Lambdina fiscellaria somniaria' OR 'Lamia textor' OR 'Lamprodila decipiens' OR 'Lamprodila mirifica' OR 'Lamprotettix nitidulus' OR 'Laethoe populi' OR 'Lasiochampa quercus' OR 'Lasiodiplodia theobromae' OR 'Lasionycta perplexa' OR 'Lasiosphaeria ovina' OR 'Lasiosphaeria pallida' OR 'Lasiosphaeria hirsuta' OR 'Lasiosphaeria hispida' OR 'Latgerina orizabaensis' OR 'Latgerina orizabaensis ssp. Mexicana' OR 'Laxitextum bicolor' OR 'Ledra aurita' OR 'Leiopus linnei' OR 'Leiopus nebulosus' OR 'Lentinellus cochleatus' OR 'Lentinellus flabelliformis' OR 'Lentinellus micheneri' OR 'Lentinus arcularius' OR 'Lentinus brumalis' OR 'Lentinus substrictus' OR 'Lentithecium aquaticum' OR 'Lenzites betulinus' OR 'Lenzites saeparia' OR 'Lenzites trabea' OR 'Lepidosaphes alnicola' OR 'Lepidosaphes conchiformis' OR 'Lepidosaphes kashicola' OR 'Lepidosaphes tubulorum' OR 'Lepidosaphes ulmi' OR 'Lepidosaphes ussuriensis' OR 'Lepidosaphes yanagicola' OR 'Lepiota clypeolaria' OR 'Lepraria lobificans' OR 'Leptogium hildenbrandii' OR 'Leptographium alneum' OR 'Leptographium piriforme' OR 'Leptographium tardum' OR 'Leptographium trypodendri' OR 'Leptosphaeria lonicerina' OR 'Leptosphaeria vagabunda' OR 'Leptosporomyces galzinii' OR 'Leptothyrium alneum' OR 'Leptoxiphium fumago' OR 'Leptura aethiops' OR 'Leptura annularis' OR 'Leptura aurulenta' OR 'Leptura quadrifasciata' OR 'Lepyrus capucinus' OR 'Leucobrepheos brephoides' OR 'Leucoptera malifoliella' OR 'Libythea celtis' OR 'Licrostoma subgiganteum' OR 'Lindbergina aurovittata' OR 'Lindtneria chordulata' OR 'Lindtneria leucobryophila' OR 'Linnavuoriana intercedens' OR 'Liothrips pragensis' OR 'Lipartrum mandibulare' OR 'Lirimiris truncata' OR 'Litholomia napaea' OR 'Lithomoia solidaginis' OR 'Lithophane amanda' OR 'Lithophane consocia' OR 'Lithophane dilatocula' OR 'Lithophane fagina' OR 'Lithophane furcifera' OR 'Lithophane furcifera suffusa' OR 'Lithophane georgii' OR 'Lithophane innominata' OR 'Lithophane merckii' OR 'Lithophane petulca' OR 'Lithophane pexata' OR 'Lithophane socia' OR 'Lithophane thaxteri' OR 'Lithophane vivida' OR 'Lobesia reliquana' OR 'Lobophora nivigerata' OR 'Lobulomyces poculatus' OR 'Lochmaea caprea' OR 'Lomanaltes eductalis' OR 'Lomographa semioclara' OR 'Lomographa temerata' OR 'Longidorus aetnaeus' OR 'Longidorus distinctus' OR 'Longidorus elongatus' OR 'Longidorus iliturgiensis' OR 'Longidorus paralaskaensis' OR 'Lophiostoma aquaticum' OR 'Lophiostoma glabrotunicatum' OR 'Lophiostoma rugulosum' OR 'Lophiotrema boreale' OR 'Lophocampa argentata' OR 'Lophocampa caryae' OR 'Lophocampa maculata' OR 'Lophocosma atriplaga' OR 'Lopholeucaspis japonica' OR 'Loweomyces fractipes' OR 'Lunulospora curvula' OR 'Luperus flavipes' OR 'Luperus longicornis' OR 'Luperus luperus' OR 'Luperus viridipennis' OR 'Lycia hirtaria' OR 'Lycia pomonaria' OR 'Lycia rachelae' OR 'Lycia ursaria' OR 'Lycogala epidendrum' OR 'Lycoperdon excipuliforme' OR 'Lycophotia phyllophora' OR 'Lycorma delicatula' OR 'Lygocoris contaminatus' OR 'Lygocoris pabulinus' OR 'Lygocoris rugicollis' OR 'Lygocoris viridis' OR 'Lymantria dispar' OR 'Lymantria monacha' OR 'Lymantria obfuscata' OR 'Lyomyces crustosus' OR 'Lyomyces pruni' OR 'Lyomyces sambuci' OR 'Lyonetia alniella' OR 'Lyonetia saliciella' OR 'Macaria alternata' OR 'Macaria notata' OR 'Macrodiaportha occulta' OR 'Macrolabis alnicola' OR 'Macroleptura thoracica' OR 'Macrophomina phaseolina' OR 'Macrophya montana' OR 'Macrotyphula phacorrhiza' OR 'Maireina ochracea' OR 'Malacosoma americanum' OR 'Malacosoma californica' OR 'Malacosoma californica californica' OR 'Malacosoma californica pluvialis' OR 'Malacosoma disstria' OR 'Malacosoma neustria' OR 'Marasmiellus phaeophyllus' OR 'Marasmiellus roseipallens' OR 'Margaritispora aquatica' OR 'Massaria alpina' OR 'Massaria ulmi' OR 'Massaria eburnea' OR 'Massaria leucosarca' OR 'Massariovalsa megalospora' OR 'Megacoelum infusum' OR 'Megacolbyia platyphylla' OR 'Megalocystidium leucoxanthum' OR 'Megalocystidium leucoxanthum' OR 'Megapenthes lugens' OR 'Melampsoridium alni' OR 'Melampsoridium betulinum' OR 'Melampsoridium hiratsukanum' OR 'Melanchra adjuncta' OR 'Melanchra assimilis' OR 'Melanchra persicariae' OR 'Melanchra pisi' OR 'Melanconiopsis megalospora' OR 'Melanconis alni' OR 'Melanconis marginalis' OR 'Melanconis pacifica' OR 'Melanconis stilbostoma' OR 'Melanconium apiocarpum' OR 'Melanconium bicolor' OR 'Melanconium magnum' OR 'Melanconium sphaeroideum' OR 'Melangyna arctica' OR 'Melanolophia canadaria' OR 'Melanolophia imitata' OR 'Melanolophia signataria' OR 'Melanomma pulvis-pyrius' OR 'Melanophila acuminata' OR 'Melanopsamma pomiformis' OR 'Melanopsamma pomiformis' OR 'Melasma alni' OR 'Meliscaevae auricollis' OR 'Melogramma campylosporum' OR 'Melioidodera sitkhotealiniensis' OR 'Melzericium udicola' OR 'Menispora caesia' OR 'Menispora ciliata' OR 'Menispora glauca' OR 'Merismodes anomala' OR 'Merismodes anomala' OR 'Merismodes connivens' OR 'Merismodes fasciculata' OR 'Merlinius brevidens' OR 'Merlinius microdorus' OR 'Merlinius tartuensis' OR 'Mesites tardii' OR 'Mesocallis alnicola' OR 'Mesocallis obtusirostris' OR 'Mesocallis pteleae' OR 'Mesocallis taoi' OR 'Mesogona oxalina' OR 'Mesosa curculionoides' OR 'Mesosa nebulosa' OR 'Mesothea incertata viridipennata' OR 'Metarranthis duaria' OR

(Continues)

(Continued)

'Metendothenia atropunctana' OR 'Metriotes lutarea' OR 'Microascus brevicaulis' OR 'Microdiplodia alni' OR 'Micropeltella grummanniana' OR 'Microporus longisporus' OR 'Microsebacina microbasidia' OR 'Mimas tiliae' OR 'Mimocoris rugicollis' OR 'Miris striatus' OR 'Mitrula borealis' OR 'Mitrula lunulatospora' OR 'Mniotype adusta' OR 'Mniotype bathensis' OR 'Mniotype miniota' OR 'Mollisia alnicola' OR 'Mollisia amenticola' OR 'Mollisia caespiticia' OR 'Mollisia cinerea' OR 'Mollisia ramealis' OR 'Mollisia uda' OR 'Mollisia ventosa' OR 'Mollitrichosiphum montanum' OR 'Mollitrichosiphum nandii' OR 'Mollitrichosiphum niitakaensis' OR 'Mollitrichosiphum tenuicorpus' OR 'Moniliopsis foliicola' OR 'Monochaetia alnea' OR 'Monosoma pulverata' OR 'Monosteira unicostata' OR 'Monostichella alni' OR 'Monosynamma bohemanni' OR 'Monsoma pulveratum' OR 'Morimus asper' OR 'Mormo maura' OR 'Morrisonia latex' OR 'Mycena abramsii' OR 'Mycena algeriensis' OR 'Mycena alnicola' OR 'Mycena citrinomarginata' OR 'Mycena galericulata' OR 'Mycena galericulata' OR 'Mycena haematopus' OR 'Mycena leaiana' OR 'Mycena longiseta' OR 'Mycena maculata' OR 'Mycena rhenana' OR 'Mycena viridimarginata' OR 'Mycenella margaritispora' OR 'Mycoacia aurea' OR 'Mycoacia fuscoatra' OR 'Mycoacia gilvescens' OR 'Mycoacia uda' OR 'Mycoaciella bispora' OR 'Mycobernardia incrustans' OR 'Mycocalia denudata' OR 'Mycoglaena alni' OR 'Mycoleptodon dichroum' OR 'Mycomicrothelia confusa' OR 'Mycopappus alni' OR 'Mycosphaerella alnicola' OR 'Mycosphaerella alni-viridis' OR 'Mycosphaerella conglomerata' OR 'Mycosphaerella incompta' OR 'Mycosphaerella latebrosa' OR 'Mycosphaerella maculiformis' OR 'Mycosphaerella perparva' OR 'Mycosphaerella punctiformis' OR 'Mycterothrips consociatus' OR 'Mycterothrips latus' OR 'Mycterothrips salicis' OR 'Myelopsis minutularia' OR 'Myrmaecium rubricosum' OR 'Mytilodiscus alnicola' OR 'Myxofusicoccum alni' OR 'Myxosporium roumeguerei' OR 'Nadata gibbosa' OR 'Naematelia aurantia' OR 'Naenia typica' OR 'Natantiella ligneola' OR 'Nathrius brevipennis' OR 'Nealgedonia extricalis' OR 'Nealgedonia extricalis dionalis' OR 'Nectria cinnabrina' OR 'Nectria lugdunensis' OR 'Necydalis major' OR 'Nellymyces megaceros' OR 'Nemania diffusa' OR 'Nemania serpens' OR 'Nematinus abdominalis' OR 'Nematinus acuminatus' OR 'Nematinus bilineatus' OR 'Nematinus fuscipennis' OR 'Nematinus luteus' OR 'Nematinus steini' OR 'Nematinus willigkiae' OR 'Nematocampa filamentaria' OR 'Nematogonum ferrugineum' OR 'Nematus alniastri' OR 'Nematus capreae' OR 'Nematus latipes' OR 'Nematus oligospilus' OR 'Nematus pavidus' OR 'Nematus polypilus' OR 'Nematus septentrionalis' OR 'Nematus umbratus' OR 'Nemoria mimosaria' OR 'Neobetulaphis chaetosiphon' OR 'Neobetulaphis pusilla' OR 'Neobulgaria pura' OR 'Neocrepidodera peirolierii' OR 'Neocurbitaria rhamni' OR 'Neodasyscypha cerina' OR 'Neofusicoccum ribis' OR 'Neohelicosporium griseum' OR 'Neolyngus contaminatus' OR 'Neolyngus viridis' OR 'Neonectria coccinea' OR 'Neonectria ditissima' OR 'Neonectria major' OR 'Neonectria punicea' OR 'Neopolvinaria innumerabilis innumerabilis' OR 'Neottiella vitellina' OR 'Neozephyrus helenae' OR 'Neozephyrus japonica' OR 'Neozephyrus taiwanus' OR 'Neozephyrus taxila japonicus' OR 'Neurospora sitophila' OR 'Neurospora tetraspora' OR 'Nipaecoccus vectis' OR 'Nipterella parksii' OR 'Nites betulella' OR 'Nites grotella' OR 'Nitschka grevillii' OR 'Nivellia sanguinosa' OR 'Nola clethrae' OR 'Nola confusalis' OR 'Nola minna' OR 'Notodonta dromedarius' OR 'Notodonta stigmatica' OR 'Notodonta ziczac' OR 'Nymphalis antiopa' OR 'Nymphalis vaualbum' OR 'Oberea linearis' OR 'Occultocarpon ailaoshanense' OR 'Ocellaria aurantiaca' OR 'Ochropacha duplaris' OR 'Octospora sydowii' OR 'Odonestis pruni' OR 'Odontia calcicola' OR 'Odonticium septocystidia' OR 'Odontopera bidentata' OR 'Odontoplatys bidentulus' OR 'Odontosia carmelita' OR 'Odontosia sieversii' OR 'Oemona hirta' OR 'Oestlundia flava' OR 'Ohleria rugulosa' OR 'Oidium betulacearum' OR 'Olethreutes albiciania' OR 'Olethreutes appendiceum' OR 'Olethreutes brunneopurpuratum' OR 'Olethreutes fraternana' OR 'Olethreutes submissana' OR 'Oligia illocata' OR 'Oligocentria pallida' OR 'Oligocentria semirufescens' OR 'Oligonychus biharensis' OR 'Oligonychus coffeae' OR 'Oligonychus mcgregori' OR 'Oligonychus perseae' OR 'Oligonychus punicae' OR 'Oligonychus ununguis' OR 'Oliveonia pauxilla' OR 'Ombrophila janthina' OR 'Oncopodiella robusta' OR 'Oncopodiella trigonella' OR 'Oncopsis alni' OR 'Oncopsis flavicollis' OR 'Oncopsis planiuscula' OR 'Onnia tomentosa' OR 'Operophtera fagata' OR 'Operophtera occidentalis' OR 'Ophiognomonia alni-cordatae' OR 'Ophiognomonia alni-viridis' OR 'Ophiognomonia apiospora' OR 'Ophiognomonia bugabensis' OR 'Ophiognomonia gardienneti' OR 'Ophiognomonia ibarakiensis' OR 'Ophiognomonia intermedia' OR 'Ophiognomonia ischnostyla' OR 'Ophiognomonia michiganensis' OR 'Ophiognomonia multirostrata' OR 'Ophiognomonia naganoensis' OR 'Ophiognomonia pseudoischnostyla' OR 'Ophiognomonia setacea' OR 'Ophiognomonia trientensis' OR 'Ophiognomonia tucumanensis' OR 'Ophiostoma pseudokarelicum' OR 'Ophiostoma signatum' OR 'Ophiostoma sparsianulatum' OR 'Orbilia crenatomarginata' OR 'Orbilia leucostigma' OR 'Orbilia pyrifera' OR 'Orbilia xanthostigma' OR 'Orchestes jota' OR 'Orchestes testaceus' OR 'Orgya antiqua' OR 'Orgya antiqua badia' OR 'Orgya antiqua nova' OR 'Orgya leucostigma' OR 'Orgya leucostigma intermedia' OR 'Orgya leucostigma plagiata' OR 'Orgya recens' OR 'Orgya vetusta' OR 'Orientus ishidae' OR 'Orthosia cerasi' OR 'Orthosia gothica' OR 'Orthosia hibisci' OR 'Orthosia incerta' OR 'Orthosia opima' OR 'Orthosia rubescens' OR 'Orthotaenia undulana' OR 'Orthotylus flavinervis' OR 'Orthotylus interpositus' OR 'Orthotylus marginalis' OR 'Ossiannilssonola callosa' OR 'Osteina obducta' OR 'Otiorhynchus apenninus' OR 'Otiorhynchus armadillo' OR 'Otiorhynchus aurifer' OR 'Otiorhynchus carinatopunctatus' OR 'Otiorhynchus coecus' OR 'Otiorhynchus desertus' OR 'Otiorhynchus fagi' OR 'Otiorhynchus fullo' OR 'Otiorhynchus morio' OR 'Otiorhynchus multipunctatus' OR 'Otiorhynchus pyrenaeus' OR 'Otiorhynchus scaber' OR 'Otiorhynchus singularis' OR 'Otiorhynchus tenebricosus' OR 'Otiorhynchus uncinatus' OR 'Ourapteryx obtusicauda' OR 'Ourapteryx sambucaria' OR 'Oxycarenus modestus' OR 'Oxymirus cursor' OR 'Oxyporus corticola' OR 'Oxyporus populinus' OR 'Oxyporus ravidus' OR 'Pachyella hydrophila' OR 'Pachytodes cerambyciformis' OR 'Palatinate grapevine yellows' OR 'Palomena prasina' OR 'Palthis angulalis' OR 'Pamphilus fumipennis' OR 'Pamphilus kontuniemii' OR 'Pamphilus nigrifemoratus' OR 'Pamphilus pallipes' OR 'Pamphilus vafer' OR 'Pamphilus varius' OR 'Pandemis canadana' OR 'Pandemis cerasana' OR 'Pandemis corylana' OR 'Pandemis dumetana' OR 'Pandemis heparana' OR 'Pandemis limitata' OR 'Panellus longinquus' OR 'Panellus oralis' OR 'Panellus ringens' OR 'Panellus ringens' OR 'Panellus stipticus' OR 'Panonychus ulmi' OR 'Pantilius tunicatus' OR 'Panus conchatus' OR 'Panus rufus' OR 'Papestra cristifera' OR 'Papestra invalida' OR 'Papestra quadrata' OR 'Papilio eurymedon' OR 'Papilio glaucus' OR 'Papilio glaucus canadensis' OR 'Papilio glaucus glaucus' OR 'Papilio rutulus' OR 'Paracolax tristalis' OR 'Paracrania chrysolepidella' OR 'Paradarisa consonaria' OR 'Paradarisa extersaria' OR 'Paradiarsia littoralis' OR 'Paralongidorus maximus' OR 'Paranthrene asilipennis' OR 'Paranthrene tabaniformis' OR 'Parasaissetia nigra' OR 'Parasyrphus nigritarsis' OR 'Paratrichodorus pachydermus' OR 'Paratrichodorus ramblensis' OR 'Paratrichodorus teres' OR 'Paratrichodorus tunisiensis' OR 'Paratylenchus hamatus' OR 'Paratylenchus macrophallus' OR 'Paratylenchus projectus' OR 'Paratylenchus straeleni' OR 'Paratylenchus veruculatus' OR 'Parectropis similaria' OR 'Parlatoria crypta' OR 'Parornix alni' OR 'Parthenolecanium corni' OR 'Passalora alni' OR 'Passalora bacilligera' OR 'Passalora microsperma' OR 'Passalora nepalensis' OR 'Patinella flavobrunnea' OR 'Pechipogo strigilata' OR 'Pellidiscus pallidus' OR 'Pelosia muscerala' OR 'Penicillago nodositata' OR 'Peniophora albobadia' OR 'Peniophora aurantiaca' OR 'Peniophora cinerea' OR 'Peniophora erikssonii' OR 'Peniophora incarnata' OR 'Peniophora pithya' OR 'Peniophora polygonia' OR 'Peniophora

(Continued)

pseudoversicolor' OR 'Peniophora quercina' OR 'Peniophora rhodochroa' OR 'Peniophora roumeguerii' OR 'Peniophora versiformis' OR 'Peniophora violaceolivida' OR 'Peniophorella guttulifera' OR 'Peniophorella praetermissa' OR 'Peniophorella pubera' OR 'Pentarthrum huttoni' OR 'Pentastiridius beieri' OR 'Pentatoma rufipes' OR 'Perconia strigillaria' OR 'Perenniporia amylocladina' OR 'Perenniporia tenuis' OR 'Perenniporia tenuis' OR 'Peridea gigantea' OR 'Peridroma saucia' OR 'Pero gigantea' OR 'Pero hubneraria' OR 'Pero morrisonaria' OR 'Peroettia flammea' OR 'Pertusaria carneopallida' OR 'Pertusaria pupillaris' OR 'Pestalotia alnea' OR 'Pestalotiopsis glandicola' OR 'Pestalotiopsis microspora' OR 'Peyronellaea obtusa' OR 'Pezicula alni' OR 'Pezicula alnicola' OR 'Pezicula cinnamomea' OR 'Pezicula frangulae' OR 'Pezicula heterochroma' OR 'Pezicula livida' OR 'Phaeoacremonium croatiense' OR 'Phaeoacremonium fraxinopennsylvanicum' OR 'Phaeoacremonium iranianum' OR 'Phaeoacremonium minimum' OR 'Phaeocalcium compressulum' OR 'Phaeohelotium nobile' OR 'Phaeoisaria sparsa' OR 'Phaeomarasmius erinaceus' OR 'Phaeophlebiopsis ravenelii' OR 'Phaeosaccardinula penzigi' OR 'Phaeosphaerella borealis' OR 'Phaeotremella frondosa' OR 'Phalera bucephala' OR 'Phanerochaete affinis' OR 'Phanerochaete burtii' OR 'Phanerochaete carnosa' OR 'Phanerochaete laevis' OR 'Phanerochaete sordida' OR 'Phanerochaete velutina' OR 'Phellinopsis conchata' OR 'Phellinus igniarius' OR 'Phellinus laevigatus' OR 'Phellinus lundellii' OR 'Phellinus prunicola' OR 'Phellinus viticola' OR 'Phenacoccus aceris' OR 'Pheosia gnoma' OR 'Phialocephala compacta' OR 'Phialophora verrucosa' OR 'Phigalia pilosaria' OR 'Phlebia albomellea' OR 'Phlebia cinnabarinia' OR 'Phlebia cystidiata' OR 'Phlebia ludoviciana' OR 'Phlebia radiata' OR 'Phlebia rufa' OR 'Phlebia tremellosa' OR 'Phleospora dearnessii' OR 'Phleospora borealis' OR 'Phlogophora meticulosa' OR 'Phlogophora periculosa' OR 'Phlyctaenia coronata tertiialis' OR 'Pholiota adiposa' OR 'Pholiota alniphila' OR 'Pholiota aurivella' OR 'Pholiota aurivelloides' OR 'Pholiota occidentalis var. occidentalis' OR 'Pholiota populnea' OR 'Pholiota terrestris' OR 'Phoma pomorum var. pomorum' OR 'Phomopsis rhodophila' OR 'Phragmoporthe conformis' OR 'Phragmoporthe ploettneriana' OR 'Phragmotrichum rivoclarinum' OR 'Phragmotrichum vassiljevae' OR 'Phyllactinia alni' OR 'Phyllactinia alnicola' OR 'Phyllactinia fraxini' OR 'Phyllactinia guttata' OR 'Phyllobius alpinus' OR 'Phyllobius arborator' OR 'Phyllobius argentatus' OR 'Phyllobius calcaratus' OR 'Phyllobius fessus' OR 'Phyllobius glaucus' OR 'Phyllobius maculicornis' OR 'Phyllobius pomaceus' OR 'Phyllobius pyri' OR 'Phyllobius viridicollis' OR 'Phyllocoptes alniborealis' OR 'Phyllocoptes alniincanae' OR 'Phyllocoptes punctatus' OR 'Phylloclada americana' OR 'Phyllonorycter alaskana' OR 'Phyllonorycter alni' OR 'Phyllonorycter alnicella' OR 'Phyllonorycter alnivorella' OR 'Phyllonorycter alpina' OR 'Phyllonorycter auronitens' OR 'Phyllonorycter chrysella' OR 'Phyllonorycter durangensis' OR 'Phyllonorycter faginella' OR 'Phyllonorycter froelichiella' OR 'Phyllonorycter groenlieni' OR 'Phyllonorycter hancola' OR 'Phyllonorycter incanella' OR 'Phyllonorycter kisoensis' OR 'Phyllonorycter klemannella' OR 'Phyllonorycter longispinata' OR 'Phyllonorycter maculata' OR 'Phyllonorycter messaniella' OR 'Phyllonorycter nepalensis' OR 'Phyllonorycter populifoliella' OR 'Phyllonorycter rajella' OR 'Phyllonorycter stettinensis' OR 'Phyllonorycter strigulatella' OR 'Phyllonorycter suaveolentis' OR 'Phyllonorycter takagii' OR 'Phyllonorycter tristrigella' OR 'Phyllonorycter vulturella' OR 'Phyllosticta allantospora' OR 'Phyllosticta alnea' OR 'Phyllosticta alnigena' OR 'Phyllosticta alni-glutinosae' OR 'Phyllosticta alniperda' OR 'Phyllosticta capitalensis' OR 'Phyllosticta carpini' OR 'Phyllosticta frangulae' OR 'Phyllostopsis nidulans' OR 'Phylus coryli' OR 'Phylus plagiatus' OR 'Phymatodes alni' OR 'Phymatotrichopsis omnivore' OR 'Physalospora abdita' OR 'Physarum leucophaeum' OR 'Physarum nutans' OR 'Physatocheila costata' OR 'Physisorinus vitreus' OR 'Phytobia cambii' OR 'Phytobia carbonaria' OR 'Phytocoris longipennis' OR 'Phytocoris populi' OR 'Phytocoris reuteri' OR 'Phytophthora acerina' OR 'Phytophthora alni' OR 'Phytophthora alni subsp. alni' OR 'Phytophthora amnicola' OR 'Phytophthora asparagi' OR 'Phytophthora bilorbang' OR 'Phytophthora cactorum' OR 'Phytophthora cambivora' OR 'Phytophthora chlamydospora' OR 'Phytophthora cinnamomi' OR 'Phytophthora citricola' OR 'Phytophthora crassamira' OR 'Phytophthora cryptogea' OR 'Phytophthora europaea' OR 'Phytophthora gallica' OR 'Phytophthora gonapodyides' OR 'Phytophthora gregata' OR 'Phytophthora inundata' OR 'Phytophthora lacustris' OR 'Phytophthora megasperma' OR 'Phytophthora multivora' OR 'Phytophthora parsiana' OR 'Phytophthora plurivora' OR 'Phytophthora polonica' OR 'Phytophthora pseudocryptogea' OR 'Phytophthora pseudosyringae' OR 'Phytophthora quercina' OR 'Phytophthora rosacearum' OR 'Phytophthora siskiyouensis' OR 'Phytophthora syringae' OR 'Phytoptus laevis' OR 'Phytoptus nalepai' OR 'Picipes badius' OR 'Picipes melanopus' OR 'Picipes tubaeformis' OR 'Pilidium lythri' OR 'Pirex concentricus' OR 'Plagiognathus arbustorum' OR 'Plagiosterna aenea' OR 'Plagiostoma jenseii' OR 'Plagiostoma salicellum' OR 'Plagiostoma samuelsii' OR 'Plagodis alcoolaria' OR 'Plagodis phlogosaria' OR 'Plagodis phlogosaria approximaria' OR 'Plagodis pulveraria' OR 'Planstromella conglomeratiformis' OR 'Platarcia parthenos' OR 'Platycampus luridiventris' OR 'Platychora alni' OR 'Platynota nigrocervina' OR 'Platypus cylindrus' OR 'Platypus simulans' OR 'Plemyria georgii' OR 'Plemyria rubiginata' OR 'Pleohelicon richonis' OR 'Pleomassaria holoschista' OR 'Pleospora alnea' OR 'Pleospora alnicola' OR 'Pleospora pygmaea' OR 'Pleuroflammula tuberculosa' OR 'Pleurophragmium varieseptatum' OR 'Pleurotus cornucopiae' OR 'Pleurotus limpidus' OR 'Pleurotus ostreatus' OR 'Plicatura nivea' OR 'Plicaturopsis crispa' OR 'Pluteus latifolius' OR 'Pluteus nanus' OR 'Pluteus romellii' OR 'Pluteus semibulbosus' OR 'Pochazia shantungensis' OR 'Pococera aplastella' OR 'Podofomes mollis' OR 'Podofomes stereoides' OR 'Podosphaera clandestina' OR 'Poecilium alni' OR 'Poecilocampa populi' OR 'Pogonocherus hispidulus' OR 'Pogonocherus hispidus' OR 'Polia bombycina' OR 'Polia imbrifera' OR 'Polia nimbosa' OR 'Polia purpurissata' OR 'Polydrusus amoenus' OR 'Polydrusus cervinus' OR 'Polydrusus corruscus' OR 'Polydrusus flavipes' OR 'Polydrusus formosus' OR 'Polydrusus fulvicornis' OR 'Polydrusus impressifrons' OR 'Polydrusus picus' OR 'Polydrusus pilosulus' OR 'Polydrusus pilosus' OR 'Polydrusus prasinus' OR 'Polydrusus pterygomalis' OR 'Polydrusus sericeus' OR 'Polydrusus sparsus' OR 'Polydrusus tereticollis' OR 'Polydrusus undatus' OR 'Polygonia faunus' OR 'Polygonia faunus faunus' OR 'Polygonia oreas' OR 'Polygonia satyrus' OR 'Polyozellus humicola' OR 'Polyporus lepideus' OR 'Polyporus pargamensis' OR 'Polyporus picipes' OR 'Popillia japonica' OR 'Poria conferta' OR 'Poria tenuis var. tenuis' OR 'Poriella subacida' OR 'Porodaedalea pinii' OR 'Porophilomyces poricola' OR 'Porostereum spadiceum' OR 'Porothelium fimbriatum' OR 'Postia immitis' OR 'Postia tephroleuca' OR 'Praetumpfia obducens' OR 'Pratylenchus crenatus' OR 'Pratylenchus flakkensis' OR 'Pratylenchus neglectus' OR 'Pratylenchus penetrans' OR 'Pratylenchus vulnus' OR 'Prinobius myardi' OR 'Prionus coriarius' OR 'Probolus amicaria' OR 'Prochoerodes transversata' OR 'Prociphilus baicalensis' OR 'Prociphilus mexicanus' OR 'Prociphilus tessellatus' OR 'Propolis farinosa' OR 'Prosthemium alni' OR 'Prosthemium stellare' OR 'Protantigius superans' OR 'Protantigius superans ginzii' OR 'Protitame matilda' OR 'Protitame virginalis' OR 'Protodiaspis parvula' OR 'Psallus ambiguus' OR 'Psallus betuleti' OR 'Psallus haematoxylon' OR 'Psallus henschi' OR 'Psallus perrisi' OR 'Psallus salicis' OR 'Psallus variabilis' OR 'Psallus varians' OR 'Psathyrella alboalutacea' OR 'Psathyrella candidissima' OR 'Psathyrella griseifolia' OR 'Psathyrella immaculata' OR 'Psathyrella maculata' OR 'Psepholax sulcatus' OR 'Pseudaonidia duplex' OR 'Pseudaulacaspis celtis' OR 'Pseudaulacaspis prunicola prunicola' OR 'Pseuderannis lomozemias' OR 'Pseudociboria umbrina' OR 'Pseudococcus comstocki' OR 'Pseudocosmospora pitoides' OR

(Continues)

(Continued)

'Pseudoglaea olivata' OR 'Pseudoips praninana' OR 'Pseudoloxops coccineus' OR 'Pseudomonas syringae' OR 'Pseudopityophthorus pruinosis' OR 'Pseudorthodes irrorata' OR 'Pseudosciaphila duplex' OR 'Pseudosigmoidea alnicola' OR 'Pseudotelphusa belangerella' OR 'Pseudothyatira cymatophoroides' OR 'Pseudotrichia mutabilis' OR 'Pseudovalsaria ferruginea' OR 'Pseudovalsella thelebola' OR 'Psyche casta' OR 'Psyche crassiorella' OR 'Psyche rotunda' OR 'Psylla alni' OR 'Psylla alpina' OR 'Psylla cordata' OR 'Psylla foersteri' OR 'Psylla fusca' OR 'Pterocallis albida' OR 'Pterocallis alni' OR 'Pterocallis alnifoliae' OR 'Pterocallis alnijaponicae' OR 'Pterocallis essigi' OR 'Pterocallis maculata' OR 'Pterocallis nigrostriata' OR 'Pterocallis pseudoalni' OR 'Pterocallis rhombifoliae' OR 'Pterostoma palpina' OR 'Ptilinus pectinicornis' OR 'Ptilodon capucina' OR 'Ptilodon robusta' OR 'Puccinia coronata' OR 'Pulvinaria borchsenii' OR 'Pulvinaria costata' OR 'Pulvinaria idesiae' OR 'Pulvinaria inconspecta' OR 'Pulvinaria occidentalis' OR 'Pulvinaria regalis' OR 'Pulvinaria vitis' OR 'Punctularia strigosozonata' OR 'Pycnopeziza americana' OR 'Pyrenopeziza benesuada' OR 'Pyrenopeziza benesuada' OR 'Pyrenopeziza foliocola' OR 'Pyrigemmula aurantiaca' OR 'Pyrrhia umbra' OR 'Pyrrhoglossum recedens' OR 'Racodium therryanum' OR 'Radulomyces confluens' OR 'Radulomyces molaris' OR 'Ramaricium albo-ochraceum' OR 'Ramichloridium anceps' OR 'Ramphus pulicarius' OR 'Ramularia alnicola' OR 'Ramularia iwayamensis' OR 'Ramularia taleshina' OR 'Ramularia unterseheri' OR 'Ranullospora alni' OR 'Raphia frater' OR 'Rectipilus fasciculatus' OR 'Refractohilum achromaticum' OR 'Repetophragma wroblewskii' OR 'Resinicium bicolor' OR 'Resinicium praeteritum' OR 'Resinomyces saccharifera' OR 'Resinoporia crassa' OR 'Resupinatus conglobatus' OR 'Reticularia lycoperdon' OR 'Rhabdospora maculans' OR 'Rhagium bifasciatum' OR 'Rhagium mordax' OR 'Rhagium sycophanta' OR 'Raphigaster nebulosa' OR 'Rheumaptera hastata' OR 'Rheumaptera subhastata albodecorata' OR 'Rhizina undulata' OR 'Rhizobium rhizogenes' OR 'Rhizoctonia ochracea' OR 'Rhizoctonia pseudocornigera' OR 'Rhizoctonia solani' OR 'Rhizoctonia stridii' OR 'Rhizonemella sequoiae' OR 'Rhodofomes cajanderi' OR 'Rhogogaster chlorosoma' OR 'Rhogogaster punctulata' OR 'Rhogogaster scalaris' OR 'Rhogogaster viridis' OR 'Rhynchaenus alni' OR 'Rhynchaenus iota' OR 'Rhynchaenus stigma' OR 'Rhynchaenus testaceus' OR 'Rhynchites nanus' OR 'Rhynchites tomentosus' OR 'Rhytidodus decimusquartus' OR 'Rhytisma salicinum' OR 'Ribautiana cruciata' OR 'Ribautiana debilis' OR 'Ribautiana tenerima' OR 'Ribautiana ulmi' OR 'Rigidoporus microporus' OR 'Rigidoporus undatus' OR 'Ropalopus clavipes' OR 'Ropalopus femoratus' OR 'Ropalopus ungaricus' OR 'Rosalia alpina' OR 'Rosellinia abscondita' OR 'Rosellinia aquila' OR 'Rosellinia callosa' OR 'Rosellinia corticum' OR 'Rosellinia desmazieri' OR 'Rosellinia helvetica' OR 'Rosellinia marcucciana' OR 'Rosellinia necrotioides' OR 'Rosellinia thelena' OR 'Rotylenchus buxophilus' OR 'Rutpela maculata' OR 'Rutstroemia bolaris' OR 'Rutstroemia conformata' OR 'Ruzenia spermoides' OR 'Sabra harpagula' OR 'Sabulodes aegrotata' OR 'Sabulodes caberata' OR 'Saccothecium sepincula' OR 'Salebriopsis albicilla' OR 'Saperda carcharias' OR 'Saperda octopunctata' OR 'Saperda populnea' OR 'Saperda scalaris' OR 'Saphanus piceus' OR 'Sarcodontia electans' OR 'Sarcomyxa serotina' OR 'Sarcoscypha austriaca' OR 'Sarocladium strictum' OR 'Saturnia atlantica' OR 'Saturnia pavonia' OR 'Saturnia pyri' OR 'Satyrium w-album' OR 'Schizophyllum amplum' OR 'Schizophyllum commune' OR 'Schizopora paradoxa' OR 'Schizotetranychus alni' OR 'Schizothryium jamaicense' OR 'Schizura concinna' OR 'Schizura ipomoeae' OR 'Schizura unicornis' OR 'Sclerococcum stygium' OR 'Sclerotium nervale' OR 'Scolioneura betuleti' OR 'Scoliopteryx libatrix' OR 'Scolytus intricatus' OR 'Scolytus rugulosus' OR 'Scopinella caulincola' OR 'Scopulariopsis asperula' OR 'Scopulariopsis brumptii' OR 'Scopuloides rimosus' OR 'Scorias spongiosa' OR 'Scutellinia hirta' OR 'Scutellinia scutellata' OR 'Scytalidium lignicola' OR 'Scytinostroma portentosum' OR 'Scytinostroma protrusum subsp. protrusum' OR 'Sebacina burtii' OR 'Sebacina epigaea var. epigaea' OR 'Sebacina grisea' OR 'Sebacina incrustans' OR 'Sebacina macrospora' OR 'Seimatosporium alneum' OR 'Selatosomus bipustulatus' OR 'Selenia alciphearia' OR 'Selenia dentaria' OR 'Selenia lunularia' OR 'Selenia tetralunaria' OR 'Semiorthisa aemulataria' OR 'Semiorthisa alternaria' OR 'Semiorthisa granitata-group' OR 'Semiorthisa hebetata' OR 'Semiorthisa neptaria' OR 'Semiorthisa ulsterata' OR 'Septobasidium aligerum' OR 'Septobasidium bogoriense' OR 'Septonema secedens' OR 'Septonema subramosum' OR 'Septoria alni' OR 'Septoria alnifolia' OR 'Septoria frangulae' OR 'Septoria taleshana' OR 'Septoria weiri' OR 'Septotrullula bacilligera' OR 'Sertulicium niveocremeum' OR 'Sesia apiformis' OR 'Setagrotis pallidicollis' OR 'Setagrotis planifrons' OR 'Sicya crocea' OR 'Sicya macularia' OR 'Sidera lenis' OR 'Sierraphytopus alnivagrans' OR 'Sinodendron cylindricum' OR 'Sirodthis inversa' OR 'Sistotrema brinkmannii' OR 'Sistotrema brinkmannii' OR 'Sistotrema farinaceum' OR 'Sistotrema porulosum' OR 'Skeletocutis alutacea' OR 'Skeletocutis semipileata' OR 'Skeletocutis subincarnata' OR 'Smerinthus ocellata' OR 'Soltanea mariae' OR 'Spadicoides atra' OR 'Spadicoides bina' OR 'Spadicoides klotzschii' OR 'Sparganothis pettitana' OR 'Sparganothis reticulatana' OR 'Speudotettix subfuscus' OR 'Sphaeloma alni' OR 'Sphaerobolus stellatus' OR 'Sphaeronema alni' OR 'Sphinx gordius' OR 'Sphinx luscitiosa' OR 'Spilonota ocellana' OR 'Spilosoma lutea' OR 'Spilosoma virginica' OR 'Spiramater grandis' OR 'Spiramater lutra' OR 'Spodoptera ornithogalli' OR 'Spongiporus perdelicatus' OR 'Sporendodladia fumosa' OR 'Sporidesmium folliculatum' OR 'Stachybotrys echinatus' OR 'Stanjehughesia hormiscioides' OR 'Stathmopoda pedella' OR 'Stauropus fagi' OR 'Steccherinum alaskense' OR 'Steccherinum bourdotii' OR 'Steccherinum fimbriatellum' OR 'Steccherinum fimbriatum' OR 'Steccherinum fimbriatum' OR 'Steccherinum laeticolor' OR 'Steccherinum ochraceum' OR 'Steccherinum ochraceum' OR 'Stegania cararia' OR 'Stenocephalopsis subalutacea' OR 'Stenocorus meridianus' OR 'Stenocybe pullatula' OR 'Stenopterus flavidicornis' OR 'Stenopterus rufus' OR 'Stephanitis pyri' OR 'Stereum complicatum' OR 'Stereum frustulatum' OR 'Stereum gausapatum' OR 'Stereum hirsutum' OR 'Stereum ochraceoflavum' OR 'Stereum ostrea' OR 'Stereum rugosum' OR 'Stereum sanguinolentum' OR 'Stereum subtomentosum' OR 'Sterrhopterix standfussi' OR 'Sthenarus rotermundi' OR 'Sthenopis argenteomaculatus' OR 'Stictoleptura scutellata' OR 'Stigmella alnetella' OR 'Stigmella canadensis' OR 'Stigmella confusella' OR 'Stigmella continua' OR 'Stigmella glutinosae' OR 'Stigmella laponica' OR 'Stigmella luteella' OR 'Stigmella marginicolella' OR 'Stigmella microtheriella' OR 'Stigmella rubescens' OR 'Stilbella byssisida' OR 'Stilbella clavispora' OR 'Stomaphis alni' OR 'Stomaphis quercus' OR 'Stomaphis radicicola' OR 'Stomaphis wojciechowski' OR 'Strangalia attenuata' OR 'Strangalia aurulenta' OR 'Strangalia quadrifasciata' OR 'Stromatium auratum' OR 'Strossmayeria alnicala' OR 'Strossmayeria atriseta' OR 'Strossmayeria bakeriana' OR 'Stygnocoris sabulosus' OR 'Subacronicta megacephala' OR 'Subulicystidium longisporum' OR 'Svrcekomyces pallidus' OR 'Symphytocarpus flaccidus' OR 'Symydobius aliarius' OR 'Symydobius aliarius ssp. Nipponicus' OR 'Symydobius kabae' OR 'Symydobius minutus' OR 'Symydobius oblongus' OR 'Symydobius quednau' OR 'Synanthoned culiciformis' OR 'Synanthoned mesiaeformis' OR 'Synanthoned multitarsus' OR 'Synanthoned pseudoscoliaeforme' OR 'Synanthoned spheciiformis' OR 'Synanthoned talischense' OR 'Synanthoned tenuis' OR 'Synaxis jubararia' OR 'Syndemis afflictana' OR 'Syngrapha epigaea' OR 'Szczepkamyces campestris' OR 'Tachyerges pseudostigma' OR 'Tachyerges stigma' OR 'Tacparia detersa' OR 'Taeniolella alta' OR 'Taeniolella stilbospora' OR 'Taeniolella stricta' OR 'Taeniolina scripta' OR 'Takahashia japonica' OR 'Taoia chuansiensis' OR 'Taoia indica' OR 'Tapesia fusca' OR 'Tapesia lividofusca' OR 'Tapesia mollisioides' OR 'Tapesia

(Continued)

villosa' OR 'Taphrina alni' OR 'Taphrina epiphylla' OR 'Taphrina japonica' OR 'Taphrina macrophylla' OR 'Taphrina occidentalis' OR 'Taphrina populina' OR 'Taphrina robinsoniana' OR 'Taphrina rugosa' OR 'Taphrina sadebeckii' OR 'Taphrina tosquinetii' OR 'Taphrina viridis' OR 'Taphrorychus siculos' OR 'Taphrorychus villifrons' OR 'Tectella patellaris' OR 'Tegonotus borealis' OR 'Tegonotus heptacanthus' OR 'Tegonotus keiferi' OR 'Tegonotus platynaspis' OR 'Tegonotus trouessartii' OR 'Teichospora winteriana' OR 'Teleiodes proximella' OR 'Temnocerus coeruleus' OR 'Temnocerus nanus' OR 'Tenthredo ferruginea' OR 'Tenthredo velox' OR 'Tenuiappendicula alnica' OR 'Tetheella fluctuosa' OR 'Tetrachaetum elegans' OR 'Tetracis cachexiata' OR 'Tetracis crocallata' OR 'Tetranychus urticae' OR 'Tetranychopsis horridus' OR 'Thallophaga hyperborea' OR 'Thecotheus rivicola' OR 'Thelephora atra' OR 'Thelephora ellisia' OR 'Thelephora wakefieldiae' OR 'Thrips alni' OR 'Thrips calcaratus' OR 'Thrips major' OR 'Thrips minutissimus' OR 'Thrips viminalis' OR 'Thyridaria macrostomoides' OR 'Thyridium vestitum' OR 'Thyronectria coryli' OR 'Tinocallis ulmicola' OR 'Tinocallis zelkovae' OR 'Tomasellia diffusa' OR 'Tomentella badia' OR 'Tomentella bryophila' OR 'Tomentella cinerascens' OR 'Tomentella coerulea' OR 'Tomentella crinalis' OR 'Tomentella donkii' OR 'Tomentella ferruginea' OR 'Tomentella fuscocinerea' OR 'Tomentella lapida' OR 'Tomentella lilacinogrisea' OR 'Tomentella puberula' OR 'Tomentella punicea' OR 'Tomentella viridis' OR 'Tortilispora aurantiaca' OR 'Tortricidia testacea' OR 'Tortrix viridana' OR 'Torula herbarum' OR 'Torula lucifuga' OR 'Trametes cinnabarin' OR 'Trametes coccinea' OR 'Trametes cubensis' OR 'Trametes gibbosa' OR 'Trametes hirsuta' OR 'Trametes lacerata' OR 'Trametes pubescens' OR 'Trametes vernipes' OR 'Trametes versicolor' OR 'Treichispora candidissima' OR 'Treichispora farinacea' OR 'Treichispora mollusca' OR 'Trematosphaeria pertusa' OR 'Tremella mesenterica' OR 'Tremex fuscicornis' OR 'Trichaptum abietinum' OR 'Trichaptum biforme' OR 'Trichaptum byssogenum' OR 'Trichaptum laricinum' OR 'Trichiosoma lucorum' OR 'Trichiosoma vitellina' OR 'Trichiura crataegi' OR 'Trichius fasciatus' OR 'Trichocladium asperum' OR 'Trichoderma alni' OR 'Trichoderma brunneoviride' OR 'Trichoderma citrinum' OR 'Trichoderma crystallinum' OR 'Trichoderma estonicum' OR 'Trichoderma flavipes' OR 'Trichoderma lixii' OR 'Trichoderma patella' OR 'Trichoderma strictipile' OR 'Trichoderma thelephoricola' OR 'Trichoderma viride' OR 'Trichoderma voglmayri' OR 'Trichodorus californicus' OR 'Trichodorus giennensis' OR 'Trichodorus sparsus' OR 'Trichoferus campestris' OR 'Trichoferus holosericeus' OR 'Tricholomopsis streetsii' OR 'Trichopteryx carpinata' OR 'Trichosphaeria pilosa' OR 'Trichothecium roseum' OR 'Tricladium angulatum' OR 'Tricladium splendens' OR 'Trichonymus thulensis' OR 'Triposporium pannosum' OR 'Trirachys sartus' OR 'Tritomegas bicolor' OR 'Truncatella angustata' OR 'Tryptodendron domesticum' OR 'Tryptodendron signatum' OR 'Tryptophloeus alni' OR 'Tryptophloeus asperatus' OR 'Tubaria furfuracea' OR 'Tubulicrinis glebulosus' OR 'Tulasnella bifrons' OR 'Tulasnella pallida' OR 'Tulasnella violea' OR 'Tydeus californicus' OR 'Tydeus caudatus' OR 'Tylenchorhynchus dubius' OR 'Tylenchus davainei' OR 'Tylenchus vulgaris' OR 'Tympanis alnea' OR 'Tympanis pseudoalnea' OR 'Tympanis truncatula' OR 'Typhlocyba quercus' OR 'Typhula contorta' OR 'Typhula erythropus' OR 'Typhula fistulosa' OR 'Tyromyces chioneus' OR 'Tyromyces galactinus' OR 'Ulmicola spinipes' OR 'Umbelopsis vinacea' OR 'Uncinula miyabei var. alnica' OR 'Uncinula miyabei var. hermaphroditica' OR 'Unguiculella foliicola' OR 'Valsa americana' OR 'Valsa ceratophora' OR 'Valsa diatrypoides' OR 'Valsa frangulae' OR 'Valsa inconspicua' OR 'Valsa minutella' OR 'Valsa salicina' OR 'Valsa truncata' OR 'Valsalnicola oxystoma' OR 'Valsaria moroides' OR 'Valsella alnica' OR 'Valsella furva' OR 'Vararia investiens' OR 'Varicosporium elodeae' OR 'Veluticeps abietina' OR 'Venturia alnea' OR 'Venturia ditricha' OR 'Venusia cambrica' OR 'Venusia comptaria' OR 'Venusia pearsallii' OR 'Verrucaria hydrula' OR 'Vibrissea filisporia' OR 'Vibrissea truncorum' OR 'Violella fucata' OR 'Vitreoporus dichrous' OR 'Volucrispora aurantiaca' OR 'Vuilleminia alni' OR 'Vuilleminia comedens' OR 'Watsonalla binaria' OR 'Watsonalla uncinula' OR 'Wuestneia paucispora' OR 'Xanthoporia radiata' OR 'Xanthorhoe defensaria' OR 'Xanthorhoe fluctuata' OR 'Xanthorhoe spadicearia' OR 'Xanthotype urticaria' OR 'Xenasma rimicola' OR 'Xenasmatella vaga' OR 'Xestia infimatis' OR 'Xestia smithii' OR 'Xestobium rufovillosum' OR 'Xiphinema globosum' OR 'Xiphinema pachtaicum' OR 'Xiphinema pyrenaicum' OR 'Xiphydria camelus' OR 'Xiphydria longicollis' OR 'Xiphydria megapolitana' OR 'Xiphydria picta' OR 'Xiphydria prolongata' OR 'Xylaria corniformis' OR 'Xylaria cornu-damae' OR 'Xylaria digitata' OR 'Xylaria hypoxylon' OR 'Xylaria polymorpha' OR 'Xylaria subterranea' OR 'Xyleborinus alni' OR 'Xyleborinus attenuatus' OR 'Xyleborinus saxesenii' OR 'Xyleborus dispar' OR 'Xyleborus pfeili' OR 'Xyleborus saxeseni' OR 'Xylena cineritia' OR 'Xylena curvimacula' OR 'Xylena exsoleta' OR 'Xylena nupera' OR 'Xylena thoracica' OR 'Xyllococcus betulae' OR 'Xylodoccus japonicus' OR 'Xylodon asper' OR 'Xylodon brevisetus' OR 'Xylodon flavigiporus' OR 'Xylodon radula' OR 'Xylodon rimosissimus' OR 'Xylomelasma sordida' OR 'Xylosandrus crassiusculus' OR 'Xylosandrus germanus' OR 'Xylosphaera bertero' OR 'Xyloteras domesticum' OR 'Xyloteras signatum' OR 'Xylotrechus ibex' OR 'Xylotrechus namanganensis' OR 'Xylotrechus rusticus' OR 'Xylotrechus stebbingi' OR 'Xylotype arcadia' OR 'Ypsolopha parenthesella' OR 'Zale minerea' OR 'Zale minerea norda' OR 'Zalerion maritima' OR 'Zethenia albonotaria' OR 'Zeuzera pyrina' OR 'Zonocryba bifasciata' OR 'Zosteropoda hirtipes' OR 'Zygina angusta' OR 'Zygina flammigera' OR 'Zygina rhamni' OR 'Zygina suavis' OR 'Zygina tiliae'

APPENDIX C

List of pests that can potentially cause an effect not further assessed

No.	Pest name	EPPO code	Group	Pest present in the UK	Pest present in the EU	<i>Alnus</i> confirmed as a host (reference)	Pest can be associated with the commodity (NA = not assessed)	Impact	Justification for inclusion in this list
1	<i>Eriophyes axillaris</i>	ERPHSI	Acari	Yes	No	<i>A. glutinosa</i> (Database of Insects and their Food Plants, Online)	Yes	Uncertain	No information on impact
2	<i>Melampsoridium alni</i>	MELMLI	Fungi	Uncertain	Limited	<i>Alnus</i> spp. (USDA Fungal Database)	Yes	Uncertain	Uncertainty about presence in UK (one record from 1961)
3	<i>Pestalotiopsis microspora</i>	PESTDC	Fungi	Uncertain	Limited	<i>A. rubra</i> (USDA Fungal Database)	Yes	Yes	Uncertainty on the presence in the UK (only one record in GBIF)
4	<i>Septoria alnicola</i>	SEPTAP	Fungi	Yes	Limited	<i>A. glutinosa</i> , <i>Alnus</i> sp. (USDA Fungal Database)	Yes	Uncertain	No information on impact of this species

APPENDIX D

Excel file with the pest list of *Alnus* species

Appendix D can be found in the online version of this output in the 'supporting information section'.